

Asymptotic stability of a nonlinear Korteweg-de Vries equation with a critical length

Jixun Chu,^{*} Jean-Michel Coron,[†] Peipei Shang[‡]

Abstract

We study an initial-boundary-value problem of a nonlinear Korteweg-de Vries equation posed on a finite interval $(0, 2\pi)$. The whole system has Dirichlet boundary condition at the left end-point, and both of Dirichlet and Neumann homogeneous boundary conditions at the right end-point. It is known that the origin is not asymptotically stable for the linearized system around the origin. We prove that the origin is (locally) asymptotically stable for the nonlinear system.

Key words: nonlinearity, Korteweg-de Vries equation, stability, center manifold

2000 MR Subject Classification: 35Q53, 35B35

1 Introduction

This article is concerned with the following initial-boundary-value problem of the Korteweg-de Vries (KdV) equation posed on a finite interval

$$\begin{cases} y_t + y_x + yy_x + y_{xxx} = 0, \\ y(t, 0) = y(t, L) = 0, \\ y_x(t, L) = 0, \\ y(0, x) = y_0 \in L^2(0, L), \end{cases} \quad (1.1)$$

^{*}Department of Applied Mathematics, School of Mathematics and Physics, University of Science and Technology Beijing, Beijing 100083, China, Université Pierre et Marie Curie-Paris 6, UMR 7598 Laboratoire Jacques-Louis Lions, 75005 Paris, France. E-mail: chujixun@mail.bnu.edu.cn. JXC was supported by the ERC advanced grant 266907 (CPDENL) of the 7th Research Framework Programme (FP7).

[†]Institut universitaire de France and Université Pierre et Marie Curie-Paris 6, UMR 7598 Laboratoire Jacques-Louis Lions, 75005 Paris, France. E-mail: coron@ann.jussieu.fr. JMC was supported by the ERC advanced grant 266907 (CPDENL) of the 7th Research Framework Programme (FP7).

[‡]Department of Mathematics, Tongji University, Shanghai 200092, China. E-mail: peipeishang@hotmail.com. PS was partially supported by the ERC advanced grant 266907 (CPDENL) of the 7th Research Framework Programme (FP7).

with $L = 2\pi$.

The KdV equation was first derived by Boussinesq in [4] (see, in particular, equation (283 bis), p. 360) and Korteweg and de Vries in [26] in order to describe the propagation of small amplitude long water waves in a uniform channel. This equation is now commonly used to model unidirectional propagation of small amplitude long waves in nonlinear dispersive systems.

Since in many physical applications the region is finite, people are also interested in properties of the KdV equations on a finite spacial domain. Moreover, Bona and Winther pointed out in [3] that the term y_x should be incorporated in the KdV equations to model the water waves when x denotes the spatial coordinate in a fixed frame. We refer to [1, 2, 13, 18, 20, 22, 27, 35] for the well-posedness results of initial-boundary-value problems of the KdV equations posed on a finite interval. From control theory point of view, we refer to [7, 38] for an overall review and recent progress on different kinds of KdV equations. In particular, when the spacial domain is of finite interval, we refer to [6, 15, 16, 19, 36, 37, 45] for the controllability and [8, 23, 30, 31, 34] for some stabilization results. We refer to [10, 24, 25, 28, 39, 40, 41] for studies on the KdV equations with periodic boundary conditions.

Rosier introduced in [36] the following set of critical lengths

$$\mathcal{N} := \left\{ 2\pi \sqrt{\frac{j^2 + l^2 + jl}{3}}; j, l \in \mathbb{N}^* \right\}$$

for the following KdV control system

$$\begin{cases} y_t + y_x + yy_x + y_{xxx} = 0, \\ y(t, 0) = y(t, L) = 0, \\ y_x(t, L) = u(t), \\ y(0, x) = y_0, \end{cases} \quad (1.2)$$

where $u(t) \in \mathbb{R}$ is the control. We refer to [9, 15, 36] for the well-posedness and controllability of system (1.2). Especially, Rosier proved in [36] that (1.2) is locally controllable around the origin by analyzing the corresponding linearized system and by means of Banach fixed point theorem, provided that the spacial domain is not critical, i.e. $L \notin \mathcal{N}$. However, this method does not work when $L \in \mathcal{N}$, since the corresponding linearized system of (1.2) around the origin is not any more controllable in this case. By using the ‘‘power series expansion’’ method, Coron and Crépeau in [15] obtained the local exact controllability around the origin of the nonlinear KdV equation (1.2) with the critical length $L = 2k\pi$ (i.e. taking $j = l = k$ in \mathcal{N}), provided that (see [14, Theorem 8.1 and Remark 8.2])

$$(j^2 + l^2 + jl = 3k^2 \text{ and } (j, l) \in \mathbb{N} \setminus \{0\}^2) \Rightarrow (j = l = k). \quad (1.3)$$

The cases with the other critical lengths have been studied by Cerpa in [6] and by Cerpa and Crépeau in [9] with the same method, where the authors have proved that the nonlinear term yy_x gives the local exact controllability around the origin.

If $L \notin \mathcal{N}$, it is proved by Perla Menzala, Vasconcellos and Zuazua in [34] that 0 is exponentially stable for the linearized equation (1.4)

$$\begin{cases} y_t + y_x + y_{xxx} = 0, \\ y(t, 0) = y(t, L) = 0, \\ y_x(t, L) = 0, \\ y(0, x) = y_0 \in L^2(0, L), \end{cases} \quad (1.4)$$

of (1.1) around 0. Furthermore, it is also proved in [34] that 0 is locally asymptotically stable for system (1.1). However, when $L \in \mathcal{N}$, it has been proved by Rosier in [36] that (1.4) admits a family of non-trivial solutions of the form $e^{\lambda t} v_\lambda(x)$ for some $\lambda \in i\mathbb{R}$, where $v_\lambda \in C^\infty([0, L]) \setminus \{0\}$ satisfies

$$\begin{cases} \lambda v_\lambda(x) + v'_\lambda(x) + v'''_\lambda(x) = 0, \\ v_\lambda(0) = v_\lambda(L) = v'_\lambda(0) = v'_\lambda(L) = 0. \end{cases}$$

For these critical lengths, it is therefore interesting to study the influence of the nonlinear term yy_x on the local asymptotic stability of 0 for the nonlinear KdV equation (1.1). This article is concerned with the stability property for system (1.1) with special critical length $L = 2\pi$. In this particular case, by Remark 3.6 of [36], $a(1 - \cos x)$, $a \in \mathbb{R}$ are steady solutions of (1.4).

Center manifolds play an important role in studying nonlinear systems. We refer to [5, 11, 21, 29, 42] and the references therein for center manifold theories on abstract Cauchy problems in Banach spaces. The authors in [5, 21, 29] investigated directly the evolution equations and gave some sufficient conditions for the existence and smoothness of center manifolds. While, the authors in [11] presented a general result on the invariant manifolds together with associated invariant foliations of the state space, which can be applied directly to C^1 semigroups in Banach space. But the method presented in [11] has no extension to the case of C^k -smoothness with $k > 1$. In [42], by using the method of graph transforms, some classical results about smoothness of invariant manifolds for maps and the technique of “lifting”, the existence, smoothness and attractivity of invariant manifolds for evolutionary process on general Banach spaces are proved when the nonlinear perturbation has a small global Lipschitz constant and is locally C^k -smooth near the trivial solution. Because of the existence of the nonlinear term in (1.1), the results presented in [5, 29] do not work for our system. Moreover, due to the fact that the linear operator in our system (1.1) with $L = 2\pi$ does not satisfy the resolvent estimates provided by [21], we cannot apply directly the results given in [21]. Thanks to the center manifold results given in [42], in this article, we show the existence and smoothness of a center manifold of (1.1) with $L = 2\pi$, and obtain that the stability property can be determined by a reduced system of dimension one. Furthermore, by studying the stability on this reduced one dimensional system, we obtain the local asymptotic stability of 0 for the original system (1.1) when $L = 2\pi$. The main result of this article is the following theorem.

Theorem 1.1 *Let us assume that $L = 2\pi$. Then $0 \in L^2(0, L)$ is (locally) asymptotically stable for the nonlinear KdV equation (1.1). More precisely:*

(i) For every $\varepsilon > 0$, there exists $\delta = \delta(\varepsilon) > 0$ such that, if $\|y_0\|_{L^2(0,L)} < \delta$, then

$$\|y(t, \cdot)\|_{L^2(0,L)} < \varepsilon, \quad \forall t \geq 0.$$

(ii) There exists $\delta_1 > 0$ such that, if $\|y_0\|_{L^2(0,L)} < \delta_1$, then

$$\lim_{t \rightarrow +\infty} \|y(t, \cdot)\|_{L^2(0,L)} = 0.$$

Remark 1.1 The existence of $\delta(\varepsilon)$ is trivial and well known. In fact, one can take $\delta(\varepsilon) = \varepsilon$ since $t \in [0, +\infty) \mapsto \|y(t, \cdot)\|_{L^2(0,2\pi)}$ is nonincreasing (see also Lemma 3.1 below). The nontrivial part of Theorem 1.1 is property (ii).

The organization of this paper is as follows: First, in Section 2, some basic properties of the linearized system (1.4) are given. Then, in Section 3, we prove some properties of a non local modification of the KdV equation (1.1) and then deduce the existence and smoothness of the center manifold. Finally, in Section 4, we analyze the dynamic on the center manifold, which concludes the proof of the main result, i.e. Theorem 1.1.

2 Preliminary

In this section, we give some properties for the linearized system (1.4) with $L = 2\pi$.

Set $X := L^2(0, L)$. Let $A : D(A) \rightarrow X$ be the linear operator defined by

$$A\varphi = -\varphi_x - \varphi_{xxx}$$

with

$$D(A) = \{\varphi \in H^3(0, L) : \varphi(0) = \varphi(L) = \varphi_x(L) = 0\}.$$

It is easily verified that both A and its adjoint A^* are dissipative. The following proposition follows from [33, Corollary. 4.4, Chapter 1]. See also [36].

Proposition 2.1 *A generates a C_0 -semigroup of contractions on $L^2(0, L)$.*

From now on, we denote by $\{S(t)\}_{t \geq 0}$ the C_0 -semigroup associated with A . Then $S(t)y_0$ is the mild solution of the linearized system (1.4) for any given initial data $y_0 \in L^2(0, L)$. By Proposition 2.1, we obtain the following lemma directly.

Lemma 2.1 *For every $y_0 \in L^2(0, L)$, we have*

$$\|S(t)y_0\|_{L^2(0,L)} \leq \|y_0\|_{L^2(0,L)}, \quad \forall t \geq 0.$$

Furthermore, the following Kato smoothing effect is given by Rosier [36, Proposition 3.2].

Lemma 2.2 *For every $y_0 \in L^2(0, L)$ and for every $T > 0$, we have $S(t)y_0 \in L^2(0, T; H^1(0, L))$ and*

$$\|S(t)y_0\|_{L^2(0,T;H^1(0,L))} \leq \left(\frac{4T+L}{3}\right)^{\frac{1}{2}} \|y_0\|_{L^2(0,L)}.$$

Proceeding as in [32], we can prove the following two results.

Lemma 2.3 *There exists a constant $C > 0$ such that for any $y_0 \in H_0^1(0, L)$, the solution $S(t)y_0$ of (1.4) fulfills*

$$\|S(t)y_0\|_{H_0^1(0,L)} \leq C \|y_0\|_{H_0^1(0,L)}, \quad \forall t \geq 0.$$

Proof. For any $U_0 \in D(A)$, let us define $U(t) := S(t)U_0$. Let $V(t) = U_t(t) = AU(t)$. Then V is the mild solution of the system

$$\begin{cases} V_t = AV, \\ V(0) = AU_0 \in L^2(0, L). \end{cases}$$

Hence, it follows from Lemma 2.1 that

$$\|V(t)\|_{L^2(0,L)} \leq \|V(0)\|_{L^2(0,L)}, \quad \forall t \geq 0.$$

Since $V(t) = AU(t)$, $V(0) = AU_0$, and the norms $\|U\|_{L^2(0,L)} + \|AU\|_{L^2(0,L)}$ and $\|U\|_{D(A)}$ are equivalent on $D(A)$, we conclude that, for some constant $C_1 > 0$ independent of U_0 and $t \geq 0$, we have

$$\|U(t)\|_{D(A)} \leq C_1 \|U_0\|_{D(A)}.$$

Then the result of Lemma 2.3 follows by a standard interpolation argument. ■

Our next proposition shows that $\{S(t)\}_{t \geq 0}$ is a compact semigroup.

Proposition 2.2 *Let $T > 0$. There exists a constant $C > 0$ such that, for every $y_0 \in L^2(0, L)$, we have*

$$\|S(t)y_0\|_{H_0^1(0,L)} \leq \frac{C}{\sqrt{t}} \|y_0\|_{L^2(0,L)}, \quad \forall t \in (0, T]. \quad (2.1)$$

Consequently, the C_0 -semigroup $\{S(t)\}_{t \geq 0}$ generated by A is compact.

Proof. Let $T > 0$ be fixed. For every $t \in (0, T]$ and for every $y_0 \in L^2(0, L)$, by Lemma 2.2, the estimate

$$\|S(\cdot)y_0\|_{L^2(0, \frac{t}{2}; H_0^1(0,L))} \leq \left(\frac{2t+L}{3}\right)^{\frac{1}{2}} \|y_0\|_{L^2(0,L)} \quad (2.2)$$

holds. Then, arguing by contradiction, we get the existence of $\tau \in (0, t/2]$ such that

$$\|S(\tau)y_0\|_{H_0^1(0,L)} \leq \left(\frac{2t+L}{3}\right)^{\frac{1}{2}} \sqrt{\frac{2}{t}} \|y_0\|_{L^2(0,L)}, \quad \forall y_0 \in L^2(0, L). \quad (2.3)$$

Now it follows from Lemma 2.3 and (2.3) that there exists $C' = C'(T) > 0$ such that, for every $t \in (0, T]$ and every $y_0 \in L^2(0, L)$,

$$\begin{aligned} \|S(t)y_0\|_{H_0^1(0,L)} &= \|S(t-\tau)S(\tau)y_0\|_{H_0^1(0,L)} \\ &\leq C \|S(\tau)y_0\|_{H_0^1(0,L)} \\ &\leq C \left(\frac{2t+L}{3}\right)^{\frac{1}{2}} \sqrt{\frac{2}{t}} \|y_0\|_{L^2(0,L)} \\ &\leq \frac{C'}{\sqrt{t}} \|y_0\|_{L^2(0,L)}. \end{aligned}$$

Thus, for any given $T > 0$, (2.1) holds. Since $H^1(0, L)$ is compactly embedded in $L^2(0, L)$, we conclude that $S(t)$ is compact. ■

Let us now consider the spectral properties of the operator A . Firstly, we give the definition of growth bound and essential growth bound of the infinitesimal generator of a linear C_0 -semigroup.

Definition 2.1 Let $K : D(K) \subset X \rightarrow X$ be the infinitesimal generator of a linear C_0 -semigroup $\{S_K(t)\}_{t \geq 0}$ on a Banach space X . We define $\omega_0(K) \in [-\infty, +\infty)$ the growth bound of K by

$$\omega_0(K) := \lim_{t \rightarrow +\infty} \frac{\ln \left(\|S_K(t)\|_{\mathcal{L}(X)} \right)}{t}.$$

The essential growth bound $\omega_{0,ess}(K) \in [-\infty, +\infty)$ of K is defined by

$$\omega_{0,ess}(K) := \lim_{t \rightarrow +\infty} \frac{\ln \left(\|S_K(t)\|_{ess} \right)}{t},$$

where $\|S_K(t)\|_{ess}$ is the essential norm of $S_K(t)$ defined by

$$\|S_K(t)\|_{ess} = \kappa(S_K(t)B_X(0, 1)),$$

where $B_X(0, 1) := \{x \in X : \|x\|_X \leq 1\}$ and, for each bounded set $B \subset X$,

$$\kappa(B) = \inf \{ \varepsilon > 0 : B \text{ can be covered by a finite number of balls of radius } \leq \varepsilon \}$$

is the Kuratovsky measure of non-compactness.

The following result is proved by Webb [43, Proposition 4.11, p. 166, Proposition 4.13, p.170] and by Engel and Nagel [17, Corollary 2.11, p. 241].

Theorem 2.1 Let $K : D(K) \subset X \rightarrow X$ be the infinitesimal generator of a linear C_0 -semigroup $\{S_K(t)\}_{t \geq 0}$ on a Banach space X . Then

$$\omega_0(K) = \max \left(\omega_{0,ess}(K), \max_{\lambda \in \sigma(K) \setminus \sigma_{ess}(K)} \operatorname{Re}(\lambda) \right).$$

Assume in addition that $\omega_{0,ess}(K) < \omega_0(K)$. Then for each $\gamma \in (\omega_{0,ess}(K), \omega_0(K)]$, $\{\lambda \in \sigma(K) : \operatorname{Re}(\lambda) \geq \gamma\} \subset \sigma_p(K)$ is nonempty, finite and contains only poles of the resolvent of K .

As a consequence of Proposition 2.2 and Theorem 2.1, one has the following lemma.

Lemma 2.4 All the spectrum of the linear operator A are point spectrum, i.e., $\sigma(A) = \sigma_p(A)$ and $\omega_0(A) = \max_{\lambda \in \sigma(A)} \operatorname{Re}(\lambda)$. Moreover, for each $\gamma \in (-\infty, \omega_0(A)]$, $\{\lambda \in \sigma(A) : \operatorname{Re}(\lambda) \geq \gamma\}$ is nonempty, finite and contains only poles of the resolvent of A .

From Lemma 2.1 and Lemma 2.4, one has

Lemma 2.5 *For every $\lambda \in \sigma(A)$, $\operatorname{Re}(\lambda) \leq 0$.*

Let us now prove the following lemma.

Lemma 2.6 *One has $\sigma_p(A) \cap i\mathbb{R} = \{0\}$. Moreover, the kernel of A is $a(1 - \cos x)$, $a \in \mathbb{R}$.*

Proof. We have $\lambda \in \sigma_p(A) \cap i\mathbb{R}$ if and only if there exists $\varphi \in H^3(0, L) \setminus \{0\}$ such that

$$\begin{cases} \lambda\varphi + \varphi_x + \varphi_{xxx} = 0, \\ \varphi(0) = \varphi(L) = \varphi_x(L) = 0. \end{cases} \quad (2.4)$$

Multiplying equation (2.4) by $\bar{\varphi}$, and then integrating over $[0, L]$, we obtain

$$\lambda \int_0^L \varphi \bar{\varphi} dx + \int_0^L \varphi_x \bar{\varphi} dx + \int_0^L \varphi_{xxx} \bar{\varphi} dx = 0. \quad (2.5)$$

Taking the real part of (2.5), we have

$$\int_0^L \frac{\varphi_x \bar{\varphi} + \bar{\varphi}_x \varphi}{2} dx + \int_0^L \frac{\varphi_{xxx} \bar{\varphi} + \bar{\varphi}_{xxx} \varphi}{2} dx = 0. \quad (2.6)$$

Integrating by parts in (2.6) and using (2.4), we get

$$\varphi_x(0) = 0.$$

Hence, $\lambda \in \sigma_p(A) \cap i\mathbb{R}$ if and only if there exists $\varphi \in H^3(0, L) \setminus \{0\}$ such that

$$\begin{cases} \lambda\varphi + \varphi_x + \varphi_{xxx} = 0, \\ \varphi(0) = \varphi(L) = \varphi_x(0) = \varphi_x(L) = 0, \end{cases}$$

and the result of this lemma follows directly from the proof of Rosier [36, Lemma 3.5]. ■

Combining Lemma 2.4, Lemma 2.5 and Lemma 2.6, we obtain the following corollary.

Corollary 2.2 *$0 \in \sigma(A) = \sigma_p(A)$ and the other eigenvalues of A have negative real parts which are bounded away from 0.*

3 Existence and smoothness of the center manifold

This section is devoted to show the existence and smoothness of the center manifold for system (1.1) with $L = 2\pi$ by applying the results given in [42]. We would like to mention that the linear operator A in our system (1.1) with $L = 2\pi$ does not satisfy the resolvent estimates required in [21]. In particular, A does not generate an analytic semigroup, but a C_0 -semigroup with a Gevrey property. We refer to [12] and [41] for this result. Hence, we cannot apply the results given in [21] to show the existence and smoothness of the center manifold.

In order to apply the results given in [42], we need to show that the nonlinear perturbation has a small global Lipschitz constant. To that end, we modify the nonlinear part of the original system (1.1) by using some smooth cut-off mapping, and consider the following equation

$$\begin{cases} y_t + y_x + y_{xxx} + \Phi_\varepsilon(\|y\|_{L^2(0,L)})yy_x = 0, \\ y(t, 0) = y(t, L) = 0, \\ y_x(t, L) = 0, \\ y(0, x) = y_0(x) \in L^2(0, L). \end{cases} \quad (3.1)$$

Here $\varepsilon > 0$ is small enough, and $\Phi_\varepsilon : [0, +\infty) \rightarrow [0, 1]$ is defined by

$$\Phi_\varepsilon(x) = \Phi\left(\frac{x}{\varepsilon}\right), \quad \forall x \in [0, +\infty),$$

where $\Phi \in C^\infty([0, +\infty); [0, 1])$ satisfies

$$\Phi(x) = \begin{cases} 1, & \text{when } x \in [0, \frac{1}{2}], \\ 0, & \text{when } x \in [1, +\infty), \end{cases}$$

and

$$\Phi' \leq 0.$$

It can be readily checked that

$$\begin{aligned} \Phi_\varepsilon(x) &= 1, & \text{when } x \in [0, \frac{1}{2}], \\ \Phi_\varepsilon(x) &= 0, & \text{when } x \in [\varepsilon, +\infty). \end{aligned} \quad (3.2)$$

Moreover, there exists some constant $C > 0$ such that

$$0 \leq -\Phi'_\varepsilon(x) \leq \frac{C}{\varepsilon}, \quad \forall x \in [0, +\infty). \quad (3.3)$$

In (3.3) and in the following, C denotes various positive constants, which may vary from line to line, but do not depend on $\varepsilon \in (0, 1]$ and $y_0 \in L^2(0, L)$.

3.1 Well-posedness of (3.1)

In this section, we prove the following proposition on the global (in positive time) existence and uniqueness of the solution to system (3.1).

Proposition 3.1 *For every $y_0 \in L^2(0, L)$, there exists a unique mild solution*

$$y \in C([0, +\infty); L^2(0, L)) \cap L^2_{loc}([0, +\infty); H^1_0(0, L))$$

of (3.1).

In order to prove this proposition, one first points out that

Lemma 3.1 *Let $T > 0$. If*

$$y \in C([0, T]; L^2(0, L)) \cap L^2(0, T; H^1_0(0, L))$$

is a mild solution of (3.1), then

$$\frac{d}{dt} \left(\int_0^L y^2(t, x) dx \right) \leq 0.$$

Proof. We multiply $y_t + y_x + y_{xxx} + \Phi_\varepsilon(\|y\|_{L^2(0, L)})yy_x = 0$ by y and integrate over $[0, L]$. Using the boundary conditions in (3.1) and integrations by parts, we get

$$\frac{1}{2} \frac{d}{dt} \int_0^L y^2 dx + \frac{1}{2} y_x^2(t, 0) = 0.$$

The lemma follows. ■

By Lemma 3.1, in order to prove Proposition 3.1, it is sufficient to prove local (in positive time) existence and uniqueness of the solution to system (3.1).

Proposition 3.2 *Let $\varepsilon > 0, \eta > 0$. There exists $T > 0$ such that for every $y_0 \in L^2(0, L)$ with $\|y_0\|_{L^2(0, L)} \leq \eta$, there exists a unique solution $y \in C([0, T]; L^2(0, L)) \cap L^2(0, T; H^1_0(0, L))$ of (3.1).*

Proof. The case where $\Phi_\varepsilon \equiv 1$ is proved in [34]. Adapting the proof given in [34], we get the existence of T together with the existence and uniqueness of mild solution y . We briefly give the proof since some estimates given in the proof will be used later on.

Using the variation of constants formula, system (3.1) can be written in the following integral form:

$$\begin{aligned} y(t, \cdot) &= S(t) y_0 + \int_0^t S(t-s) \Phi_\varepsilon \left(\|y(s, \cdot)\|_{L^2(0, L)} \right) y(s, \cdot) y_x(s, \cdot) ds \\ &:= [\phi(y)](t). \end{aligned} \tag{3.4}$$

We will show that the nonlinear map ϕ is a contraction from $Y_T := C([0, T]; L^2(0, L)) \cap L^2(0, T; H_0^1(0, L))$ into itself when $T > 0$ is small enough.

Firstly, we prove that ϕ maps continuously Y_T into itself. Let us first show that if $y \in Y_T$, $\Phi_\varepsilon(\|y\|_{L^2(0, L)}) yy_x \in L^1(0, T; L^2(0, L))$ and the map $y \rightarrow \Phi_\varepsilon(\|y\|_{L^2(0, L)}) yy_x$ is continuous. Indeed, let $y, z \in Y_T$. Applying the triangular inequality, Hölder's inequality and Sobolev's embedding $H_0^1(0, L) \subset C^0([0, L])$ together with (3.3), we get

$$\begin{aligned}
& \left\| \Phi_\varepsilon(\|y\|_{L^2(0, L)}) yy_x - \Phi_\varepsilon(\|z\|_{L^2(0, L)}) zz_x \right\|_{L^1(0, T; L^2(0, L))} \\
& \leq \|(yy_x - zz_x)\|_{L^2(0, T; L^2(0, L))} + \left\| \left[\Phi_\varepsilon(\|y\|_{L^2(0, L)}) - \Phi_\varepsilon(\|z\|_{L^2(0, L)}) \right] zz_x \right\|_{L^1(0, T; L^2(0, L))} \\
& \leq \|(y - z)y_x + (y_x - z_x)z\|_{L^1(0, T; L^2(0, L))} + \frac{C}{\varepsilon} \left\| \|y - z\|_{L^2(0, L)} zz_x \right\|_{L^1(0, T; L^2(0, L))} \\
& \leq \int_0^T \|(y - z)y_x\|_{L^2(0, L)} dt + \int_0^T \|(y_x - z_x)z\|_{L^2(0, L)} dt \\
& \quad + \frac{C}{\varepsilon} \int_0^T \|y - z\|_{L^2(0, L)} \|zz_x\|_{L^2(0, L)} dt \\
& \leq C \int_0^T \|y - z\|_{L^\infty(0, L)} \|y_x\|_{L^2(0, L)} dt + C \int_0^T \|z\|_{L^\infty(0, L)} \|y_x - z_x\|_{L^2(0, L)} dt \\
& \quad + \frac{C}{\varepsilon} \|y - z\|_{L^\infty(0, T; L^2(0, L))} \int_0^T \|z\|_{L^\infty(0, L)} \|z_x\|_{L^2(0, L)} dt \\
& \leq C \|y - z\|_{L^2(0, T; L^\infty(0, L))} \|y_x\|_{L^2(0, T; L^2(0, L))} \\
& \quad + C \|z\|_{L^2(0, T; L^\infty(0, L))} \|y_x - z_x\|_{L^2(0, T; L^2(0, L))} \\
& \quad + \frac{C}{\varepsilon} \|y - z\|_{L^\infty(0, T; L^2(0, L))} \|z\|_{L^2(0, T; L^\infty(0, L))} \|z_x\|_{L^2(0, T; L^2(0, L))}. \tag{3.5}
\end{aligned}$$

By the classical Gagliardo-Nirenberg inequality, we have

$$\|u\|_{L^\infty(0, L)} \leq C \|u\|_{L^2(0, L)}^{\frac{1}{2}} \|u_x\|_{L^2(0, L)}^{\frac{1}{2}}, \quad \forall u \in H_0^1(0, L). \tag{3.6}$$

Hence,

$$\begin{aligned}
\int_0^T \|u\|_{L^\infty(0, L)}^2 dt & \leq C \int_0^T \|u\|_{L^2(0, L)} \|u_x\|_{L^2(0, L)} dt \\
& \leq C \|u\|_{L^\infty(0, T; L^2(0, L))} \int_0^T \|u_x\|_{L^2(0, L)} dt \\
& \leq C \|u\|_{L^\infty(0, T; L^2(0, L))} T^{\frac{1}{2}} \|u_x\|_{L^2(0, T; L^2(0, L))}.
\end{aligned}$$

Consequently, we get

$$\begin{aligned}
\|u\|_{L^2(0, T; L^\infty(0, L))} & \leq C \|u\|_{L^\infty(0, T; L^2(0, L))}^{\frac{1}{2}} T^{\frac{1}{4}} \|u_x\|_{L^2(0, T; L^2(0, L))}^{\frac{1}{2}} \\
& \leq CT^{\frac{1}{4}} \|u\|_{Y_T}, \quad \forall u \in Y_T.
\end{aligned}$$

Thus, it follows from (3.5) that

$$\begin{aligned}
& \left\| \Phi_\varepsilon \left(\|y\|_{L^2(0,L)} \right) yy_x - \Phi_\varepsilon \left(\|z\|_{L^2(0,L)} \right) zz_x \right\|_{L^1(0,T;L^2(0,L))} \\
& \leq CT^{\frac{1}{4}} \|y - z\|_{Y_T} \|y_x\|_{L^2(0,T;L^2(0,L))} + CT^{\frac{1}{4}} \|z\|_{Y_T} \|y_x - z_x\|_{L^2(0,T;L^2(0,L))} \\
& \quad + \frac{C}{\varepsilon} \|y - z\|_{L^\infty(0,T;L^2(0,L))} T^{\frac{1}{4}} \|z\|_{Y_T} \|z_x\|_{L^2(0,T;L^2(0,L))} \\
& \leq \|y - z\|_{Y_T} T^{\frac{1}{4}} C \left(\|y\|_{Y_T} + \|z\|_{Y_T} + \frac{1}{\varepsilon} \|z\|_{Y_T}^2 \right), \tag{3.7}
\end{aligned}$$

which implies that $\Phi_\varepsilon \left(\|y\|_{L^2(0,L)} \right) yy_x \in L^1(0, T; L^2(0, L))$ and that the map

$$y \rightarrow \Phi_\varepsilon \left(\|y\|_{L^2(0,L)} \right) yy_x$$

is continuous from Y_T to $L^1(0, T; L^2(0, L))$.

By Proposition 4.1 in [36], we obtain that

$$\int_0^t S(t-s) \Phi_\varepsilon \left(\|y(s, \cdot)\|_{L^2(0,L)} \right) y(s, \cdot) y_x(s, \cdot) ds$$

lies in Y_T , and the map

$$\Phi_\varepsilon \left(\|y\|_{L^2(0,L)} \right) yy_x \rightarrow \int_0^t S(t-s) \Phi_\varepsilon \left(\|y(s, \cdot)\|_{L^2(0,L)} \right) y(s, \cdot) y_x(s, \cdot) ds$$

is continuous. This fact, together with the continuity of the map $y \rightarrow \Phi_\varepsilon \left(\|y\|_{L^2(0,L)} \right) yy_x$ from Y_T to $L^1(0, T; L^2(0, L))$ and $S(t)y_0 \in Y_T$ (thanks to Lemma 2.1 and Lemma 2.2), leads to the conclusion that ϕ maps continuously Y_T into itself.

Let us now prove that ϕ is a contraction in a suitable ball B_R of Y_T when $T > 0$ is small enough. Obviously,

$$\phi(y) - \phi(z) = \int_0^t S(t-s) \left[\Phi_\varepsilon \left(\|y\|_{L^2(0,L)} \right) yy_x(s) - \Phi_\varepsilon \left(\|z\|_{L^2(0,L)} \right) zz_x(s) \right] ds.$$

In view of the proof of Proposition 4.1 in [36] and (3.7), we deduce that

$$\begin{aligned}
& \|\phi(y) - \phi(z)\|_{Y_T} \\
& \leq \left(1 + \left(\frac{T + 2L}{3} \right)^{\frac{1}{2}} \right) \left\| \Phi_\varepsilon \left(\|y\|_{L^2(0,L)} \right) yy_x - \Phi_\varepsilon \left(\|z\|_{L^2(0,L)} \right) zz_x \right\|_{L^1(0,T;L^2(0,L))} \\
& \leq C \left(1 + \sqrt{T} \right) \left\| \Phi_\varepsilon \left(\|y\|_{L^2(0,L)} \right) yy_x - \Phi_\varepsilon \left(\|z\|_{L^2(0,L)} \right) zz_x \right\|_{L^1(0,T;L^2(0,L))} \\
& \leq C \left(1 + \sqrt{T} \right) \|y - z\|_{Y_T} T^{\frac{1}{4}} \left(\|y\|_{Y_T} + \|z\|_{Y_T} + \frac{1}{\varepsilon} \|z\|_{Y_T}^2 \right), \tag{3.8}
\end{aligned}$$

which shows that ϕ is a contraction in the ball B_R of Y_T if

$$C \left(1 + \sqrt{T}\right) T^{\frac{1}{4}} \left(2R + \frac{1}{\varepsilon} R^2\right) < 1. \quad (3.9)$$

Therefore, the proof will be complete if we could show that for a suitable choice of R and T satisfying (3.9), the map ϕ sends B_R into itself.

It can be deduced from the definition of $\phi(y)$ given in (3.4), Lemma 2.1, Lemma 2.2 and (3.8) with $z = 0$ that there exists $\bar{C} > 0$ independent of $\varepsilon \in (0, 1]$, $y_0 \in L^2(0, L)$ and $T > 0$, such that

$$\begin{aligned} \|\phi(y)\|_{Y_T} &\leq \left(1 + \left(\frac{4T + L}{3}\right)^{\frac{1}{2}}\right) \|y_0\|_{L^2(0,L)} + \|y\|_{Y_T}^2 T^{\frac{1}{4}} C \left(1 + \sqrt{T}\right) \\ &\leq \left(1 + \left(\frac{4T + L}{3}\right)^{\frac{1}{2}}\right) \|y_0\|_{L^2(0,L)} + R^2 T^{\frac{1}{4}} C \left(1 + \sqrt{T}\right) \\ &\leq \bar{C} \left(1 + \sqrt{T}\right) \left(\|y_0\|_{L^2(0,L)} + R^2 T^{\frac{1}{4}}\right), \quad \forall y \in B_R. \end{aligned}$$

Now let $\|y_0\|_{L^2(0,L)} \leq \eta$, and set $R := 2\eta\bar{C}$. Then

$$\|\phi(y)\|_{Y_T} \leq \eta\bar{C} \left(1 + \sqrt{T}\right) \left(1 + 4\bar{C}^2\eta T^{\frac{1}{4}}\right), \quad \forall y \in B_R. \quad (3.10)$$

It is clear that we can choose $T > 0$ sufficiently small such that

$$\left(1 + \sqrt{T}\right) \left(1 + 4\eta\bar{C}^2 T^{\frac{1}{4}}\right) \leq 2,$$

which, together with (3.10) implies that ϕ maps B_R into itself. Moreover, decreasing T if necessary allows us to guarantee (3.9) as well. The proof of Proposition 3.2 is complete. ■

Proposition 3.3 *There exists $C > 0$ such that for every $\varepsilon > 0$, for every $y_0 \in L^2(0, L)$ and for every $T > 0$, the unique solution of (3.1) satisfies*

$$\|y\|_{L^2(0,T;H_0^1(0,L))}^2 \leq \frac{8T + 2L}{3} \|y_0\|_{L^2(0,L)}^2 + CT \|y_0\|_{L^2(0,L)}^4. \quad (3.11)$$

Proof. Proceeding as in [36], we multiply the first equation in (3.1) by xy and integrate over $(0, L) \times (0, T)$. Then, by Lemma 3.1, we obtain

$$\begin{aligned} &\int_0^T \int_0^L y_x^2 dx dt + \frac{1}{3} \int_0^L xy^2(x, T) dx \\ &= \frac{1}{3} \int_0^T \int_0^L y^2 dx dt + \frac{1}{3} \int_0^L xy_0^2 dx - \frac{2}{3} \int_0^T \Phi_\varepsilon(\|y\|_{L^2(0,L)}) \int_0^L xy^2 y_x dx dt \\ &\leq \frac{T + L}{3} \|y_0\|_{L^2(0,L)}^2 + \frac{2}{3} \int_0^T \Phi_\varepsilon(\|y\|_{L^2(0,L)}) \left| \int_0^L xy^2 y_x dx \right| dt. \end{aligned} \quad (3.12)$$

Since

$$\int_0^L xy^2 y_x dx = -\frac{1}{3} \int_0^L y^3 dx,$$

it follows from (3.12) that

$$\begin{aligned} & \int_0^T \int_0^L y_x^2 dx dt + \frac{1}{3} \int_0^L xy^2(x, T) dx \\ & \leq \frac{T+L}{3} \|y_0\|_{L^2(0,L)}^2 + \frac{2}{9} \int_0^T \Phi_\varepsilon(\|y\|_{L^2(0,L)}) \int_0^L |y|^3 dx dt \\ & \leq \frac{T+L}{3} \|y_0\|_{L^2(0,L)}^2 + \frac{2}{9} \int_0^T \int_0^L |y|^3 dx dt. \end{aligned}$$

Hence,

$$\|y\|_{L^2(0,T;H_0^1(0,L))}^2 \leq \frac{4T+L}{3} \|y_0\|_{L^2(0,L)}^2 + \frac{2}{9} \int_0^T \int_0^L |y|^3 dx dt. \quad (3.13)$$

Furthermore, by Lemma 3.1, the continuous Sobolev embedding $H_0^1(0, L) \subset C^0([0, L])$, Poincaré inequality and Hölder's inequality, we have

$$\begin{aligned} \int_0^T \int_0^L |y|^3 dx dt & \leq C \int_0^T \|y\|_{H_0^1(0,L)} \left(\int_0^L |y|^2 dx \right) dt \\ & \leq C \|y_0\|_{L^2(0,L)}^2 \int_0^T \|y\|_{H_0^1(0,L)} dt \\ & \leq C \|y_0\|_{L^2(0,L)}^2 \sqrt{T} \left(\int_0^T \|y\|_{H_0^1(0,L)}^2 dt \right)^{\frac{1}{2}} \\ & = C\sqrt{T} \|y_0\|_{L^2(0,L)}^2 \|y\|_{L^2(0,T;H_0^1(0,L))}. \end{aligned}$$

Now, using the above inequality in (3.13) we have

$$\begin{aligned} & \|y\|_{L^2(0,T;H_0^1(0,L))}^2 \\ & \leq \frac{4T+L}{3} \|y_0\|_{L^2(0,L)}^2 + C\sqrt{T} \|y_0\|_{L^2(0,L)}^2 \|y\|_{L^2(0,T;H_0^1(0,L))} \\ & \leq \frac{4T+L}{3} \|y_0\|_{L^2(0,L)}^2 + CT \|y_0\|_{L^2(0,L)}^4 + \frac{1}{2} \|y\|_{L^2(0,T;H_0^1(0,L))}^2. \end{aligned}$$

Therefore, we get

$$\|y\|_{L^2(0,T;H_0^1(0,L))}^2 \leq \frac{8T+2L}{3} \|y_0\|_{L^2(0,L)}^2 + CT \|y_0\|_{L^2(0,L)}^4.$$

This concludes the proof of Proposition 3.3. ■

Remark 3.1 According to Proposition 3.3, we have, for every $\tau \in [0, T]$,

$$\|y\|_{L^2(\tau, T; H_0^1(0, L))}^2 \leq \frac{8(T - \tau) + 2L}{3} \|y(\tau, \cdot)\|_{L^2(0, L)}^2 + C(T - \tau) \|y(\tau, \cdot)\|_{L^2(0, L)}^4.$$

It follows that, if $\tau \in [0, T]$ is such that $\|y(\tau, \cdot)\|_{L^2(0, L)} = \varepsilon$, then

$$\begin{aligned} \|y\|_{L^2(\tau, T; H_0^1(0, L))}^2 &\leq \frac{8(T - \tau) + 2L}{3} \varepsilon^2 + C(T - \tau) \varepsilon^4 \\ &\leq \frac{8T + 2L}{3} \varepsilon^2 + CT \varepsilon^4. \end{aligned}$$

Lemma 3.2 Let $T > 0$. There exist $\eta > 0$ and $C > 0$, such that, for every $\varepsilon \in (0, 1]$ and for every $y_0 \in L^2(0, L)$ with $\|y_0\|_{L^2(0, L)} \leq \eta$, there exists a unique mild solution $y : [0, T] \times [0, L] \rightarrow \mathbb{R}$ of (3.1) which satisfies

$$\|y(t, \cdot)\|_{H_0^1(0, L)} \leq \frac{C}{\sqrt{t}} \|y_0\|_{L^2(0, L)}, \quad \forall t \in (0, T].$$

Proof. From Proposition 2.2 and (3.4), we deduce that

$$\begin{aligned} \|y(t, \cdot)\|_{H_0^1(0, L)} &\leq \|S(t) y_0\|_{H_0^1(0, L)} + \int_0^t \left\| S(t-s) \Phi_\varepsilon \left(\|y(s, \cdot)\|_{L^2(0, L)} \right) y(s, \cdot) y_x(s, \cdot) \right\|_{H_0^1(0, L)} ds \\ &\leq \frac{C}{\sqrt{t}} \|y_0\|_{L^2(0, L)} + \int_0^t \frac{C}{\sqrt{t-s}} \|y(s, \cdot) y_x(s, \cdot)\|_{L^2(0, L)} ds. \end{aligned} \quad (3.14)$$

As a consequence of Lemma 3.1 and (3.6), we have

$$\begin{aligned} \|y(s, \cdot) y_x(s, \cdot)\|_{L^2(0, L)} &\leq \|y(s, \cdot)\|_{L^\infty(0, L)} \|y_x(s, \cdot)\|_{L^2(0, L)} \\ &\leq C \|y(s, \cdot)\|_{L^2(0, L)}^{\frac{1}{2}} \|y_x(s, \cdot)\|_{L^2(0, L)}^{\frac{3}{2}} \\ &\leq C \|y_0\|_{L^2(0, L)}^{\frac{1}{2}} \|y(s, \cdot)\|_{H_0^1(0, L)}^{\frac{3}{2}}. \end{aligned} \quad (3.15)$$

Substituting (3.15) into (3.14), we obtain

$$\|y(t, \cdot)\|_{H_0^1(0, L)} \leq \frac{C}{\sqrt{t}} \|y_0\|_{L^2(0, L)} + \|y_0\|_{L^2(0, L)}^{\frac{1}{2}} \int_0^t \frac{C}{\sqrt{t-s}} \|y(s, \cdot)\|_{H_0^1(0, L)}^{\frac{3}{2}} ds,$$

i.e.

$$\begin{aligned} &\sqrt{t} \|y(t, \cdot)\|_{H_0^1(0, L)} \\ &\leq C \|y_0\|_{L^2(0, L)} + \|y_0\|_{L^2(0, L)}^{\frac{1}{2}} \sqrt{t} \int_0^t \frac{C}{s^{\frac{3}{4}} \sqrt{t-s}} \left(\sqrt{s} \|y(s, \cdot)\|_{H_0^1(0, L)} \right)^{\frac{3}{2}} ds. \end{aligned} \quad (3.16)$$

Let $\bar{C} > C$. We claim that there exists $\eta > 0$ (small enough) such that, for every $\varepsilon \in (0, 1]$ and for every $y_0 \in L^2(0, L)$ such that $\|y_0\|_{L^2(0, L)} \leq \eta$, we have

$$\xi(t) < \bar{C} \|y_0\|_{L^2(0, L)}, \quad \forall t \in (0, T], \quad (3.17)$$

where $\xi(t) := \sqrt{t} \|y(t, \cdot)\|_{H_0^1(0, L)}$. Let us argue by contradiction. Suppose that (3.17) is not valid. Then there exists $\tau \in (0, T]$ such that

$$\xi(\tau) = \bar{C} \|y_0\|_{L^2(0, L)} \quad \text{and} \quad \xi(t) < \bar{C} \|y_0\|_{L^2(0, L)}, \quad \forall t \in (0, \tau). \quad (3.18)$$

Thus by (3.16), we have

$$\begin{aligned} \xi(\tau) &\leq C \|y_0\|_{L^2(0, L)} + \|y_0\|_{L^2(0, L)}^{\frac{1}{2}} \sqrt{\tau} \int_0^\tau \frac{C}{s^{\frac{3}{4}} \sqrt{\tau - s}} \left(\bar{C} \|y_0\|_{L^2(0, L)} \right)^{\frac{3}{2}} ds \\ &= C \|y_0\|_{L^2(0, L)} + \|y_0\|_{L^2(0, L)}^2 \sqrt{\tau} C (\bar{C})^{\frac{3}{2}} \int_0^\tau \frac{1}{s^{\frac{3}{4}} \sqrt{\tau - s}} ds \\ &= \|y_0\|_{L^2(0, L)} \left(C + \|y_0\|_{L^2(0, L)} \sqrt{\tau} C (\bar{C})^{\frac{3}{2}} \int_0^\tau \frac{1}{s^{\frac{3}{4}} \sqrt{\tau - s}} ds \right). \end{aligned}$$

It can be readily checked that if $\|y_0\|_{L^2(0, L)}$ is small enough, we get $\xi(\tau) < \bar{C} \|y_0\|_{L^2(0, L)}$, which leads to a contradiction with (3.18). This concludes the proof of Lemma 3.2. ■

3.2 Properties of the semigroup generated by (3.1)

Let

$$\mathcal{S}(t) : L^2(0, L) \rightarrow L^2(0, L), \quad t \geq 0$$

be the semigroup on $L^2(0, L)$ defined by

$$\mathcal{S}(t)(y_0) := y(t, x),$$

where $y(t, x)$ is the unique solution of (3.1) with respect to the initial value $y_0 \in L^2(0, L)$. Let $T > 0$. Then, for every $t \in [0, T]$, $\mathcal{S}(t)$ can be decomposed as

$$\mathcal{S}(t) = S(t) + R(t),$$

or equivalently,

$$y(t, x) = z(t, x) + \alpha(t, x),$$

where, as above, for every $y_0 \in L^2(0, L)$, $z(t, \cdot) := S(t)y_0$ is the unique solution of

$$\begin{cases} z_t + z_x + z_{xxx} = 0, \\ z(t, 0) = z(t, L) = 0, \\ z_x(t, L) = 0 \\ z(0, x) = y_0 \end{cases}$$

and $\alpha(t, \cdot) := R(t)y_0$ is the unique solution of

$$\begin{cases} \alpha_t + \alpha_x + \alpha_{xxx} + \Phi_\varepsilon \left(\|z + \alpha\|_{L^2(0,L)} \right) (z_x \alpha + \alpha_x z + z_x z + \alpha_x \alpha) = 0, \\ \alpha(t, 0) = \alpha(t, L) = 0, \\ \alpha_x(t, L) = 0, \\ \alpha(0, x) = 0. \end{cases}$$

Let

$$M := \{\alpha\varphi : \alpha \in \mathbb{R}\},$$

where

$$\varphi(x) = \frac{1}{\sqrt{3\pi}} (1 - \cos x). \quad (3.19)$$

Let us recall that, by Lemma 2.6, $\varphi(x)$ is an eigenfunction of the linear operator A for the linearized system (1.4) corresponding to the eigenvalue 0 and M is the eigenspace corresponding to this eigenvalue. Then we can do the following decomposition of $X = L^2(0, L)$:

$$X = M \oplus M^\perp.$$

The projection $P : X \rightarrow M$ is given by

$$Py(t, x) = p(t)\varphi(x),$$

where

$$p(t) := \int_0^L y(t, x)\varphi(x)dx, \quad (3.20)$$

and the projection $Q : X \rightarrow M^\perp$ is given by $I - P$.

It is clear that $S(t)$ leaves M and M^\perp invariant and $S(t)$ commutes with P and Q . Denote by $S_1(t) : M \rightarrow M$ and $S_2(t) : M^\perp \rightarrow M^\perp$ the restriction of $S(t)$ on M and M^\perp respectively. Then $S_1(t) = Id$. Moreover, by Corollary 2.2, there exist $N \geq 1$ and $\omega > 0$ such that $\|S_2(t)\| \leq Ne^{-\omega t}, \forall t \geq 0$.

3.2.1 Global Lipschitzianity of the map $R(t) : L^2(0, L) \rightarrow L^2(0, L)$

The aim of this part is to prove and estimate the global Lipschitzianity of the map $R(t) : L^2(0, L) \rightarrow L^2(0, L)$. To that end, we consider

$$\begin{cases} \alpha_t + \alpha_x + \alpha_{xxx} + \Phi_\varepsilon \left(\|\alpha + z\|_{L^2(0,L)} \right) (z_x \alpha + \alpha_x z + z_x z + \alpha_x \alpha) = 0, \\ \alpha(t, 0) = \alpha(t, L) = 0, \\ \alpha_x(t, L) = 0, \\ \alpha(0, x) = 0, \end{cases}$$

and

$$\begin{cases} \bar{\alpha}_t + \bar{\alpha}_x + \bar{\alpha}_{xxx} + \Phi_\varepsilon \left(\|\bar{z} + \bar{\alpha}\|_{L^2(0,L)} \right) (\bar{z}_x \bar{\alpha} + \bar{\alpha}_x \bar{z} + \bar{z}_x \bar{z} + \bar{\alpha}_x \bar{\alpha}) = 0, \\ \bar{\alpha}(t, 0) = \bar{\alpha}(t, L) = 0, \\ \bar{\alpha}_x(t, L) = 0, \\ \bar{\alpha}(0, x) = 0, \end{cases}$$

where z is the solution of

$$\begin{cases} z_t + z_x + z_{xxx} = 0, \\ z(t, 0) = z(t, L) = 0, \\ z_x(t, L) = 0, \\ z(0, x) = y_0 \in L^2(0, L), \end{cases}$$

and \bar{z} is the solution of

$$\begin{cases} \bar{z}_t + \bar{z}_x + \bar{z}_{xxx} = 0, \\ \bar{z}(t, 0) = \bar{z}(t, L) = 0, \\ \bar{z}_x(t, L) = 0, \\ z(0, x) = \bar{y}_0 \in L^2(0, L). \end{cases}$$

Set

$$\begin{aligned} \Delta &:= \alpha - \bar{\alpha}, \quad y := \alpha + z, \quad \bar{y} := \bar{z} + \bar{\alpha}, \\ \Phi_1 &:= \Phi_\varepsilon \left(\|y\|_{L^2(0,L)} \right), \quad \Phi_2 := \Phi_\varepsilon \left(\|\bar{y}\|_{L^2(0,L)} \right). \end{aligned}$$

Then we obtain

$$\begin{cases} \Delta_t + \Delta_x + \Delta_{xxx} = -\Phi_1 y y_x + \Phi_2 \bar{y} \bar{y}_x = \Phi_1 [-(\alpha + z) \Delta_x - (\bar{\alpha}_x + z_x) \Delta - \bar{\alpha} (z - \bar{z})_x \\ \quad - \bar{\alpha}_x (z - \bar{z}) - z_x z + \bar{z}_x \bar{z}] - (\Phi_1 - \Phi_2) (\bar{z}_x \bar{\alpha} + \bar{\alpha}_x \bar{z} + \bar{z}_x \bar{z} + \bar{\alpha}_x \bar{\alpha}), \\ \Delta(t, 0) = \Delta(t, L) = 0, \\ \Delta_x(t, L) = 0, \\ \Delta(0, x) = 0. \end{cases} \tag{3.21}$$

Moreover, by the definition of Φ_1 , Φ_2 and (3.2), we get

$$\Phi_1 = \Phi_2 = 0, \quad \forall \|y\|_{L^2(0,L)} \geq \varepsilon, \quad \forall \|\bar{y}\|_{L^2(0,L)} \geq \varepsilon. \tag{3.22}$$

We first give the following estimate of the L^2 -norm of Δ .

Lemma 3.3 *Let $T > 0$. Then there exists $C > 0$ such that*

$$\|\Delta(t, \cdot)\|_{L^2(0,L)} \leq C, \quad \forall t \in [0, T], \quad \forall \varepsilon \in (0, 1], \quad \forall y_0 \in L^2(0, L), \quad \forall \bar{y}_0 \in L^2(0, L).$$

Proof. By integrating by parts in

$$\int_0^L \Delta (\Delta_t + \Delta_x + \Delta_{xxx} + \Phi_1 y y_x - \Phi_2 \bar{y} \bar{y}_x) dx = 0,$$

we get

$$\frac{1}{2} \frac{d}{dt} \int_0^L \Delta^2 dx + \frac{1}{2} \Delta_x^2(t, 0) = -\Phi_1 \int_0^L \Delta y y_x dx + \Phi_2 \int_0^L \Delta \bar{y} \bar{y}_x dx. \quad (3.23)$$

Note that $\Delta(t, 0) = \Delta(t, L) = 0$, by the continuous Sobolev embedding $H_0^1(0, L) \subset C^0([0, L])$ and Poincaré inequality, we obtain

$$\begin{aligned} \left| \int_0^L \Delta \bar{y} \bar{y}_x dx \right| &\leq \|\bar{y}\|_{L^\infty(0, L)} \int_0^L |\Delta \bar{y}_x| dx \\ &\leq C \|\bar{y}\|_{H_0^1(0, L)} \int_0^L |\Delta \bar{y}_x| dx \\ &\leq C \|\bar{y}_x\|_{L^2(0, L)} \int_0^L |\Delta \bar{y}_x| dx. \end{aligned}$$

In the above inequalities and in the following, C , unless otherwise specified, denotes various positive constants which may vary from line to line but are independent of $t \in [0, T]$, $\varepsilon \in (0, 1]$, $y_0 \in L^2(0, L)$ and $\bar{y}_0 \in L^2(0, L)$. Thus,

$$\left| \int_0^L \Delta \bar{y} \bar{y}_x dx \right| \leq C \|\bar{y}_x\|_{L^2(0, L)}^2 \|\Delta\|_{L^2(0, L)}.$$

Similarly, we have

$$\left| \int_0^L \Delta y y_x dx \right| \leq C \|y_x\|_{L^2(0, L)}^2 \|\Delta\|_{L^2(0, L)}.$$

Hence, it follows from (3.23) that

$$\frac{d}{dt} \int_0^L \Delta^2 dx + \Delta_x^2(t, 0) \leq C \left(\Phi_1 \|y_x\|_{L^2(0, L)}^2 + \Phi_2 \|\bar{y}_x\|_{L^2(0, L)}^2 \right) \|\Delta\|_{L^2(0, L)}.$$

In particular,

$$\frac{d}{dt} \int_0^L \Delta^2 dx \leq C \left(\Phi_1 \|y_x\|_{L^2(0, L)}^2 + \Phi_2 \|\bar{y}_x\|_{L^2(0, L)}^2 \right) \|\Delta\|_{L^2(0, L)}.$$

By Lemma 17 in [15] and Remark 3.1, we get

$$\begin{aligned} \int_0^L \Delta^2 dx &\leq 3 \left(\int_0^t C \left(\Phi_1 \|y_x\|_{L^2(0, L)}^2 + \Phi_2 \|\bar{y}_x\|_{L^2(0, L)}^2 \right) dt \right)^2 \\ &\leq 3C^2 \left(2 \left(\frac{8T + 2L}{3} \varepsilon^2 + CT\varepsilon^4 \right) \right)^2, \forall t \in [0, T]. \end{aligned}$$

The result follows. ■

For the sake of simplicity, we denote from now on by $L^2(L^2)$ the norm $L^2(0, T; L^2(0, L))$.

Lemma 3.4 *Let $T > 0$. Then there exists $C > 0$ such that*

$$\begin{aligned} & \|\Delta(t, \cdot)\|_{L^2(0,L)} \\ & \leq \int_0^T \left[\Phi_1 \left(\|\bar{\alpha}_x\|_{L^2(0,L)} + \|z_x\|_{L^2(0,L)} + \|\bar{z}_x\|_{L^2(0,L)} \right) \|(z - \bar{z})_x\|_{L^2(0,L)} \right. \\ & \quad \left. + |\Phi_1 - \Phi_2| \|\bar{z} + \bar{\alpha}\|_{L^2(0,L)}^{\frac{1}{2}} \|(\bar{z} + \bar{\alpha})_x\|_{L^2(0,L)}^{\frac{3}{2}} \right] dt \\ & \quad \times \exp \left[C \left(1 + \left\| \sqrt{\Phi_1} \alpha_x \right\|_{L^2(L^2)}^2 + \left\| \sqrt{\Phi_1} \bar{\alpha}_x \right\|_{L^2(L^2)}^2 + \left\| \sqrt{\Phi_1} z_x \right\|_{L^2(L^2)}^2 \right) \right], \end{aligned}$$

for every $t \in [0, T]$, for every $\varepsilon \in (0, 1]$, for every $y_0 \in L^2(0, L)$ and for every $\bar{y}_0 \in L^2(0, L)$.

Proof. We multiply the first equation of (3.21) by $2x\Delta$ and then integrate over $[0, L]$. By integrating by parts and using the boundary conditions of (3.21), we get

$$\begin{aligned} & \frac{d}{dt} \int_0^L x \Delta^2 dx + 3 \int_0^L \Delta_x^2 dx \\ & = \int_0^L \Delta^2 dx + \Phi_1 \times \left(-2 \int_0^L x \alpha \Delta \Delta_x dx + 4 \int_0^L x \bar{\alpha} \Delta \Delta_x dx + 2 \int_0^L x z \Delta \Delta_x dx \right. \\ & \quad + 2 \int_0^L \bar{\alpha} \Delta^2 dx + 2 \int_0^L z \Delta^2 dx - 2 \int_0^L x \Delta \bar{\alpha} (z - \bar{z})_x dx \\ & \quad - 2 \int_0^L x \Delta \bar{\alpha}_x (z - \bar{z}) dx - 2 \int_0^L x \Delta z_x (z - \bar{z}) dx - 2 \int_0^L x \Delta \bar{z} (z - \bar{z})_x dx \left. \right) \\ & \quad - (\Phi_1 - \Phi_2) \int_0^L 2x \Delta (\bar{z}_x \bar{\alpha} + \bar{\alpha}_x \bar{z} + \bar{z}_x \bar{z} + \bar{\alpha}_x \bar{\alpha}) dx. \end{aligned} \tag{3.24}$$

Note that $\alpha(t, 0) = \alpha(t, L) = 0$, by the continuous Sobolev embedding $H_0^1(0, L) \subset C^0([0, L])$ and Poincaré inequality, there exists $C = C(L) > 0$ such that

$$2 \left| \int_0^L x \alpha \Delta \Delta_x dx \right| \leq C \|\alpha_x\|_{L^2(0,L)} \int_0^L |x \Delta \Delta_x| dx.$$

Thus,

$$\begin{aligned} 2 \left| \int_0^L x \alpha \Delta \Delta_x dx \right| & \leq C \|\alpha_x\|_{L^2(0,L)} \|\Delta_x\|_{L^2(0,L)} \|x \Delta\|_{L^2(0,L)} \\ & \leq \frac{1}{2} \|\Delta_x\|_{L^2(0,L)}^2 + \frac{1}{2} \left(C \|\alpha_x\|_{L^2(0,L)} \|x \Delta\|_{L^2(0,L)} \right)^2 \\ & \leq \frac{1}{2} \int_0^L \Delta_x^2 dx + C \|\alpha_x\|_{L^2(0,L)}^2 \int_0^L x \Delta^2 dx. \end{aligned} \tag{3.25}$$

Similarly,

$$4 \left| \int_0^L x \bar{\alpha} \Delta \Delta_x dx \right| \leq \frac{1}{2} \int_0^L \Delta_x^2 dx + C \|\bar{\alpha}_x\|_{L^2(0,L)}^2 \int_0^L x \Delta^2 dx, \tag{3.26}$$

$$2 \left| \int_0^L xz \Delta \Delta_x dx \right| \leq \frac{1}{2} \int_0^L \Delta_x^2 dx + C \|z_x\|_{L^2(0,L)}^2 \int_0^L x \Delta^2 dx. \quad (3.27)$$

Note that $\bar{\alpha}(t, 0) = \bar{\alpha}(t, L) = 0$, by the continuous Sobolev embedding $H_0^1(0, L) \subset C^0([0, L])$ and Poincaré inequality, we have

$$2 \left| \int_0^L \bar{\alpha} \Delta^2 dx \right| \leq C \|\bar{\alpha}_x\|_{L^2(0,L)} \int_0^L \Delta^2 dx. \quad (3.28)$$

From (3.28) and Lemma 16 in [15] with $a := \min \left\{ \frac{1}{\sqrt{2}} C^{-\frac{1}{2}} \|\bar{\alpha}_x\|_{L^2(0,L)}^{-\frac{1}{2}}, L \right\}$, there exists $C = C(L) > 0$ such that

$$2 \left| \int_0^L \bar{\alpha} \Delta^2 dx \right| \leq \frac{1}{4} \int_0^L \Delta_x^2 dx + C \left(\|\bar{\alpha}_x\|_{L^2(0,L)}^{\frac{3}{2}} + \|\bar{\alpha}_x\|_{L^2(0,L)} \right) \int_0^L x \Delta^2 dx. \quad (3.29)$$

Similarly, we have

$$2 \left| \int_0^L z \Delta^2 dx \right| \leq \frac{1}{4} \int_0^L \Delta_x^2 dx + C \left(\|z_x\|_{L^2(0,L)}^{\frac{3}{2}} + \|z_x\|_{L^2(0,L)} \right) \int_0^L x \Delta^2 dx. \quad (3.30)$$

By Lemma 16 in [15], there exists $C = C(L) > 0$ such that

$$\int_0^L \Delta^2 dx \leq \frac{1}{2} \int_0^L \Delta_x^2 dx + C \int_0^L x \Delta^2 dx. \quad (3.31)$$

We have

$$\begin{aligned} 2 \left| \int_0^L x \Delta \bar{\alpha} (z - \bar{z})_x dx \right| &\leq C \|\bar{\alpha}_x\|_{L^2(0,L)} \left| \int_0^L x \Delta (z - \bar{z})_x dx \right| \\ &\leq C \|\bar{\alpha}_x\|_{L^2(0,L)} \left(\int_0^L x^2 \Delta^2 dx \right)^{\frac{1}{2}} \|(z - \bar{z})_x\|_{L^2(0,L)} \\ &\leq C \|\bar{\alpha}_x\|_{L^2(0,L)} \left(\int_0^L x \Delta^2 dx \right)^{\frac{1}{2}} \|(z - \bar{z})_x\|_{L^2(0,L)}. \end{aligned} \quad (3.32)$$

Similarly, we can obtain

$$2 \left| \int_0^L x \Delta \bar{\alpha}_x (z - \bar{z}) dx \right| \leq C \|(z - \bar{z})_x\|_{L^2(0,L)} \left(\int_0^L x \Delta^2 dx \right)^{\frac{1}{2}} \|\bar{\alpha}_x\|_{L^2(0,L)}, \quad (3.33)$$

$$2 \left| \int_0^L x \Delta z_x (z - \bar{z}) dx \right| \leq C \|(z - \bar{z})_x\|_{L^2(0,L)} \left(\int_0^L x \Delta^2 dx \right)^{\frac{1}{2}} \|z_x\|_{L^2(0,L)}, \quad (3.34)$$

$$2 \left| \int_0^L x \Delta \bar{z} (z - \bar{z})_x dx \right| \leq C \|\bar{z}_x\|_{L^2(0,L)} \left(\int_0^L x \Delta^2 dx \right)^{\frac{1}{2}} \|(z - \bar{z})_x\|_{L^2(0,L)}. \quad (3.35)$$

Moreover, we have

$$\begin{aligned}
& \left| \int_0^L 2x\Delta (\bar{z}_x\bar{\alpha} + \bar{\alpha}_x\bar{z} + \bar{z}_x\bar{z} + \bar{\alpha}_x\bar{\alpha}) dx \right| \\
&= 2 \left| \int_0^L x\Delta (\bar{z}_x + \bar{\alpha}_x) (\bar{z} + \bar{\alpha}) dx \right| \\
&\leq 2 \|\bar{z} + \bar{\alpha}\|_{L^\infty(0,L)} \int_0^L |x\Delta (\bar{z} + \bar{\alpha})_x| dx \\
&\leq 2\sqrt{L} \|\bar{z} + \bar{\alpha}\|_{L^\infty(0,L)} \|(\bar{z} + \bar{\alpha})_x\|_{L^2(0,L)} \left(\int_0^L x\Delta^2 dx \right)^{\frac{1}{2}}. \tag{3.36}
\end{aligned}$$

Then, using the Gagliardo-Nirenberg inequality (3.6), it follows from (3.36) that

$$\begin{aligned}
& \left| \int_0^L 2x\Delta (\bar{z}_x\bar{\alpha} + \bar{\alpha}_x\bar{z} + \bar{z}_x\bar{z} + \bar{\alpha}_x\bar{\alpha}) dx \right| \\
&\leq C \|\bar{z} + \bar{\alpha}\|_{L^2(0,L)}^{\frac{1}{2}} \|(\bar{z} + \bar{\alpha})_x\|_{L^2(0,L)}^{\frac{3}{2}} \left(\int_0^L x\Delta^2 dx \right)^{\frac{1}{2}}. \tag{3.37}
\end{aligned}$$

Thus, using (3.24) to (3.37), we get

$$\begin{aligned}
& \frac{d}{dt} \int_0^L x\Delta^2 dx + \frac{1}{2} \int_0^L \Delta_x^2 dx \\
&\leq C \left(1 + \Phi_1 \left(\|\alpha_x\|_{L^2(0,L)}^2 + \|\bar{\alpha}_x\|_{L^2(0,L)}^2 + \|z_x\|_{L^2(0,L)}^2 \right) \right) \int_0^L x\Delta^2 dx \\
&\quad + C \left[\Phi_1 \left(\|\bar{\alpha}_x\|_{L^2(0,L)} + \|z_x\|_{L^2(0,L)} + \|\bar{z}_x\|_{L^2(0,L)} \right) \|(z - \bar{z})_x\|_{L^2(0,L)} \right. \\
&\quad \left. + |\Phi_1 - \Phi_2| \|\bar{z} + \bar{\alpha}\|_{L^2(0,L)}^{\frac{1}{2}} \|(\bar{z} + \bar{\alpha})_x\|_{L^2(0,L)}^{\frac{3}{2}} \right] \left(\int_0^L x\Delta^2 dx \right)^{\frac{1}{2}}. \tag{3.38}
\end{aligned}$$

In particular,

$$\begin{aligned}
& \frac{d}{dt} \int_0^L x\Delta^2 dx \\
&\leq C \left(1 + \Phi_1 \left(\|\alpha_x\|_{L^2(0,L)}^2 + \|\bar{\alpha}_x\|_{L^2(0,L)}^2 + \|z_x\|_{L^2(0,L)}^2 \right) \right) \int_0^L x\Delta^2 dx \\
&\quad + C \left[\Phi_1 \left(\|\bar{\alpha}_x\|_{L^2(0,L)} + \|z_x\|_{L^2(0,L)} + \|\bar{z}_x\|_{L^2(0,L)} \right) \|(z - \bar{z})_x\|_{L^2(0,L)} \right. \\
&\quad \left. + |\Phi_1 - \Phi_2| \|\bar{z} + \bar{\alpha}\|_{L^2(0,L)}^{\frac{1}{2}} \|(\bar{z} + \bar{\alpha})_x\|_{L^2(0,L)}^{\frac{3}{2}} \right] \left(\int_0^L x\Delta^2 dx \right)^{\frac{1}{2}}.
\end{aligned}$$

Then, by Lemma 17 in [15], we get

$$\int_0^L x\Delta^2 dx \leq W, \quad \forall t \in [0, T] \tag{3.39}$$

with

$$\begin{aligned}
W &:= 3C^2 \left[\int_0^T \left(\left(\|\Phi_1 \bar{\alpha}_x\|_{L^2(0,L)} + \|\Phi_1 z_x\|_{L^2(0,L)} + \|\Phi_1 \bar{z}_x\|_{L^2(0,L)} \right) \|(z - \bar{z})_x\|_{L^2(0,L)} \right. \right. \\
&\quad \left. \left. + |\Phi_1 - \Phi_2| \|\bar{z} + \bar{\alpha}\|_{L^2(0,L)}^{\frac{1}{2}} \|(\bar{z} + \bar{\alpha})_x\|_{L^2(0,L)}^{\frac{3}{2}} \right) dt \right]^2 \\
&\quad \times \exp \left[C \left(T + \left\| \sqrt{\Phi_1} \alpha_x \right\|_{L^2(L^2)}^2 + \left\| \sqrt{\Phi_1} \bar{\alpha}_x \right\|_{L^2(L^2)}^2 + \left\| \sqrt{\Phi_1} z_x \right\|_{L^2(L^2)}^2 \right) \right]. \tag{3.40}
\end{aligned}$$

Now integrating (3.38) over $[0, T]$ and using (3.39), we have

$$\begin{aligned}
&\int_0^L x \Delta^2(T, x) dx + \frac{1}{2} \int_0^T \int_0^L \Delta_x^2 dx dt \\
&\leq C \int_0^T \left(1 + \Phi_1 \left(\|\alpha_x\|_{L^2(0,L)}^2 + \|\bar{\alpha}_x\|_{L^2(0,L)}^2 + \|z_x\|_{L^2(0,L)}^2 \right) \right) dt W \\
&\quad + C \int_0^T \left[\Phi_1 \left(\|\bar{\alpha}_x\|_{L^2(0,L)} + \|z_x\|_{L^2(0,L)} + \|\bar{z}_x\|_{L^2(0,L)} \right) \|(z - \bar{z})_x\|_{L^2(0,L)} \right. \\
&\quad \left. + |\Phi_1 - \Phi_2| \|\bar{z} + \bar{\alpha}\|_{L^2(0,L)}^{\frac{1}{2}} \|(\bar{z} + \bar{\alpha})_x\|_{L^2(0,L)}^{\frac{3}{2}} \right] dt W^{\frac{1}{2}}.
\end{aligned}$$

Then it follows that

$$\frac{1}{2} \int_0^T \int_0^L \Delta_x^2 dx dt \tag{3.41}$$

$$\begin{aligned}
&\leq C \int_0^T \left(1 + \Phi_1 \left(\|\alpha_x\|_{L^2(0,L)}^2 + \|\bar{\alpha}_x\|_{L^2(0,L)}^2 + \|z_x\|_{L^2(0,L)}^2 \right) \right) dt W + \frac{1}{2} W \\
&\quad + \frac{1}{2} \left[C \int_0^T \left(\Phi_1 \left(\|\bar{\alpha}_x\|_{L^2(0,L)} + \|z_x\|_{L^2(0,L)} + \|\bar{z}_x\|_{L^2(0,L)} \right) \|(z - \bar{z})_x\|_{L^2(0,L)} \right. \right. \\
&\quad \left. \left. + |\Phi_1 - \Phi_2| \|\bar{z} + \bar{\alpha}\|_{L^2(0,L)}^{\frac{1}{2}} \|(\bar{z} + \bar{\alpha})_x\|_{L^2(0,L)}^{\frac{3}{2}} \right) dt \right]^2. \tag{3.42}
\end{aligned}$$

Hence, combining (3.42) with (3.40), we obtain

$$\begin{aligned}
&\int_0^T \int_0^L \Delta_x^2 dx dt \\
&\leq C \left[\int_0^T \left(\left(\|\Phi_1 \bar{\alpha}_x\|_{L^2(0,L)} + \|\Phi_1 z_x\|_{L^2(0,L)} + \|\Phi_1 \bar{z}_x\|_{L^2(0,L)} \right) \|(z - \bar{z})_x\|_{L^2(0,L)} \right. \right. \\
&\quad \left. \left. + |\Phi_1 - \Phi_2| \|\bar{z} + \bar{\alpha}\|_{L^2(0,L)}^{\frac{1}{2}} \|(\bar{z} + \bar{\alpha})_x\|_{L^2(0,L)}^{\frac{3}{2}} \right) dt \right]^2 \\
&\quad \times \exp \left[C \left(T + \left\| \sqrt{\Phi_1} \alpha_x \right\|_{L^2(L^2)}^2 + \left\| \sqrt{\Phi_1} \bar{\alpha}_x \right\|_{L^2(L^2)}^2 + \left\| \sqrt{\Phi_1} z_x \right\|_{L^2(L^2)}^2 \right) \right]. \tag{3.43}
\end{aligned}$$

We multiply the first equation of (3.21) by Δ and integrate over $[0, L]$. Using the boundary conditions of (3.21) and integrations by parts, we get

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \int_0^L \Delta^2 dx + \frac{1}{2} \Delta_x^2(t, 0) \\
&= \Phi_1 \times \left(- \int_0^L \alpha \Delta_x \Delta dx + 2 \int_0^L \bar{\alpha} \Delta_x \Delta dx + \int_0^L z \Delta_x \Delta dx \right. \\
&\quad - \int_0^L \bar{\alpha} (z - \bar{z})_x \Delta dx - \int_0^L \bar{\alpha}_x (z - \bar{z}) \Delta dx \\
&\quad \left. - \int_0^L z_x (z - \bar{z}) \Delta dx - \int_0^L \bar{z} (z - \bar{z})_x \Delta dx \right) \\
&\quad - (\Phi_1 - \Phi_2) \int_0^L \Delta (\bar{z}_x \bar{\alpha} + \bar{\alpha}_x \bar{z} + \bar{z}_x \bar{z} + \bar{\alpha}_x \bar{\alpha}) dx.
\end{aligned} \tag{3.44}$$

It can be readily checked that

$$\begin{aligned}
\left| \int_0^L \alpha \Delta_x \Delta dx \right| &\leq \frac{1}{2} \int_0^L \Delta_x^2 dx + \frac{1}{2} \int_0^L \Delta^2 \alpha^2 dx \\
&\leq \frac{1}{2} \int_0^L \Delta_x^2 dx + C \|\alpha_x\|_{L^2(0,L)}^2 \int_0^L \Delta^2 dx.
\end{aligned} \tag{3.45}$$

Similarly, we have

$$\left| 2 \int_0^L \bar{\alpha} \Delta_x \Delta dx \right| \leq \frac{1}{2} \int_0^L \Delta_x^2 dx + C \|\bar{\alpha}_x\|_{L^2(0,L)}^2 \int_0^L \Delta^2 dx, \tag{3.46}$$

and

$$\left| \int_0^L z \Delta_x \Delta dx \right| \leq \frac{1}{2} \int_0^L \Delta_x^2 dx + C \|z_x\|_{L^2(0,L)}^2 \int_0^L \Delta^2 dx. \tag{3.47}$$

Similarly to (3.32), we get the following inequalities

$$\left| \int_0^L \bar{\alpha} (z - \bar{z})_x \Delta dx \right| \leq C \|\bar{\alpha}_x\|_{L^2(0,L)} \|(z - \bar{z})_x\|_{L^2(0,L)} \left(\int_0^L \Delta^2 dx \right)^{\frac{1}{2}}, \tag{3.48}$$

$$\left| \int_0^L \bar{\alpha}_x (z - \bar{z}) \Delta dx \right| \leq C \|(z - \bar{z})_x\|_{L^2(0,L)} \|\bar{\alpha}_x\|_{L^2(0,L)} \left(\int_0^L \Delta^2 dx \right)^{\frac{1}{2}}, \tag{3.49}$$

$$\left| \int_0^L z_x (z - \bar{z}) \Delta dx \right| \leq C \|(z - \bar{z})_x\|_{L^2(0,L)} \|z_x\|_{L^2(0,L)} \left(\int_0^L \Delta^2 dx \right)^{\frac{1}{2}}, \tag{3.50}$$

$$\left| \int_0^L \bar{z} (z - \bar{z})_x \Delta dx \right| \leq C \|\bar{z}_x\|_{L^2(0,L)} \|(z - \bar{z})_x\|_{L^2(0,L)} \left(\int_0^L \Delta^2 dx \right)^{\frac{1}{2}}. \tag{3.51}$$

Moreover, for the last term on the right-hand side of (3.44), using the same argument as for (3.37), we have

$$\begin{aligned} & \left| \int_0^L \Delta (\bar{z}_x \bar{\alpha} + \bar{\alpha}_x \bar{z} + \bar{z}_x \bar{z} + \bar{\alpha}_x \bar{\alpha}) dx \right| \\ & \leq C \|\bar{z} + \bar{\alpha}\|_{L^2(0,L)}^{\frac{1}{2}} \|(\bar{z} + \bar{\alpha})_x\|_{L^2(0,L)}^{\frac{3}{2}} \left(\int_0^L \Delta^2 dx \right)^{\frac{1}{2}}. \end{aligned} \quad (3.52)$$

Hence, by (3.44) to (3.52), we deduce that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_0^L \Delta^2 dx \\ & \leq \frac{3}{2} \int_0^L \Delta_x^2 dx + C \Phi_1 \left(\|\alpha_x\|_{L^2(0,L)}^2 + \|\bar{\alpha}_x\|_{L^2(0,L)}^2 + \|z_x\|_{L^2(0,L)}^2 \right) \int_0^L \Delta^2 dx \\ & \quad + C \left[\Phi_1 \left(2 \|\bar{\alpha}_x\|_{L^2(0,L)} + \|z_x\|_{L^2(0,L)} + \|\bar{z}_x\|_{L^2(0,L)} \right) \|(z - \bar{z})_x\|_{L^2(0,L)} \right. \\ & \quad \left. + |\Phi_1 - \Phi_2| \|\bar{z} + \bar{\alpha}\|_{L^2(0,L)}^{\frac{1}{2}} \|(\bar{z} + \bar{\alpha})_x\|_{L^2(0,L)}^{\frac{3}{2}} \right] \left(\int_0^L \Delta^2 dx \right)^{\frac{1}{2}}. \end{aligned}$$

Therefore, by (3.43) and Lemma 17 in [15], we get that, for every $t \in [0, T]$,

$$\begin{aligned} & \|\Delta(t, \cdot)\|_{L^2(0,L)} \\ & \leq \left[\int_0^T \left(\Phi_1 \left(\|\bar{\alpha}_x\|_{L^2(0,L)} + \|z_x\|_{L^2(0,L)} + \|\bar{z}_x\|_{L^2(0,L)} \right) \|(z - \bar{z})_x\|_{L^2(0,L)} \right. \right. \\ & \quad \left. \left. + |\Phi_1 - \Phi_2| \|\bar{z} + \bar{\alpha}\|_{L^2(0,L)}^{\frac{1}{2}} \|(\bar{z} + \bar{\alpha})_x\|_{L^2(0,L)}^{\frac{3}{2}} \right) dt \right]^2 \\ & \quad \times \exp \left[C \left(1 + \left\| \sqrt{\Phi_1} \alpha_x \right\|_{L^2(L^2)}^2 + \left\| \sqrt{\Phi_1} \bar{\alpha}_x \right\|_{L^2(L^2)}^2 + \left\| \sqrt{\Phi_1} z_x \right\|_{L^2(L^2)}^2 \right) \right]. \end{aligned}$$

This completes the proof of Lemma 3.4. ■

Now we are in a position to prove the following proposition on the global Lipschitzianity of the map $R(t)$. With our notation, we have

$$R(t)y_0 - R(t)\bar{y}_0 = \alpha(t, \cdot) - \bar{\alpha}(t, \cdot) = \Delta.$$

Proposition 3.4 *Let $T > 0$. There exists $\varepsilon_0 \in (0, 1]$ and $\tilde{C} : (0, \varepsilon_0] \rightarrow (0, +\infty)$ such that*

$$\|\Delta\|_{L^2(0,L)} \leq \tilde{C}(\varepsilon) \|y_0 - \bar{y}_0\|_{L^2(0,L)}, \quad \forall \bar{y}_0, y_0 \in L^2(0, L), \forall t \in [0, T], \forall \varepsilon \in (0, \varepsilon_0], \quad (3.53)$$

$$\tilde{C}(\varepsilon) \rightarrow 0 \text{ as } \varepsilon \rightarrow 0^+. \quad (3.54)$$

Proof. Let

$$\Delta_{\max} := \sup_{\substack{t \in [0, T] \\ \varepsilon \in (0, 1]}} \|\Delta(t, \cdot)\|_{L^2(0,L)}.$$

Let us point out that, by Lemma 3.3, $\Delta_{\max} < +\infty$. We claim that

$$\Delta_{\max} \leq \varepsilon C \left(\|y_0 - \bar{y}_0\|_{L^2(0,L)} + \Delta_{\max} \right), \quad \forall \bar{y}_0, y_0 \in L^2(0,L). \quad (3.55)$$

Then let ε be small enough such that $1 - \varepsilon C > 0$, we obtain

$$\Delta_{\max} \leq \frac{\varepsilon C}{1 - \varepsilon C} \|y_0 - \bar{y}_0\|_{L^2(0,L)}, \quad \forall \bar{y}_0, y_0 \in L^2(0,L).$$

Consequently, we get

$$\|\Delta\|_{L^2(0,L)} \leq \frac{\varepsilon C}{1 - \varepsilon C} \|y_0 - \bar{y}_0\|_{L^2(0,L)}, \quad \forall t \in [0, T], \quad \forall \bar{y}_0, y_0 \in L^2(0,L),$$

and the result follows. Hence, in order to prove Proposition 3.4, we only need to show that (3.55) holds in the following cases:

- (i) $\|\bar{y}_0\|_{L^2(0,L)}, \|y_0\|_{L^2(0,L)} \geq \varepsilon$, and $\|\bar{y}\|_{L^2(0,L)}, \|y\|_{L^2(0,L)} \geq \varepsilon, \forall t \in [0, T]$;
- (ii) $\|\bar{y}_0\|_{L^2(0,L)}, \|y_0\|_{L^2(0,L)} \geq \varepsilon$, there exists $\tau \in [0, T]$ such that $\|\bar{y}(\tau, \cdot)\|_{L^2(0,L)} = \varepsilon$, and $\|y(t, \cdot)\|_{L^2(0,L)} \geq \varepsilon, \forall t \in [0, T]$;
- (iii) $\|\bar{y}_0\|_{L^2(0,L)}, \|y_0\|_{L^2(0,L)} \geq \varepsilon$, there exists $\tau, \varsigma \in [0, T], \varsigma > \tau$, such that $\|\bar{y}(\tau, \cdot)\|_{L^2(0,L)} = \varepsilon$, and $\|y(\varsigma, \cdot)\|_{L^2(0,L)} = \varepsilon$;
- (iv) $\|\bar{y}_0\|_{L^2(0,L)} \leq \varepsilon$ and $\|y\|_{L^2(0,L)} \geq \varepsilon, \forall t \in [0, T]$;
- (v) $\|\bar{y}_0\|_{L^2(0,L)} \leq \varepsilon, \|y_0\|_{L^2(0,L)} \geq \varepsilon$, and there exists $\tau \in [0, T]$ such that $\|y(\tau, \cdot)\|_{L^2(0,L)} = \varepsilon$;
- (vi) $\|\bar{y}_0\|_{L^2(0,L)} \leq \varepsilon, \|y_0\|_{L^2(0,L)} \leq \varepsilon$.

By Lemma 3.4, for every $t \in [0, T]$, we have

$$\begin{aligned} & \|\Delta(t, \cdot)\|_{L^2(0,L)} \\ & \leq \int_0^t \left[\Phi_1 \left(\|\bar{\alpha}_x\|_{L^2(0,L)} + \|z_x\|_{L^2(0,L)} + \|\bar{z}_x\|_{L^2(0,L)} \right) \|(z - \bar{z})_x\|_{L^2(0,L)} \right. \\ & \quad \left. + |\Phi_1 - \Phi_2| \|\bar{z} + \bar{\alpha}\|_{L^2(0,L)}^{\frac{1}{2}} \|(\bar{z} + \bar{\alpha})_x\|_{L^2(0,L)}^{\frac{3}{2}} \right] dt \\ & \quad \times \exp \left[C \left(1 + \left\| \sqrt{\Phi_1} \alpha_x \right\|_{L^2(L^2)}^2 + \left\| \sqrt{\Phi_1} \bar{\alpha}_x \right\|_{L^2(L^2)}^2 + \left\| \sqrt{\Phi_1} z_x \right\|_{L^2(L^2)}^2 \right) \right]. \end{aligned} \quad (3.56)$$

Furthermore, by using Hölder's inequality and Lemma 2.2, we have

$$\begin{aligned} & \int_0^t \Phi_1 \left(\|\bar{\alpha}_x\|_{L^2(0,L)} + \|z_x\|_{L^2(0,L)} + \|\bar{z}_x\|_{L^2(0,L)} \right) \|(z - \bar{z})_x\|_{L^2(0,L)} dt \\ & \leq \|(z - \bar{z})_x\|_{L^2(L^2)} \left(\|\Phi_1 \bar{\alpha}_x\|_{L^2(L^2)} + \|\Phi_1 z_x\|_{L^2(L^2)} + \|\Phi_1 \bar{z}_x\|_{L^2(L^2)} \right) \\ & \leq C \|y_0 - \bar{y}_0\|_{L^2(0,L)} \left(\|\Phi_1 \bar{\alpha}_x\|_{L^2(L^2)} + \|\Phi_1 z_x\|_{L^2(L^2)} + \|\Phi_1 \bar{z}_x\|_{L^2(L^2)} \right). \end{aligned} \quad (3.57)$$

Applying the mean value theorem, noticing that $\|\alpha - \bar{\alpha}\|_{L^2(0,L)} \leq \Delta_{\max}, \forall t \in [0, T]$, and by Lemma 2.1,

$$\|z - \bar{z}\|_{L^2(0,L)} \leq \|y_0 - \bar{y}_0\|_{L^2(0,L)}, \quad \forall t \in [0, T],$$

we get

$$\begin{aligned} |\Phi_1 - \Phi_2| &\leq |\Phi'_\varepsilon(\theta)| \left| \|z + \alpha\|_{L^2(0,L)} - \|\bar{z} + \bar{\alpha}\|_{L^2(0,L)} \right| \\ &\leq |\Phi'_\varepsilon(\theta)| \|(z + \alpha) - (\bar{z} + \bar{\alpha})\|_{L^2(0,L)} \\ &\leq |\Phi'_\varepsilon(\theta)| \left(\|z - \bar{z}\|_{L^2(0,L)} + \|\alpha - \bar{\alpha}\|_{L^2(0,L)} \right) \\ &\leq |\Phi'_\varepsilon(\theta)| \left(\|y_0 - \bar{y}_0\|_{L^2(0,L)} + \Delta_{\max} \right), \end{aligned} \quad (3.58)$$

where

$$\theta = \theta(t) \in \left(\min\{\|z + \alpha\|_{L^2(0,L)}, \|\bar{z} + \bar{\alpha}\|_{L^2(0,L)}\}, \max\{\|z + \alpha\|_{L^2(0,L)}, \|\bar{z} + \bar{\alpha}\|_{L^2(0,L)}\} \right).$$

Thus, combining (3.56), (3.57) and (3.58), we arrive at

$$\begin{aligned} &\|\Delta(t, \cdot)\|_{L^2(0,L)} \\ &\leq \left(C \|y_0 - \bar{y}_0\|_{L^2(0,L)} \left(\|\Phi_1 \bar{\alpha}_x\|_{L^2(L^2)} + \|\Phi_1 z_x\|_{L^2(L^2)} + \|\Phi_1 \bar{z}_x\|_{L^2(L^2)} \right) \right. \\ &\quad \left. + \left(\|y_0 - \bar{y}_0\|_{L^2(0,L)} + \Delta_{\max} \right) \int_0^T |\Phi'_\varepsilon(\theta)| \|\bar{z} + \bar{\alpha}\|_{L^2(0,L)}^{\frac{1}{2}} \|(\bar{z} + \bar{\alpha})_x\|_{L^2(0,L)}^{\frac{3}{2}} dt \right) \\ &\quad \times \exp \left(C \left(1 + \left\| \sqrt{\Phi_1} \alpha_x \right\|_{L^2(L^2)}^2 + \left\| \sqrt{\Phi_1} \bar{\alpha}_x \right\|_{L^2(L^2)}^2 + \left\| \sqrt{\Phi_1} z_x \right\|_{L^2(L^2)}^2 \right) \right). \end{aligned}$$

Consequently, we obtain

$$\begin{aligned} \Delta_{\max} &\leq \left[C \|y_0 - \bar{y}_0\|_{L^2(0,L)} \left(\|\Phi_1 \bar{\alpha}_x\|_{L^2(L^2)} + \|\Phi_1 z_x\|_{L^2(L^2)} + \|\Phi_1 \bar{z}_x\|_{L^2(L^2)} \right) \right. \\ &\quad \left. + \left(\|y_0 - \bar{y}_0\|_{L^2(0,L)} + \Delta_{\max} \right) \int_0^T |\Phi'_\varepsilon(\theta)| \|\bar{y}\|_{L^2(0,L)}^{\frac{1}{2}} \|\bar{y}_x\|_{L^2(0,L)}^{\frac{3}{2}} dt \right] \\ &\quad \times \exp \left(C \left(1 + \left\| \sqrt{\Phi_1} \alpha_x \right\|_{L^2(L^2)}^2 + \left\| \sqrt{\Phi_1} \bar{\alpha}_x \right\|_{L^2(L^2)}^2 + \left\| \sqrt{\Phi_1} z_x \right\|_{L^2(L^2)}^2 \right) \right). \end{aligned} \quad (3.59)$$

For case (i), by (3.22), we have $\Phi_1 = \Phi_2 = 0, \forall t \in [0, T]$, it follows directly from (3.56) that

$$\Delta_{\max} = 0.$$

For case (ii), by (3.22), we have

$$\Phi_1 \equiv 0, \quad \forall t \in [0, T]. \quad (3.60)$$

In view of Lemma 3.1, we have

$$\|\bar{y}(t, \cdot)\|_{L^2(0,L)} \geq \varepsilon, \quad \forall t \in [0, \tau], \quad (3.61)$$

and

$$\|\bar{y}(t, \cdot)\|_{L^2(0,L)} \leq \|\bar{y}(\tau, \cdot)\|_{L^2(0,L)} = \varepsilon, \quad \forall t \in [\tau, T]. \quad (3.62)$$

Consequently, it follows from (3.22) and (3.61) that

$$\Phi_2 \equiv 0, \quad \forall t \in [0, \tau].$$

From (3.3), (3.59), (3.60), (3.61) and (3.62), we get that

$$\Delta_{\max} \leq \exp(C) \left(\|y_0 - \bar{y}_0\|_{L^2(0,L)} + \Delta_{\max} \right) \int_{\tau}^T \frac{C}{\varepsilon} \varepsilon^{\frac{1}{2}} \|\bar{y}_x\|_{L^2(0,L)}^{\frac{3}{2}} dt. \quad (3.63)$$

From now on, we assume that $\varepsilon \in (0, \eta]$, where $\eta > 0$ be chosen as in Lemma 3.2. Thanks to Lemma 3.2 and (3.62), we have

$$\|\bar{y}_x(t, \cdot)\|_{L^2(0,L)} \leq \frac{C}{\sqrt{t-\tau}} \|\bar{y}(\tau, \cdot)\|_{L^2(0,L)} = \frac{C}{\sqrt{t-\tau}} \varepsilon, \quad \forall t \in [\tau, T]. \quad (3.64)$$

Replacing (3.64) into (3.63), we obtain

$$\begin{aligned} \Delta_{\max} &\leq \varepsilon C \left(\|y_0 - \bar{y}_0\|_{L^2(0,L)} + \Delta_{\max} \right) \int_{\tau}^T \frac{1}{(t-\tau)^{\frac{3}{4}}} dt \\ &\leq \varepsilon C \left(\|y_0 - \bar{y}_0\|_{L^2(0,L)} + \Delta_{\max} \right). \end{aligned}$$

For case (iii), by (3.22) and Lemma 3.1, we have

$$\Phi_1 = 0, \quad \forall t \in [0, \varsigma], \quad (3.65)$$

$$\Phi_2 = 0, \quad \forall t \in [0, \tau], \quad (3.66)$$

and (3.62) still holds. In particular,

$$\|\bar{y}(\varsigma, \cdot)\|_{L^2(0,L)} \leq \varepsilon. \quad (3.67)$$

It follows from Lemma 2.2, (3.65) and (3.67) that

$$\begin{aligned} \|\Phi_1 \bar{z}_x\|_{L^2(0,T;L^2(0,L))} &= \|\Phi_1 \bar{z}_x\|_{L^2(\varsigma,T;L^2(0,L))} \\ &\leq \|\bar{z}_x\|_{L^2(\varsigma,T;L^2(0,L))} \leq C \|\bar{y}(\varsigma, \cdot)\|_{L^2(0,L)} \leq \varepsilon C, \end{aligned} \quad (3.68)$$

$$\begin{aligned} \|\Phi_1 z_x\|_{L^2(0,T;L^2(0,L))} &= \|\Phi_1 z_x\|_{L^2(\varsigma,T;L^2(0,L))} \\ &\leq \|z_x\|_{L^2(\varsigma,T;L^2(0,L))} \leq C \|y(\varsigma, \cdot)\|_{L^2(0,L)} = \varepsilon C, \end{aligned} \quad (3.69)$$

and

$$\left\| \sqrt{\Phi_1} z_x \right\|_{L^2(0,T;L^2(0,L))} \leq \varepsilon C. \quad (3.70)$$

By Remark 3.1, (3.65), (3.67) and (3.68), we have

$$\begin{aligned}\|\Phi_1 \bar{\alpha}_x\|_{L^2(0,T;L^2(0,L))} &\leq \|\Phi_1 \bar{y}_x\|_{L^2(0,T;L^2(0,L))} + \|\Phi_1 \bar{z}_x\|_{L^2(0,T;L^2(0,L))} \\ &\leq \|\bar{y}_x\|_{L^2(\varsigma,T;L^2(0,L))} + \|\Phi_1 \bar{z}_x\|_{L^2(0,T;L^2(0,L))} \\ &\leq \varepsilon C,\end{aligned}\tag{3.71}$$

and

$$\left\| \sqrt{\Phi_1} \bar{\alpha}_x \right\|_{L^2(0,T;L^2(0,L))} \leq \varepsilon C.\tag{3.72}$$

Similarly, we obtain

$$\left\| \sqrt{\Phi_1} \alpha_x \right\|_{L^2(0,T;L^2(0,L))} \leq \varepsilon C.\tag{3.73}$$

Moreover, for this case, (3.64) still holds. Now it follows from (3.3), (3.59), (3.62), (3.64), (3.65), (3.66), (3.68) to (3.73) that

$$\begin{aligned}\Delta_{\max} &\leq C \left(C\varepsilon \|y_0 - \bar{y}_0\|_{L^2(0,L)} + \varepsilon \left(\|y_0 - \bar{y}_0\|_{L^2(0,L)} + \Delta_{\max} \right) \int_{\tau}^T \frac{1}{(t-\tau)^{\frac{3}{4}}} dt \right) \\ &\leq \varepsilon C \left(\|y_0 - \bar{y}_0\|_{L^2(0,L)} + \Delta_{\max} \right).\end{aligned}$$

For case (iv), by (3.22), we have $\Phi_1 \equiv 0, \forall t \in [0, T]$. It follows from (3.59) that

$$\Delta_{\max} \leq C \left(\|y_0 - \bar{y}_0\|_{L^2(0,L)} + \Delta_{\max} \right) \int_0^T |\Phi'(\theta)| \|\bar{y}\|_{L^2(0,L)}^{\frac{1}{2}} \|\bar{y}_x\|_{L^2(0,L)}^{\frac{3}{2}} dt.\tag{3.74}$$

By Lemma 3.1, we have

$$\|\bar{y}(t, \cdot)\|_{L^2(0,L)} \leq \|\bar{y}_0\|_{L^2(0,L)} \leq \varepsilon, \quad \forall t \in [0, T].\tag{3.75}$$

Moreover, thanks to Lemma 3.2, we have

$$\|\bar{y}_x(t, \cdot)\|_{L^2(0,L)} \leq \frac{C}{\sqrt{t}} \|\bar{y}_0\|_{L^2(0,L)} \leq \frac{C}{\sqrt{t}} \varepsilon, \quad \forall t \in [0, T].\tag{3.76}$$

Then it follows from (3.3), (3.74) to (3.76) that

$$\begin{aligned}\Delta_{\max} &\leq \varepsilon C \left(\|y_0 - \bar{y}_0\|_{L^2(0,L)} + \Delta_{\max} \right) \int_0^T \frac{1}{t^{\frac{3}{4}}} dt \\ &\leq \varepsilon C \left(\|y_0 - \bar{y}_0\|_{L^2(0,L)} + \Delta_{\max} \right).\end{aligned}$$

For case (v), similarly to case (iii), we have

$$\|\Phi_1 \bar{\alpha}_x\|_{L^2(0,T;L^2(0,L))} \leq \varepsilon C, \quad \|\Phi_1 z_x\|_{L^2(0,T;L^2(0,L))} \leq \varepsilon C,\tag{3.77}$$

$$\|\Phi_1 \bar{z}_x\|_{L^2(0,T;L^2(0,L))} \leq \varepsilon C, \quad \left\| \sqrt{\Phi_1} \alpha_x \right\|_{L^2(0,T;L^2(0,L))} \leq \varepsilon C,\tag{3.78}$$

$$\left\| \sqrt{\Phi_1} \bar{\alpha}_x \right\|_{L^2(0,T;L^2(0,L))} \leq \varepsilon C, \quad \left\| \sqrt{\Phi_1} z_x \right\|_{L^2(0,T;L^2(0,L))} \leq \varepsilon C,\tag{3.79}$$

Moreover, (3.75) and (3.76) still hold. Thanks to (3.3), (3.75) and (3.76), we have

$$\int_0^T |\Phi'(\theta)| \|\bar{y}\|_{L^2(0,L)}^{\frac{1}{2}} \|\bar{y}_x\|_{L^2(0,L)}^{\frac{3}{2}} dt \leq \varepsilon C \int_0^T \frac{1}{t^{\frac{3}{4}}} dt. \quad (3.80)$$

Then, by (3.59), (3.77) to (3.80), we obtain

$$\Delta_{\max} \leq \varepsilon C \left(\|y_0 - \bar{y}_0\|_{L^2(0,L)} + \Delta_{\max} \right).$$

For the last case (vi), (3.75)-(3.80) hold, and (3.55) follows. Above all, we have proved (3.55) for all the cases (i)-(vi), which completes the proof of Proposition 3.4.

■

3.2.2 Smoothness of the semigroup

Lemma 3.5 *Let $\varepsilon > 0$ and $T > 0$ be given. Then the nonlinear map $\mathcal{S}(t)$ defined by the unique solution of (3.1) is of class C^3 from $L^2(0, L)$ to $C([0, T]; L^2(0, L))$. Moreover, its derivative $\mathcal{S}^{(1)}$ at $y_0 \in L^2(0, L)$ is given by*

$$\mathcal{S}^{(1)}(y_0)(h) := \mathcal{K}^{(1)}(y)(h), \quad \forall h \in L^2(0, L), \quad (3.81)$$

where $\mathcal{K}^{(1)}(y)(h)$ is defined by the following system (3.82) with $y = \mathcal{S}(y_0)$.

$$\begin{cases} \Delta_t + \Delta_x + \Delta_{xxx} + \Phi'_\varepsilon(\|y\|_{L^2(0,L)}) \frac{\int_0^L y \Delta dx}{\|y\|_{L^2(0,L)}} y y_x + \Phi_\varepsilon(\|y\|_{L^2(0,L)})(y \Delta_x + \Delta y_x) = 0, \\ \Delta(t, 0) = \Delta(t, L) = 0, \\ \Delta_x(t, L) = 0, \\ \Delta(0, x) = h(x), \end{cases} \quad (3.82)$$

Proof. We refer to [44] and [1, Theorem 5.4] for a detailed argument in related circumstances.

■

3.2.3 Center manifold

Combining [42, Remark 2.3], Corollary 2.2 and Proposition 3.4, we are in a position to apply [42, Theorem 2.19] and [42, Theorem 2.28]. This gives, if $\varepsilon > 0$ is small enough which will be always assumed from now on, the existence of an invariant center manifold for (3.1) which is of class C^3 . (In fact this center manifold is called a center-unstable manifold in [42, Theorem 2.19]; however, in our situation, with the notations of [42], $P_2(t)$, $t \in \mathbb{R}$, are trivial projections and then the name of center manifold can be adopted: see [42, Remark 2.20].) More precisely, there exists a map $g : M \rightarrow M^\perp$ of class C^3 satisfying $g(0) = 0$ and $g'(0) = 0$, such that, if

$$G := \{x_1 + g(x_1) : x_1 \in M\},$$

then, for every $y_0 \in G$ and for every $t \in [0, +\infty)$, $\mathcal{S}(t)y_0 \in G$. Moreover, Theorem 1.1 holds, if (and only if),

$$\mathcal{S}(t)y_0 \rightarrow 0 \text{ as } t \rightarrow +\infty, \forall y_0 \in G \text{ such that } \|y_0\|_{L^2(0,L)} \text{ is small enough.} \quad (3.83)$$

(For this last statement, see (2.42) in [42].) We prove (3.83) in the next section.

4 Dynamic on the center manifold

In this section, we prove (3.83), which concludes the proof of Theorem 1.1.

Proof. Let $y_0 \in G$. Let, for $t \in [0, +\infty)$, $y(t)(x) := y(t, x) := (\mathcal{S}(t)y_0)(x)$. We write

$$y(t, x) = p(t)\varphi(x) + y^*(t, x), \quad (4.1)$$

where $\phi(x)$ is defined in (3.19) and $y^*(t, x) \in M^\perp$. By (3.19) and (3.20), we have, at least if $\|y(t)\|_{L^2(0,L)}$ is small enough which will be always assumed in this proof,

$$\begin{aligned} \frac{dp(t)}{dt} &= \int_0^L y_t(t, x) \varphi(x) dx = \int_0^L (-y_x - yy_x - y_{xxx}) \varphi(x) dx \\ &= \int_0^L y(t, x) \varphi_x(x) dx - \int_0^L y(t, x) y_x(t, x) \varphi(x) dx + \int_0^L y(t, x) \varphi_{xxx}(x) dx \\ &= \frac{1}{2} \int_0^L y^2(t, x) \varphi_x(x) dx. \end{aligned} \quad (4.2)$$

We can also obtain the system for $y^*(t, x)$ as the following

$$\begin{cases} y_t^* + y_x^* + (I - P)yy_x + y_{xxx}^* = 0, \\ y^*(t, 0) = y^*(t, L) = 0, \\ y_x^*(t, L) = 0. \end{cases} \quad (4.3)$$

It follows from (4.1) that

$$\begin{aligned} yy_x &= (p(t)\varphi(x) + y^*(t, x)) (p(t)\varphi_x(x) + y_x^*(t, x)) \\ &= p^2(t)\varphi(x)\varphi_x(x) + p(t)y^*(t, x)\varphi_x(x) + p(t)\varphi(x)y_x^*(t, x) + y^*(t, x)y_x^*(t, x). \end{aligned}$$

Consequently, we have

$$\begin{aligned} (I - P)yy_x &= p^2(t)\varphi(x)\varphi_x(x) + p(t)y^*(t, x)\varphi_x(x) + p(t)\varphi(x)y_x^*(t, x) + y^*(t, x)y_x^*(t, x) \\ &\quad - p^2(t)\varphi(x) \int_0^L \varphi^2(x)\varphi_x(x) dx - p(t)\varphi(x) \int_0^L y^*(t, x)\varphi(x)\varphi_x(x) dx \\ &\quad - p(t)\varphi(x) \int_0^L \varphi^2(x)y_x^*(t, x) dx - \varphi(x) \int_0^L \varphi(x)y^*(t, x)y_x^*(t, x) dx. \end{aligned} \quad (4.4)$$

By using (3.19) and integrations by parts, we have

$$\int_0^L \varphi^2(x) \varphi_x(x) dx = 0, \quad (4.5)$$

$$\int_0^L y^*(t, x) \varphi(x) \varphi_x(x) dx = -\frac{1}{2} \int_0^L \varphi^2(x) y_x^*(t, x) dx, \quad (4.6)$$

$$\int_0^L \varphi(x) y^*(t, x) y_x^*(t, x) dx = -\frac{1}{2} \int_0^L \varphi_x(x) (y^*(t, x))^2 dx. \quad (4.7)$$

It can be deduced from (4.4), (4.5), (4.6) and (4.7) that

$$\begin{aligned} (I - P)yy_x &= p^2(t)\varphi(x)\varphi_x(x) + p(t)y^*(t, x)\varphi_x(x) + p(t)\varphi(x)y_x^*(t, x) + y^*(t, x)y_x^*(t, x) \\ &\quad - \frac{1}{2}p(t)\varphi(x) \int_0^L \varphi^2(x)y_x^*(t, x)dx + \frac{1}{2}\varphi(x) \int_0^L \varphi_x(x) (y^*(t, x))^2 dx. \end{aligned} \quad (4.8)$$

According to the existence and smoothness of the center manifold, we can set

$$y^*(t, x) = a(x)p^2(t) + O(p^3(t)), \quad \text{as } |p(t)| \rightarrow 0. \quad (4.9)$$

Then, by using (4.3), (4.8) and by comparing the coefficients of $p^2(t)$, we obtain

$$\begin{cases} a_x(x) + a_{xxx}(x) + \varphi(x)\varphi_x(x) = 0, \\ a(0) = a(L) = 0, \\ a_x(L) = 0. \end{cases} \quad (4.10)$$

The solution of (4.10) is

$$a(x) = C_1 + C_2 \cos x - \frac{1}{3} \sin x + \frac{1}{6\pi} x \sin x + \frac{1}{36\pi} \cos(2x),$$

where

$$C_1 + C_2 = -\frac{1}{36\pi}. \quad (4.11)$$

Note that $y^*(t, x) \in M^\perp$, we have

$$\int_0^L a(x)\varphi(x)dx = 0,$$

i.e.

$$C_1 \frac{2\pi}{\sqrt{3\pi}} + C_2 \frac{-\pi}{\sqrt{3\pi}} + \frac{1}{6\pi} \times \frac{-3\pi}{2\sqrt{3\pi}} = 0,$$

which leads to

$$2\pi C_1 - \pi C_2 - \frac{1}{4} = 0. \quad (4.12)$$

Combining (4.11) and (4.12), we get

$$C_1 = \frac{2}{27\pi}, \quad C_2 = -\frac{11}{108\pi}.$$

Therefore,

$$a(x) = \frac{2}{27\pi} - \frac{11}{108\pi} \cos x - \frac{1}{3} \sin x + \frac{1}{6\pi} x \sin x + \frac{1}{36\pi} \cos(2x). \quad (4.13)$$

Combining (4.1), (4.2), (4.9) and (4.13), we obtain

$$\begin{aligned} \frac{dp(t)}{dt} &= \frac{1}{2} \int_0^L (p(t)\varphi(x) + a(x)p^2(t) + O(p^3(t)))^2 \varphi_x(x) dx \\ &= p^3(t) \int_0^L a(x)\varphi(x)\varphi_x(x) dx + O(p^4(t)) \\ &= \frac{p^3(t)}{3\pi} \left(-\frac{1}{3}\pi + \frac{1}{6\pi}\pi^2 \right) + O(p^4(t)) \\ &= -\frac{p^3(t)}{18} + O(p^4(t)), \quad \text{as } |p(t)| \rightarrow 0. \end{aligned}$$

This concludes the proof of (3.83) and the proof of Theorem 1.1. ■

Acknowledgement

We thank Thierry Gallay, Gérard Iooss, Lionel Rosier and Bing-Yu Zhang for their valuable advices during the preparation of this work.

References

- [1] Jerry L. Bona, Shu Ming Sun, and Bing-Yu Zhang. A nonhomogeneous boundary-value problem for the Korteweg-de Vries equation posed on a finite domain. Comm. Partial Differential Equations, 28(7-8):1391–1436, 2003.
- [2] Jerry L. Bona, Shu Ming Sun, and Bing-Yu Zhang. A non-homogeneous boundary-value problem for the Korteweg-de Vries equation posed on a finite domain. II. J. Differential Equations, 247(9):2558–2596, 2009.
- [3] Jerry L. Bona and Ragnar Winther. The Korteweg-de Vries equation, posed in a quarter-plane. SIAM J. Math. Anal., 14(6):1056–1106, 1983.
- [4] Joseph Boussinesq. Essai sur la théorie des eaux courantes. Mémoires présentés par divers savants à l'Acad. des Sci. Inst. Nat. France, XXIII, pp. 1–680, 1877.
- [5] Jack Carr. Applications of centre manifold theory, volume 35 of Applied Mathematical Sciences. Springer-Verlag, New York, 1981.
- [6] Eduardo Cerpa. Exact controllability of a nonlinear Korteweg-de Vries equation on a critical spatial domain. SIAM J. Control Optim., 46(3):877–899 (electronic), 2007.
- [7] Eduardo Cerpa. Control of a Korteweg-de Vries equation: a tutorial. Preprint, 2012.
- [8] Eduardo Cerpa and Jean-Michel Coron. Rapid stabilization for a Korteweg-de Vries equation from the left Dirichlet boundary condition. Preprint, hal.archives-ouvertes.fr:hal-00730190, 2012.

- [9] Eduardo Cerpa and Emmanuelle Crépeau. Boundary controllability for the nonlinear Korteweg-de Vries equation on any critical domain. Ann. Inst. H. Poincaré Anal. Non Linéaire, 26(2):457–475, 2009.
- [10] Eduardo Cerpa and Emmanuelle Crépeau. Rapid exponential stabilization for a linear Korteweg-de Vries equation. Discrete Contin. Dyn. Syst. Ser. B, 11(3):655–668, 2009.
- [11] Xu-Yan Chen, Jack K. Hale, and Bin Tan. Invariant foliations for C^1 semigroups in Banach spaces. J. Differential Equations, 139(2):283–318, 1997.
- [12] Jixun Chu, Jean-Michel Coron, and Peipei Shang. A note on the Gevrey property of the linear operator for KdV equation. In preparation.
- [13] Thierry Colin and Jean-Michel Ghidaglia. An initial-boundary value problem for the Korteweg-de Vries equation posed on a finite interval. Adv. Differential Equations, 6(12):1463–1492, 2001.
- [14] Jean-Michel Coron. Control and nonlinearity, volume 136 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2007.
- [15] Jean-Michel Coron and Emmanuelle Crépeau. Exact boundary controllability of a nonlinear KdV equation with critical lengths. J. Eur. Math. Soc. (JEMS), 6(3):367–398, 2004.
- [16] Emmanuelle Crépeau. Exact boundary controllability of the Korteweg-de Vries equation around a non-trivial stationary solution. Internat. J. Control, 74(11):1096–1106, 2001.
- [17] Klaus-Jochen Engel and Rainer Nagel. One-parameter semigroups for linear evolution equations, volume 194 of Graduate Texts in Mathematics. Springer-Verlag, New York, 2000. With contributions by S. Brendle, M. Campiti, T. Hahn, G. Metafune, G. Nickel, D. Pallara, C. Perazzoli, A. Rhandi, S. Romanelli and R. Schnaubelt.
- [18] Andrei V. Faminskii. Global well-posedness of two initial-boundary-value problems for the Korteweg-de Vries equation. Differential Integral Equations, 20(6):601–642, 2007.
- [19] Olivier Glass and Sergio Guerrero. Controllability of the Korteweg-de Vries equation from the right Dirichlet boundary condition. Systems Control Lett., 59(7):390–395, 2010.
- [20] Olivier Goubet and Jie Shen. On the dual Petrov-Galerkin formulation of the KdV equation on a finite interval. Adv. Differential Equations, 12(2):221–239, 2007.
- [21] Mariana Haragus and Gérard Iooss. Local bifurcations, center manifolds, and normal forms in infinite-dimensional dynamical systems. Universitext. Springer-Verlag London Ltd., London, 2011.
- [22] Justin Holmer. The initial-boundary value problem for the Korteweg-de Vries equation. Comm. Partial Differential Equations, 31(7-9):1151–1190, 2006.
- [23] Chaohua Jia and Bing-Yu Zhang. Boundary stabilization of the Korteweg-de Vries equation and the Korteweg-de Vries-Burgers equation. Acta Appl. Math., 118:25–47, 2012.
- [24] V. Komornik. Exact controllability and stabilization. RAM: Research in Applied Mathematics. Masson, Paris, 1994. The multiplier method.
- [25] V. Komornik, David. L. Russell, and Bing-Yu Zhang. Stabilisation de l'équation de Korteweg-de Vries. C. R. Acad. Sci. Paris, 312:841–843, 1991.
- [26] Diederik J. Korteweg and Gustav de Vries. On the change of form of long waves advancing in a rectangular canal, and on a new type of long stationary waves. Philos. Mag., 39(5):422–

- 443, 1895.
- [27] Eugene F. Kramer and Bing-Yu Zhang. Nonhomogeneous boundary value problems for the Korteweg-de Vries equation on a bounded domain. J. Syst. Sci. Complex., 23(3):499–526, 2010.
 - [28] Camille Laurent, Lionel Rosier, and Bing-Yu Zhang. Control and stabilization of the Korteweg-de Vries equation on a periodic domain. Comm. Partial Differential Equations, 35(4):707–744, 2010.
 - [29] Pierre Magal and Shigui Ruan. Center manifolds for semilinear equations with non-dense domain and applications to Hopf bifurcation in age structured models. Mem. Amer. Math. Soc., 202(951):vi+71, 2009.
 - [30] Claiton Petris Massarolo, Gustavo Alberto Perla Menzala, and Ademir Fernando Pazoto. On the uniform decay for the Korteweg-de Vries equation with weak damping. Math. Methods Appl. Sci., 30(12):1419–1435, 2007.
 - [31] Ademir Fernando Pazoto. Unique continuation and decay for the Korteweg-de Vries equation with localized damping. ESAIM Control Optim. Calc. Var., 11(3):473–486 (electronic), 2005.
 - [32] Ademir Fernando Pazoto and Lionel Rosier. Stabilization of a Boussinesq system of KdV-KdV type. Systems Control Lett., 57(8):595–601, 2008.
 - [33] Amnon Pazy. Semigroups of linear operators and applications to partial differential equations, volume 44 of Applied Mathematical Sciences. Springer-Verlag, New York, 1983.
 - [34] Gustavo Alberto Perla Menzala, Carlos F. Vasconcellos, and Enrique Zuazua. Stabilization of the Korteweg-de Vries equation with localized damping. Quart. Appl. Math., 60(1):111–129, 2002.
 - [35] Ivonne Rivas, Muhammad Usman, and Bing-Yu Zhang. Global well-posedness and asymptotic behavior of a class of initial-boundary-value problem of the Korteweg-de Vries equation on a finite domain. Math. Control Relat. Fields, 1(1):61–81, 2011.
 - [36] Lionel Rosier. Exact boundary controllability for the Korteweg-de Vries equation on a bounded domain. ESAIM Control Optim. Calc. Var., 2:33–55 (electronic), 1997.
 - [37] Lionel Rosier. Control of the surface of a fluid by a wavemaker. ESAIM Control Optim. Calc. Var., 10(3):346–380 (electronic), 2004.
 - [38] Lionel Rosier and Bing-Yu Zhang. Control and stabilization of the Korteweg-de Vries equation: recent progresses. J. Syst. Sci. Complex., 22(4):647–682, 2009.
 - [39] David. L. Russell and Bing-Yu Zhang. Smoothing and decay properties of solutions of the Korteweg-de Vries equation on a periodic domain with point dissipation. J. Math. Anal. Appl., 190(2):449–488, 1995.
 - [40] David. L. Russell and Bing-Yu Zhang. Exact controllability and stabilizability of the Korteweg-de Vries equation. Trans. Amer. Math. Soc., 348(9):3643–3672, 1996.
 - [41] Shu Ming Sun. The Korteweg-de Vries equation on a periodic domain with singular-point dissipation. SIAM J. Control Optim., 34(3):892–912, 1996.
 - [42] Nguyen Van Minh and Jianhong Wu. Invariant manifolds of partial functional differential equations. J. Differential Equations, 198(2):381–421, 2004.

- [43] Glenn F. Webb. Theory of nonlinear age-dependent population dynamics, volume 89 of Monographs and Textbooks in Pure and Applied Mathematics. Marcel Dekker Inc., New York, 1985.
- [44] Bing-Yu Zhang. Analyticity of solutions of the generalized Kortweg-de Vries equation with respect to their initial values. SIAM J. Math. Anal., 26(6):1488–1513, 1995.
- [45] Bing-Yu Zhang. Exact boundary controllability of the Korteweg-de Vries equation. SIAM J. Control Optim., 37(2):543–565, 1999.