

# On a variational theory of image amodal completion

Simon Masnou<sup>(1)</sup>  
Jean-Michel Morel<sup>(2)</sup>

August 23, 2005

## Abstract

We study a variational model for image *amodal completion*, i.e. the recovery of missing or damaged portions of a digital image by technics inspired by the well-known *amodal completion* process in human vision. Representing the image by a real-valued function and following an idea initially proposed in [28], our approach consists in finding a set of interpolating level lines which is optimal with respect to an appropriate criterion. We prove that this method is theoretically well-founded and we show the equivalence with a more classical approach based on a direct interpolation of the function.

## 1 Introduction

Digital images can be represented as grey level functions  $u(x, y)$  defined on a simple open subset of  $\mathbb{R}^2$  (usually a rectangle), which we call “image domain”. Of course, digital images are given as a discrete set of samples, but there are standard interpolation methods to get back to a smooth image, e.g. a trigonometric polynomial by Shannon interpolation (also called zero-padding [34]). There is no substantial difference between digital images and what we know of retinal images as rough data: in both cases, images are band-limited by an optical device and then sampled on a grid. So most questions in visual perception theory are easily translated into “computer vision” problems. This opens the way to a mathematical formalization and numerical experiments.

We shall deal in this paper with the counterpart in image processing of the “amodal completion” phenomenon that arises in human vision. This phenomenon has been widely studied by the phenomenologist Gaetano Kanizsa [24], who tried to give a consistent answer to one of the major enigmas of visual perception. Its understanding starts with the straightforward observation that the objects that we see in all day life are partially occulting each other, so that we only see parts of them. Georges Matheron [29] actually proved that, under a simple and realistic stochastic model of object occultation, the so called “dead leaves model”, we only see half of the objects in sight. To be more explicit, in any all day life image or photograph, whenever we distinguish some object, we only see, on the average, half of it. Mathematically precise versions of this statement can be found in the aforementioned book and in [23, 1]. So we only see (significant) pieces of all shapes we perceive, but these pieces change constantly according to our position with respect to all objects present in the scene. Our perception, however, does not even notice this problem: we perceive

---

<sup>(1)</sup>Lab. J.-L. Lions, Univ. Pierre et Marie Curie, B.C. 187, 75252 Paris Cedex 05, France, masnou@ann.jussieu.fr

<sup>(2)</sup>CMLA, ENS Cachan, 61 Av. du Prsident Wilson, 94235 Cachan, France, Jean-Michel.Morel@cmla.ens-cachan.fr

objects as though they were complete. The mechanism of this visual illusion was formalized by Kanizsa who formulated two geometric laws, under the names of “amodal completion” and “good continuation”.

The *amodal completion* principle applies when a perceived curve stops on another perceived curve, entailing the perception of a “T-junction”. The stopped curve is the leg of the T and the other curve is represented by the horizontal bar of the T. In such a situation, our perception tends to interpret the interrupted curve as the boundary of some object undergoing an occlusion. The leg of the T is then mentally extrapolated and, whenever possible, connected to another leg in front. This fact is illustrated in figure 1 and is called “amodal completion”. The connection of two T-legs in front obeys the “good continuation” principle, according to which the reconstructed amodal curve must be similar to the pieces of curve it interpolates (same direction, curvature, etc.)




Figure 1: T-junctions entail an amodal completion and a completely different interpretation of the image.

In figure 1 we see first four black butterfly-like shapes. By superposing to them four rectangles, the butterflies get amodally completed into disks. By adding instead to the butterflies a central white cross, the butterflies contribute to the perception of an amodal black big square. In all cases, the reconstructed amodal boundaries obey the *good continuation principle*, namely they are as homogeneous as possible to the visible parts (circles in one case, straight segments in the other case).

The work of Kanizsa and his collaborators was directed at proving that this completion mechanism can be fully formalized as an automatic geometric mechanism, which we may well call amodal completion algorithm. An amodal completion algorithm proceeds as follows: given a homogeneous grey level or color region  $\Omega$  bounded by a smooth Jordan curve, it detects “T-junctions”, namely points of  $\partial\Omega$  at which other contours stop, thus forming a typical T-shaped singularity. The leg of the T is understood as the boundary of some object in back, while the upper bar of the T is understood as the occulting contour. For instance, in the case of Figure 2, the boundary of the disk stops on the boundary of the square, thus forming two T-junctions. The amodal completion



Figure 2: A square above a disk seen by amodal completion: Kanizsa showed that the T-junctions were crucial for the perception of a full circle where only an arc of circle is actually present in the image.

phenomenon happens whenever two  $T$ -junctions turn out to face each other: in that case, both legs of both junctions tend to be perceptually connected by a “good”, smooth curve. How to decide the shape of this interpolating curve? Several models have been proposed (see an exhaustive review in [22]); most suggest more or less explicitly that the interpolating curve must offer a compromise between the good continuation of the visible edges and the length minimality. In other words, the curve must be as smooth and as short as possible. A generic enough model proposed in [31, 32] defines the completion curves as minimizers of the Euler elastica energy

$$E_2(\gamma) = \int_0^{\mathcal{L}(\gamma)} (\lambda + |\gamma''|^2(s)) ds$$

(given the positions of the extremities and the associated tangent vectors) or, by extension,

$$E_p(\gamma) = \int_0^{\mathcal{L}(\gamma)} (\lambda + |\gamma''|^p(s)) ds \quad (1)$$

where  $\gamma$  is a parameterization by length of the curve on  $[0, \mathcal{L}(\gamma)]$ . In particular,  $|\gamma'(s)| = 1$  a.e and  $\gamma''(s)$  coincides with the curvature. The parameters of the method are  $\lambda$  and  $p \geq 1$ . There is no particular reason to choose one or the other value for these parameters because they are highly context-dependent, i.e. depend on the position of T-junctions, the edge orientation, the convexity of the shape, etc., see [22, 33]. For the sake of simplicity, we shall assume that they are given once for all.

Let us see now how Kanizsa’s amodal completion principles can be translated into an image processing framework – we shall speak of image *amodal completion* – and can be used to tackle the problem of recovering missing parts in an image. This approach was initially developed in [28, 25, 26], following a previous work of Mumford and Nitzberg in a different context [32]. We shall present here a mathematical analysis of the image *amodal completion* problem that completes the partial theoretical results obtained in [28, 25, 26].

Let us first proceed to some mathematical notation. The occlusion shall be represented as an open, bounded and simply connected subset  $\Omega \subset \mathbb{R}^2$  with  $C^\infty$  boundary. For the sake of simplicity, we shall assume that the original image  $u_0$  is known on  $\mathbb{R}^2 \setminus \Omega$  but we could as well suppose that it is known only on  $\tilde{\Omega} \setminus \Omega$ , where  $\tilde{\Omega} \supset \Omega$  is open, bounded and has Lipschitz boundary. In addition, we assume that  $u_0$  is the trace on  $\mathbb{R}^2 \setminus \Omega$  of an *analytic* function  $U_0$ . This regularity assumption finds a rather natural justification in Shannon interpolation theory but is of course much stronger

than the usual  $U_0 \in \text{BV}(\mathbb{R}^2)$  that has been used in recent years to model the image geometry (see the excellent discussion on this topic in [30]). We actually make this assumption in order to simplify the proofs but it is worth noticing that the existence of an optimal solution to the image completion problem, as stated in Theorem 2, can be proved as well under the weaker hypothesis that  $U_0 \in \text{BV}(\mathbb{R}^2)$ .

Our second assumption on  $U_0$  is  $\mathcal{F}_p(U_0) < +\infty$  where the functional  $\mathcal{F}_p$  is defined as

$$\mathcal{F}_p(u) := \begin{cases} \int_{\Omega} |\nabla u| (1 + |\text{div}(\frac{\nabla u}{|\nabla u|})|^p) dx & \text{if } u \in C^2(\mathbb{R}^2) \\ +\infty & \text{if } u \in L^1(\mathbb{R}^2) \setminus C^2(\mathbb{R}^2), \end{cases}$$

with the convention that the integrand is zero wherever  $|\nabla u| = 0$ . Before justifying the use of this energy, we recall a well-known property of the curvature along level lines, namely that for almost every  $t \in \mathbb{R}$  and for every  $x \in \{u = t\}$ , the curvature  $\kappa(x)$  of the level line  $\{u = t\}$  at  $x$  satisfies

$$\kappa(x) = \text{div} \frac{\nabla u}{|\nabla u|}(x).$$

From this and a call to the change of variables formula, we get that for any  $u \in C^2(\mathbb{R}^2)$

$$\mathcal{F}_p(u) \equiv \int_{-\infty}^{+\infty} \int_{\Omega \cap \partial\{u \geq \lambda\}} (1 + |\kappa|^p) d\mathcal{H}^1 d\lambda$$

when both terms are finite. This leads us to define a broader version of  $\mathcal{F}_p$  as

$$F_p(u) = \int_{-\infty}^{+\infty} \int_{\Omega \cap \partial\{u \geq \lambda\}} (1 + |\kappa|^p) d\mathcal{H}^1 d\lambda, \quad (2)$$

this definition making of course sense only when, for almost every level  $\lambda$ , the restriction to  $\Omega$  of the level lines are a countable set of smooth enough curves.

Our regularity assumption  $\mathcal{F}_p(U_0) = F_p(U_0) < \infty$  implies that for almost every  $\lambda \in \mathbb{R}$ ,  $\partial\{U_0 \geq \lambda\}$  has finite  $E_p$  energy, thus the level lines of  $U_0$  have a “good continuation” behavior.

Following the model proposed in [28, 26], we can reinterpret Kanizsa’s amodal completion in a functional framework, where *all missing level lines of the image  $u_0 = U_0|_{\mathbb{R}^2 \setminus \Omega}$  have to be interpolated inside  $\Omega$  according to the good continuation principle*. To this aim, let us call “T-junction” every point  $x \in \partial\Omega$  where  $\nabla U_0$  does not vanish, which means that there is a level line passing by  $x$ . Let us parameterize the trace of this line on  $\mathbb{R}^2 \setminus \Omega$  near  $x$  as  $\gamma^1(t)$ ,  $t \in [-\varepsilon, 0]$ , with  $\gamma^1(0) = x$ . This is a first *T-junction leg*. This leg has to be matched to another one of the same level and arriving elsewhere at some  $y \in \partial U_0$ . Let us denote as  $\gamma^2(t)$ ,  $t \in [1, 1 + \varepsilon]$ , with  $\gamma^2(1) = y$  this second one and assume that both T-junctions are compatible, i.e.  $\det(\nabla U_0(x), (\gamma^1)'(0))$  and  $\det(\nabla U_0(y), (\gamma^2)'(0))$  have the same sign (see figure 3). This compatibility condition is necessary to ensure that we will not reconstruct a “twisted” level line that could not be considered as the level line of a function. Our problem is to connect  $\gamma^1$  with  $\gamma^2$  by a smooth curve which we shall denote by  $\gamma^1: [0, 1] \rightarrow \Omega$ , with the condition that the concatenated curve  $\tilde{\gamma}: t \in [-\varepsilon, 1 + \varepsilon] \rightarrow \tilde{\gamma}(t)$  coinciding with  $\gamma^1$  for  $-\varepsilon \leq t \leq 0$  and with  $\gamma^2$  for  $1 \leq t \leq 1 + \varepsilon$  is in  $W^{2,p}(-\varepsilon, 1 + \varepsilon)$  or, equivalently, that  $E_p(\tilde{\gamma}) < \infty$ .

We finally *define an amodal completion as a set of interpolating curves  $\gamma_x$  defined for all  $x$  of  $\partial\Omega$  which are T-junctions*. Each  $\gamma_x$  joins a junction  $x$  to a junction  $y$  (so that  $U_0(x) = U_0(y)$ ) and, up to a reparameterization,  $\gamma_x = \gamma_y$ . The interpolating curves must fill some requirements making them fit to become level lines, namely




Figure 3: Two T-junctions and a possible amodal completion.

- $\nabla U_0(x)$  and  $\nabla U_0(y)$  have the same orientation along the curve  $\gamma_x$  (see figure 3) ;
- if  $\gamma_x$  arrives at  $y$ , the curves  $\gamma_x$  and  $\gamma_y$  coincide up to reparameterization ;
- two curves  $\gamma_x$  and  $\gamma_y$  can meet only tangentially and never cross each other and this applies also when  $x = y$ .

We call  $\mathcal{D}_p$  the set of all amodal completions of  $U_0$  inside  $\gamma$ . With each curve  $\gamma_x$  of an amodal completion is associated a grey level  $U_0(x)$  and the non intersection constraint makes the curves  $\gamma_x$  fit to be level lines of a function  $u_\gamma$  that we shall call the *amodal completion of  $U_0$  inside  $\Omega$* . There is a standard way to construct such a function  $u_\gamma$  inside  $\Omega$  from  $\gamma$ , so that all level lines of  $u$  are contained in a countable or finite union of curves  $\gamma_x$ . The fact that there is not necessarily identity between level curves of the reconstructed  $u_\gamma$  and the  $\gamma_x$  is illustrated in Figure 4. In this figure, we see that two level lines of the same level coincide on some interval. Since the piece of curve where they coincide shows no contrast, the reconstructed function  $u_\gamma$  loses this part of the level curve. This possibility was pointed out in [6]. We shall call this singularity creating possibility the *curve gluing phenomenon*.




Figure 4: The *gluing phenomenon*: the curves of an amodal completion and the level lines of the associated function may have totally different structures.

Introducing the measure  $\mu := |\nabla U_0| \mathcal{H}^1 \llcorner \partial\Omega$ , the energy of the amodal completion is defined as the sum of all energies of all interpolated level lines, namely,

$$E_p(\gamma) = \frac{1}{2} \int_{\partial\Omega} E_p(\gamma_x) d\mu(x),$$

where  $E_p(\gamma_x)$  has been defined above in (1). The factor  $\frac{1}{2}$  recalls that we count the energy twice, since  $E(\gamma_x) = E(\gamma_y)$  when  $x$  and  $y$  are two matching T-junctions. A numerical theory and experiments for minimizing  $E_p$  when  $p = 1$  was developed in [25]. In that case, an absolute minimum

was theoretically and computationally attained. Actually there are two kinds of numerical theories dealing with the same problem, namely the ones which minimize either  $\mathcal{F}_p(u)$  or  $F_p(u)$  and the ones which minimize  $E_p(\gamma)$ . Now, the *gluing phenomenon* explains why we may expect that sometimes

$$E_p(\gamma) \neq F_p(u_\gamma). \quad (3)$$

We shall prove, however, that with any amodal completion  $\gamma$  and for every  $\epsilon > 0$  we can associate a function  $u_{\gamma,\epsilon}$  so that  $u_{\gamma,\epsilon} = U_0$  outside  $\Omega$ ,  $\|u_\gamma - u_{\gamma,\epsilon}\|_{L^1(\Omega)} \leq \epsilon$  and  $|E_p(\gamma) - F_p(u_{\gamma,\epsilon})| \leq \epsilon$  (see Lemma 7)

All the same, (3) suggests that we cannot just solve the amodal completion problem by looking for  $u$  minimizing  $F_p(u)$  with the constraint  $u = U_0$  on  $\mathbb{R}^2 \setminus \Omega$ . Indeed, there is not necessarily a solution to either problems

$$\min_{u=U_0 \text{ on } \mathbb{R}^2 \setminus \Omega} F_p(u). \quad (4)$$

or

$$\min_{u=U_0 \text{ on } \mathbb{R}^2 \setminus \Omega} \mathcal{F}_p(u). \quad (5)$$

We shall instead prove that there is a solution to

$$\min_{\gamma \in \mathcal{D}_p} E_p(\gamma) \quad (P_1)$$

The fact that (4) and (5) are ill-posed led the authors of [4] to adopt a slightly different strategy. First, in order to incorporate an explicit reference to the good continuation requirement, they define the energy on a domain slightly bigger than  $\Omega$ . More precisely, given an open and smooth subset  $\tilde{\Omega}$  such that  $\tilde{\Omega} \supset \Omega$ , the authors consider the energy  $\mathcal{F}_p^e$  defined by

$$\mathcal{F}_p^e(u) = \begin{cases} \int_{\tilde{\Omega}} |\nabla u| (1 + |\operatorname{div}(\frac{\nabla u}{|\nabla u|})|^p) dx & \text{if } u \in C^2(\mathbb{R}^2) \\ +\infty & \text{if } u \in L^1(\mathbb{R}^2) \setminus C^2(\mathbb{R}^2) \end{cases}$$

with the convention that the integrand is zero wherever  $|\nabla u| = 0$ . The minimization process is not performed directly on  $\mathcal{F}_p^e$ , for the same reason why (5) is ill-posed, but rather on the lower semicontinuous envelope  $\overline{\mathcal{F}_p^e}$  of  $\mathcal{F}_p^e$  defined by

$$\overline{\mathcal{F}_p^e}(u) := \inf \{ \liminf_{h \rightarrow \infty} \mathcal{F}_p^e(u_h) : u_h \rightarrow u \text{ in } L^1(\mathbb{R}^2) \}.$$

Then it is proved in [4] that the problem

$$\min_{u \equiv U_0 \text{ on } \mathbb{R}^2 \setminus \Omega} \overline{\mathcal{F}_p^e}(u)$$

is well-posed.

We should work with a different definition of the relaxed functional associated with  $\mathcal{F}_p$  in order to reintroduce the good continuation requirement that does not appear in  $\mathcal{F}_p$ . Given a function  $u \in L^1(\mathbb{R}^2)$  that coincides with  $U_0$  outside  $\Omega$ , we define

$$\overline{\mathcal{F}_p}(u) := \inf \{ \liminf_{h \rightarrow \infty} \mathcal{F}_p(u_h) : u_h \rightarrow u \text{ in } L^1(\mathbb{R}^2), u_h \equiv U_0 \text{ on } \mathbb{R}^2 \setminus \Omega \}$$

Of course, this relaxed functional is still the largest lower semicontinuous functional less than  $\mathcal{F}_p$ , when restricted to the class of functions that coincide with  $U_0$  outside  $\Omega$ . Under the crucial assumptions that  $U_0$  is smooth and  $\mathcal{F}_p(U_0) < \infty$ , all results of [4] remain true when particularized to the class of functions that coincide with  $U_0$  outside  $\Omega$  and we get that

$$\min_{u \equiv U_0 \text{ on } \mathbb{R}^2 \setminus \Omega} \overline{\mathcal{F}_p}(u) \quad (P'_2)$$

is well posed.

For every  $u \in L^1(\mathbb{R}^2)$  such that  $u = U_0$  on  $\mathbb{R}^2 \setminus \Omega$ , we can also define the relaxed functional associated with  $F_p$  as

$$\overline{F_p}(u) = \inf \{ \liminf_{h \rightarrow \infty} F_p(u_h) : u_h \rightarrow u \text{ in } L^1(\mathbb{R}^2), u_h = U_0 \text{ on } \mathbb{R}^2 \setminus \Omega \}$$

It will be established in Theorem 3 that

$$\min_{u \equiv U_0 \text{ on } \mathbb{R}^2 \setminus \Omega} \overline{F_p}(u) \quad (P_2)$$

is also well posed.

Our main results in this paper are, first, that problem  $(P_1)$  is well posed (Theorem 2) and, second, that problems  $(P_1)$  and  $(P_2)$  (Theorem 4) have the same minimal energies. We also prove that there is a solution  $u$  of  $(P_2)$  satisfying  $u = u_\gamma$ , where  $\gamma$  is a solution of  $(P_1)$ . Conversely, given any  $u$  minimizing  $(P_2)$ , there is an amodal completion minimizing  $E_p$  whose curves contain all level lines of  $u$ . The amodal completion problem therefore yields a very intuitive geometric interpretation  $(E_p)$  of a relaxed functional  $(\overline{F_p})$ .

We were not able to determine whether  $(P_2)$  and  $(P'_2)$  also have the same minimizers. We actually do not know if it is true or not that  $F_p(u) = \overline{F_p}(u)$  for any  $u \in \mathcal{S}_p$ , a class of – non necessarily smooth – functions in the domain of  $F_p$  (see Section 2). It is indeed unclear whether any  $u \in \mathcal{S}_p$  can be approximated by a sequence of *smooth* functions  $(u_h)$  such that  $\lim_{h \rightarrow \infty} F_p(u_h) = \overline{F_p}(u)$ .

We end this section with a brief description of the state-of-art relative to the subject of this paper.

The first adaptation of amodal completion's principles to image processing can be found in [32]: in order to reconstruct partially occluded objects, the authors propose to interpolate their boundaries *below* the occlusions using curves that minimize the Euler elastica energy  $\int (\lambda + \kappa^2) ds$ , where  $\kappa$  is the curvature.

This idea was adapted in [28, 25, 26] to the level lines framework in order to solve the problem of recovering missing areas in an image, following the strategy that we previously described. Figure 1 illustrates the kind of results that can be obtained with this approach. The bottom left image is the result of a *global* minimization of  $E_1$  by dynamic programming (see [28, 25, 26]) whereas the bottom right image is obtained by a *global* minimization of  $E_2$ , still by dynamic programming, among the collection of all amodal completions made of Euler spirals, i.e. curves whose curvature depends linearly on the arc-length [27]. The problem of recovering missing areas in an image is addressed in a completely different way in [10]. The proposed method is inspired by the technics employed by professional restorators for the restoration of old paintings. It consists in a progressive diffusion of the information from the boundary of the domain towards the interior by means of a partial differential equation that aims at transporting along the isophotes a specific criterion




Figure 5: Top: original image with missing area shown in white. Bottom left: after amodal completion by minimization of  $E_1$  (see the description of the algorithm in [28, 26]). Bottom right: after amodal completion by minimizing  $E_2$  in the class of amodal completions made of Euler spirals [27].

of image smoothness. The connections of this model – the so-called *inpainting* model – with the classical Navier-Stokes equation of fluid dynamics are shown in [9].

In [14], the authors propose a denoising/interpolation model based on the joint minimisation of a quadratic fidelity term outside the occulting domain and a total variation criterion within a domain slightly bigger than the occlusion (see also a variant of the equation associated with this model in [15]). The model proposed in [13] aims at recovering a piecewise smooth function inside the occlusion by minimizing the classical Mumford-Shah functional with the additional constraint that the discontinuity set, whenever it exists, has minimal Euler elastica energy (see also a recent implementation of this model in [20]). Finally, the authors of [21] introduce a numerical scheme of the fourth order nonlinear flow equation associated with  $\mathcal{F}_p$  in order to find its local minimizers.

A more sophisticated criterion is derived in [5], where the authors propose a joint interpolation of image intensity and level lines directions using a functional that can be seen as a clever relaxation of  $\mathcal{F}_p$ . The resulting model offer many advantages both from a theoretical and a practical viewpoint.

All these approaches are essentially dedicated to the reconstruction of the geometric information and usually perform badly for the interpolation of texture. Very recently, a new class of methods have appeared that perform very well in many situations. All these methods rely on a very simple



“copy-paste” procedure that was introduced for the first time in [18] in the context of texture synthesis. The first adaptations to image interpolation can be found in [12, 16]. They perform remarkably well in most situations, except when the information to recover require some large scale interpretation, which indicates that these methods could be advantageously combined with an approach like in [28].

Let us finally mention two recent variational models based on a linear decomposition of the image into a geometric component and a texture component and the use of two different reconstruction methods, one for each component. The decomposition/reconstruction process is performed either independently [11] or, more interestingly, jointly [19].

## 1.1 Anterior work, novelties

Whatever is being done here can be derived from anterior works in the particular case where  $u$  is the characteristic function of a set  $A$ . In that case, G. Bellettini, G. Dal Maso and M. Paolini in [6] and, later, G. Bellettini and L. Mugnai in [7, 8] studied the relaxation of

$$F_p(\chi_A) = \int_{\partial A} (1 + |\kappa|^p) d\mathcal{H}^1,$$

where  $\chi_A$  denotes the characteristic function of  $A \subset \mathbb{R}^2$ . In particular, it is shown in [7] that if  $A$  satisfies  $\overline{F}_p(\chi_A) < \infty$  then  $A$  essentially coincides with the interior set of a limit system  $\Gamma$  of curves and  $\overline{F}_p(\chi_A) = E_p(\Gamma)$ . In the particular case where  $\partial A$  is piecewise  $W^{2,p}$  with finitely many cusps then  $\Gamma$  consists in adding to  $\partial A$  an appropriate collection of smooth curves that connect the cusps pairwise (see also a representation with varifolds in [8]) and give them a “good continuation”, thus realizing a kind of “amodal completion” like in figure 4.

What we added to these arguments:

- the slight changes necessary to deal with the amodal completion problem (which amounts to treat the kind of Dirichlet condition given by T-junctions);
- as in the mentioned papers, we obtain a geometric characterization of the relaxation of  $F_p$ , that is we translate the abstract  $\overline{F}_p(u)$  into the intuitive  $E_p(\gamma)$ ;
- now, these new functionals deal with all level sets  $A_\lambda = \{u \geq \lambda\}$  together instead of just one. The extension is not trivial as one can judge from section 4;
- we provide an existence proof for a minimizing amodal completion for  $E_p$ . This problem was left open for a long time, with the exception of  $p = 1$  proved in [26];
- we show the equivalence between the minimization of  $E_p$  on curves and the minimization of  $\overline{F}_p$  on functions.

## 1.2 Reader’s guide

We introduce some notations and the definition of T-junctions in Sections 2, 2.1. We precisely define in Section 2.2 amodal completions and the amodal energy  $E_p$  (Definition 5). The main point is to impose the non intersection constraint on the curves of the amodal completion. This permits to uniquely define from an amodal completion  $\gamma$  a function  $u_\gamma$  so that, if  $E_p(\gamma)$  is finite and  $\gamma$  has

no contact,  $F_p(u_\gamma) = E_p(\gamma)$  (Theorem 1). In section 4, Theorem 2, we prove our first main result, namely the existence of a solution to the amodal completion problem

$$\min \{E_p(\gamma) : \gamma \in \mathcal{D}_p\}. \quad (P_1).$$

The solution  $\gamma$  to this problem yields an amodal completion image  $u_\gamma$  with bounded variation. In order to identify the bridges between the functional viewpoint and the amodal completion viewpoint, we prove in Lemma 6 that every amodal completion can be approximated by a sequence of amodal completions without contact, from which we construct a sequence of continuous functions  $(u_\epsilon)$  converging to  $u_\gamma$  and whose energy  $F_p(u_\epsilon)$  is arbitrarily close to  $E_p(\gamma)$  (Lemma 7). Conversely, from any  $u \in C^2$  one can define an amodal completion  $\gamma$ , obtained by a selection of the level lines of  $u$ , such that  $E_p(\gamma) \leq F_p(u) = \mathcal{F}_p(u)$  (Lemma 8).

As we have mentioned,  $(P_1)$  is well posed while  $(P_2)$  is not, so that  $(P_1)$  turns out to be a correct formalization of the amodal completion problem. Our main second result, in section 5 is of course the equivalence of the problem  $(P_1)$  with  $(P_2)$ . We prove this equivalence in Theorem 4 and show the close relationships between the minimizers of  $(P_1)$  and those of  $(P_2)$ . In particular, if  $\gamma$  minimizes  $(P_1)$  then  $u_\gamma$  minimizes  $(P_2)$  and  $\overline{F}_p(u_\gamma) = E_p(\gamma)$ . Conversely, if  $u$  is a minimizer of  $(P_2)$  then there exists an amodal completion  $\gamma_u$  such that  $\gamma_u$  is a minimizer of  $(P_1)$  and its associated function  $u_{\gamma_u}$  coincides with  $u$  almost everywhere.

## 2 Notations and definitions

We recall from the introduction that the occlusion will be represented as an open, bounded and simply connected subset  $\Omega \subset \mathbb{R}^2$  with  $C^\infty$  boundary. The original image is supposed to be known only outside  $\Omega$ . We assume that it is the trace on  $\mathbb{R}^2 \setminus \Omega$  of an analytic function  $U_0$ . The interpolation within  $\Omega$  being trivial if  $U_0$  is constant on  $\partial\Omega$ , we can exclude this case. Then it is a straightforward consequence of Sard Lemma and the coarea formula that there exists a non empty subset  $\Lambda \subset \mathbb{R}$  with  $\mathcal{H}^1(U_0(\mathbb{R}^2 \setminus \Omega) \setminus \Lambda) = 0$  such that, for all  $\lambda \in \Lambda$ ,

$(H_1)$   $\{U_0 = \lambda\}$  is an analytic curve of finite length;

$(H_2)$   $\mathcal{H}^0(\{U_0|_{\partial\Omega} = \lambda\}) < \infty$ ;

Observing that the function  $\lambda \mapsto |\{U_0|_{\partial\Omega} \geq \lambda\}|$  is monotone and therefore admits countably many discontinuities,  $\Lambda$  can be chosen so that for all  $\lambda \in \Lambda$ ,

$(H_3)$   $\{U_0|_{\partial\Omega} \geq \lambda\} = \lim_{\mu \rightarrow \lambda} \{U_0|_{\partial\Omega} \geq \mu\}$ ;

where the convergence is meant as the convergence in measure.

**Definition 1** *A pair  $(U_0, \Lambda)$  verifying conditions  $(H_1) - (H_3)$  is called an admissible occlusion.*

We recall that, given a function  $u$  of bounded variation (see [3] for a survey on BV functions), its level sets  $\{u \geq \lambda\}$  are sets of finite perimeter for almost every  $\lambda \in \mathbb{R}$  and we can define their reduced boundary as the set

$$\begin{aligned} \partial^* \{u \geq \lambda\} := \{x \in \mathbb{R}^2 : \nu_{\{u \geq \lambda\}}(x) := \lim_{r \downarrow 0} \frac{D\chi_{\{u \geq \lambda\}}}{|D\chi_{\{u \geq \lambda\}}|}(B_r(x)) \text{ exists} \\ \text{and satisfies } |\nu_{\{u \geq \lambda\}}(x)| = 1\}, \end{aligned}$$

i.e. the set of all points where a generalized inner normal to  $\{u \geq \lambda\}$  exists.

The space  $\mathcal{S}_p(\Omega)$  that we define below is the set of all functions  $u$  of bounded variation that coincide with  $U_0$  outside  $\Omega$  and such that, for almost every  $\lambda$ ,  $\partial^*\{u \geq \lambda\} \cap \Omega$  essentially coincides with a finite union of curves that all join two points of  $\partial\Omega$  and properly extend outside  $\Omega$ . Clearly,  $\mathcal{S}_p(\Omega)$  is the space of the functions that follow Kanizsa's good continuation principle.

**Definition 2** We call  $\mathcal{S}_p(\Omega)$  the space of all functions  $u \in \text{BV}(\mathbb{R}^2)$  such that  $u = U_0$  on  $\mathbb{R}^2 \setminus \Omega$  and for almost every  $\lambda \in U_0(\partial\Omega)$ ,  $\partial^*\{u \geq \lambda\} \cap \overline{\Omega}$  coincides, up to a  $\mathcal{H}^1$ -negligible set, with the trace of a finite union of curves  $\gamma_i^\lambda : [0, 1] \rightarrow \overline{\Omega}$ ,  $i = 0, \dots, n_\lambda$  with the following properties:

- $\gamma_i^\lambda(0), \gamma_i^\lambda(1) \in \partial\Omega$ ;
- $\gamma_i^\lambda \in \text{W}^{2,p}(0, 1)$  and  $|d\gamma_i^\lambda/dt|$  is constant almost everywhere on  $[0, 1]$ ;
- there exists an extension  $\gamma_i^{\lambda, \epsilon} \in \text{W}^{2,p}(-\epsilon, 1 + \epsilon)$  of  $\gamma_i^\lambda$  such that  $\gamma_i^{\lambda, \epsilon}([-\epsilon, 0])$  and  $\gamma_i^{\lambda, \epsilon}([1, 1 + \epsilon])$  are (possibly overlapping) subsets of  $\{y : U_0(y) = U_0(x)\} \cap (\mathbb{R}^2 \setminus \Omega)$  with positive length;
- $\forall i, j$ ,  $\gamma_i^\lambda$  and  $\gamma_j^\lambda$  may intersect but only tangentially and without crossing each other.

Then we define the functional  $F_p$  acting on  $L^1(\Omega)$  by

$$F_p(u) = \begin{cases} \int_{-\infty}^{+\infty} \int_{\Omega \cap \partial^*\{u \geq \lambda\}} (1 + |\kappa|^p) d\mathcal{H}^1 d\lambda & \text{if } u \in \mathcal{S}_p(\Omega) \\ +\infty & \text{if } u \in L^1(\Omega) \setminus \mathcal{S}_p(\Omega). \end{cases}$$

where, for almost every  $\lambda \in U_0(\partial\Omega)$ , we mean

$$\int_{\Omega \cap \partial^*\{u \geq \lambda\}} (1 + |\kappa|^p) d\mathcal{H}^1 = \sum_{i=0}^{n_\lambda} \int_{\gamma_i^\lambda} (1 + |\kappa|^p) d\mathcal{H}^1$$

In Figure 6 below, we show an example of a piecewise constant element  $u$  of  $\mathcal{S}_p$  such that  $F_p(u) < \infty$  but  $\mathcal{F}_p(u) = +\infty$ .




Figure 6: A piecewise constant function  $u \in \mathcal{S}_p(\Omega)$  such that  $F_p(u) < \infty$  but  $\mathcal{F}_p(u) = +\infty$

## 2.1 Defining T-junctions

**Definition 3 (Occlusion's boundary measure)** *An admissible occlusion  $\Omega$  is endowed with a boundary measure*

$$\mu := |\nabla U_0| \mathcal{H}^1 \llcorner \partial\Omega.$$

**Definition 4 (T-junctions)** *We call T-junction any element of the set*

$$\mathcal{T} := \{x \in \partial\Omega : \exists \lambda \in \Lambda, x \in \partial\{U_0 \geq \lambda\},$$

*and we shall denote by  $\mathcal{T}_\lambda$  the set of all T-junctions associated with the level  $\lambda \in \mathbb{R}$ , i.e.*

$$\mathcal{T}_\lambda := \{x \in \mathcal{T} : U_0(x) = \lambda\}.$$

**Proposition 1**  *$\mu$ -almost every  $x \in \partial\Omega$  is a T-junction, i.e.  $\mu(\partial\Omega \setminus \mathcal{T}) = 0$ .*

PROOF: By the coarea formula for Lipschitz functions,

$$|DU_0|(\partial\Omega \setminus \mathcal{T}) = \int_{-\infty}^{+\infty} \mathcal{H}^0((\partial\Omega \setminus \mathcal{T}) \cap \{U_0 = t\}) dt.$$

Since, for almost all  $t$ ,

$$\mathcal{H}^0((\partial\Omega \setminus \mathcal{T}) \cap \{U_0 = t\}) = 0$$

by definition of  $\mathcal{T}$ , we conclude that  $|DU_0|(\partial\Omega \setminus \mathcal{T}) = 0$  and the proposition follows.  $\square$

## 2.2 Defining amodal completions and their “amodal energy”

**Definition 5 (Amodal completion on  $\Omega$  associated with  $U_0$ )** *We call amodal completion of class  $W^{2,p}$  associated with  $U_0$  a map  $\gamma$  from  $\mathcal{T}$  to  $W^{2,p}([0,1], \mathbb{R}^2)$  that associates with  $x \in \mathcal{T}$  a function  $\gamma_x$  that describes a curve in  $\overline{\Omega}$  with distinct endpoints on  $\partial\Omega$ . In addition:*

- *for  $\mu$ -almost every  $x \in \mathcal{T}$ , there exists a  $W^{2,p}$  parameterization of  $\gamma_x$  (still denoted by  $\gamma_x$ ) on  $[0,1]$  with positive and constant velocity. The endpoints conditions rewrite  $\gamma_x(0), \gamma_x(1) \in \mathcal{T}$  and  $\gamma_x(0) \neq \gamma_x(1)$ ;*
- *if two curves  $\gamma_x$  and  $\gamma_y$  have a common endpoint then  $\gamma_x(s) = \gamma_y(s)$  for every  $s \in [0,1]$ , or  $\gamma_x(s) = \gamma_y(1-s)$  for every  $s \in [0,1]$ ;*
- *each curve may have tangential self-contacts but without crossing;*
- *two curves may intersect tangentially but without crossing;*
- *for  $\mu$ -almost every  $x \in \mathcal{T}$  there exists an extension  $\gamma_x^\varepsilon \in W^{2,p}[-\varepsilon, 1+\varepsilon]$  of  $\gamma_x$  such that  $\gamma_x^\varepsilon([-\varepsilon, 0])$  and  $\gamma_x^\varepsilon([1, 1+\varepsilon])$  are (possibly overlapping) subsets of  $\{y : U_0(y) = U_0(x)\} \cap (\mathbb{R}^2 \setminus \Omega)$  with positive length and the orientation of  $\nabla U_0$  on these subsets can be continuously extended to the whole curve  $\gamma_x$  (see Figure 3).*

*The set of all such amodal completions will be denoted by  $\mathcal{D}_p$ .*

**Remark 1** Since, for every  $(x, \lambda) \in \mathcal{T}$ ,  $|\gamma'_x(t)|$  is assumed to be constant almost everywhere on  $[0, 1]$ , the arc-length parameter of the curve is  $s(t) = t\mathcal{L}(\gamma_x)$ , where  $\mathcal{L}(\gamma_x)$  denotes the curve total length. We let  $\tilde{\gamma}_x$  represent the curve by arc-length. Clearly, for every  $s \in [0, \mathcal{L}(\gamma_x)]$ ,  $\tilde{\gamma}_x(s) = \gamma_x(s/\mathcal{L}(\gamma_x))$ . Therefore

$$\tilde{\gamma}_x''(s) = \frac{\gamma_x''(t)}{[\mathcal{L}(\gamma_x)]^2}$$

Now, it is well known that the curvature along the curve satisfies, as a function of arc-length,

$$\kappa(s) = \gamma_x''(s)$$

and we deduce

$$\int_0^{\mathcal{L}(\gamma_x)} (1 + |\tilde{\gamma}_x''(s)|^p) ds = \int_0^{\mathcal{L}(\gamma_x)} (1 + |\kappa|^p) ds = \int_0^1 (|\gamma'_x(t)| + [\mathcal{L}(\gamma_x)]^{1-2p} |\gamma_x''(t)|^p) dt \quad (6)$$

Assuming that  $\gamma_x \in W^{2,p}(0, 1)$  is therefore equivalent to saying that  $\tilde{\gamma}_x$  belongs to  $W^{2,p}(0, \mathcal{L}(\gamma_x))$  and that  $\tilde{\gamma}_x$  has finite  $E_p$  energy.

**Definition 6 (Amodal completion without contact)** *An amodal completion  $\gamma \in \mathcal{D}_p$  is said to be without contact if:*

- for  $\mu$ -almost every  $x \in \partial\Omega$ ,  $\gamma_x$  is simple and  $(\gamma_x) \cap \partial\Omega = \{\gamma_x(0), \gamma_x(1)\}$ ;
- for  $\mu$ -almost every  $x, y \in \partial\Omega$  such that  $x \neq y$ ,  $(\gamma_x) \cap (\gamma_y) = \emptyset$ .

**Definition 7** *We define the amodal energy of an amodal completion  $\gamma$  as*

$$E_p(\gamma) = \frac{1}{2} \int_{\partial\Omega} E_p(\gamma_x) d\mu(x)$$

where

$$E_p(\gamma_x) = \int_0^{\mathcal{L}(\gamma_x)} (1 + |\gamma_x''|^p(s)) ds$$

### 3 From amodal completions to functions, and back

**Theorem 1 (Function associated with an amodal completion)** *Let  $(U_0, \Lambda)$  be an admissible occlusion. Any amodal completion  $\gamma$  on  $\Omega$  of class  $W^{2,p}$  satisfying  $E_p(\gamma) < \infty$  can be associated with a function  $u_\gamma \in \text{BV}(\mathbb{R}^2)$  such that  $u_\gamma \equiv U_0$  on  $\mathbb{R}^2 \setminus \Omega$  and, for almost every  $\lambda \in \mathbb{R}$ ,  $\partial^* \{u_\gamma \geq \lambda\} \cap \overline{\Omega} \subset \bigcup_{x \in \mathcal{I}_\lambda} \gamma_x$  up to a  $\mathcal{H}^1$ -negligible set. If, in addition,  $\gamma$  has no contact, then  $u_\gamma \in \mathcal{S}_p(\Omega)$  and  $F_p(u_\gamma) = E_p(\gamma)$ .*

**PROOF:** The proof is essentially based on a straightforward filling up algorithm, permitting to define uniquely from the amodal curves at level  $\lambda$  a set  $A_\lambda$  bounded by them and  $\partial\Omega$ . This set will be the  $\lambda$ -upper level set of  $u_\lambda$  inside  $\Omega$ . A consistency check must then be performed, namely that  $\mu > \lambda \Rightarrow A_\mu \subset A_\lambda$ .

**Step 1: From amodal completion curves at level  $\lambda$  to a level set  $A_\lambda$ .**

$\partial\Omega$  is provided with an orientation so that we can talk of arc intervals  $[x, y] \subset \partial\Omega$  without ambiguity. Let  $\lambda \in \Lambda$  such that  $\mathcal{T}_\lambda$  be not empty. By definition of  $\Lambda$ , it is also finite. The upper

level set of  $U_0$  on  $\partial\Omega$ ,  $\{x \in \partial\Omega, U_0(x) \geq \lambda\}$  is a finite union of disjoint arcs of  $\partial\Omega$ . Let us call  $[x_1, x_2]$  any of these arcs and set (see Figure 7)

$$x_3 = \begin{cases} \gamma_{x_2}(1) & \text{if } x_2 = \gamma_{x_2}(0), \\ \gamma_{x_2}(0) & \text{if } x_2 = \gamma_{x_2}(1). \end{cases}$$

Take for  $x_4 \in \mathcal{T}_\lambda$  the unique point such that  $[x_3, x_4]$  is a connected component of the upper level set  $\{x \in \partial\Omega, U_0(x) \geq \lambda\}$ . We can iterate this construction and after a finite number of steps we get a series of intervals  $[x_{2i+1}, x_{2i+2}]$ ,  $i = 0, \dots, j$  such that  $x_{2j+1} = x_1$ . Since each T-junction is the tip of a single amodal curve, no shorter cycle is possible in the mentioned sequence. Besides, since the arcs  $[x_{2i+1}, x_{2i+2}]$  are disjoint and the curves  $\gamma_{x_{2i+2}}$  cannot cross, we can concatenate them all into a rectifiable image of the circle into the plane, with no crossing but possibly self-contacts. We call  $\Gamma_1$  this generalized Jordan curve. Using the usual index with respect to a curve, we define the interior  $\Omega_1$  of  $\Gamma_1$  as the set of points with index 1. Notice that, by construction,  $\Omega_1$  is contained in  $\Omega$  and  $\Omega_1$  is not empty due to our assumption  $(H_3)$  at the beginning of Section 2. In addition, every point of  $[x_1, x_2]$  is the limit of a sequence of points in  $\Omega$  with index one.

This construction can be iterated until all T-junctions at level  $\lambda$  have been exhausted. The successive generalized Jordan curves  $\Gamma_1, \dots, \Gamma_k$  thus obtained do not cross. Thus, the sets  $\Omega_1, \dots, \Omega_k$  are disjoint.

We finally define

$$A_\lambda = \bigcup_{h=1}^k \Omega_h \quad (7)$$

and remark that, by construction,

$$\partial A_\lambda \subset \bigcup_{h=1}^k \Gamma_h \subset \partial\Omega \cup \bigcup_{x \in \mathcal{T}_\lambda} \gamma_x, \quad (8)$$

and

$$\mathcal{H}^1(\{x \in \partial\Omega : U_0(x) \geq \lambda\} \setminus \partial A_\lambda) = 0.$$

Besides, since each curve  $\Gamma_h$  is a finite union of  $C^1$  curves that do not cross each other, it follows that

$$\partial^* A_\lambda = \partial A_\lambda \quad \text{up to a } \mathcal{H}^1\text{-negligible set.} \quad (9)$$

The same construction is performed for every  $\lambda \in \Lambda$  (recall that  $\mathcal{H}^1(U_0(\partial\Omega) \setminus \Lambda) = 0$ ). Then we let  $A_\lambda = \Omega$  for every  $\lambda < \min_{x \in \partial\Omega} U_0(x)$  and  $A_\lambda = \emptyset$  for every  $\lambda > \max_{x \in \partial\Omega} U_0(x)$ , in order to ensure that we have associated a set  $A_\lambda$  with almost every  $\lambda \in \mathbb{R}$ .

Let us prove now that for any  $\lambda, \mu \in \Lambda$ ,

$$\lambda \leq \mu \Rightarrow A_\lambda \supset A_\mu \quad (\text{up to a Lebesgue negligible set})$$

Let  $\Gamma_1^\mu = \partial\Omega_1^\mu$  be one of the generalized Jordan curve defining  $A_\mu$  and  $[y_1, y_2]$  its first interval. By the inclusion of upper level sets property, this arc is contained in  $\{x \in \partial\Omega : U_0(x) \geq \lambda\}$  and therefore in some maximal interval of this set which we denote by  $[x_1, x_2]$ . Consider  $\Gamma_1^\lambda$ , the unique generalized Jordan curve in the preceding construction containing  $[x_1, x_2]$ . The curves  $\Gamma_1^\lambda$  and  $\Gamma_1^\mu$  do not cross. Indeed, their intersections with  $\partial\Omega$  are nested and their other parts are amodal

completion curves which do not cross each other. Thus, their associated sets  $\Omega_1^\lambda$  and  $\Omega_1^\mu$  are either disjoint, or  $\Omega_1^\mu \subset \Omega_1^\lambda$ . Now, the first possibility is ruled out because  $[y_1, y_2] \subset [x_1, x_2]$  and every point of  $[y_1, y_2]$  (resp.  $[x_1, x_2]$ ) is the limit of a sequence of points in  $\Omega$  with index 1 with respect to  $\Gamma_1^\mu$  (resp.  $\Gamma_1^\lambda$ ). Therefore, we have proved that  $\Omega_1^\mu \subset \Omega_1^\lambda$  and, by extension, that

$$\forall \lambda, \mu \in \Lambda, \quad \lambda \leq \mu \Rightarrow A_\lambda \supset A_\mu.$$

The same result is obviously true whenever  $\lambda$  or  $\mu$  are either less than  $\inf_{x \in \partial\Omega} U_0(x)$  or larger than  $\sup_{x \in \partial\Omega} U_0(x)$  and we can conclude that

$$\text{for a.e. } \lambda, \mu \in \mathbb{R}, \quad \lambda \leq \mu \Rightarrow A_\lambda \supset A_\mu$$




Figure 7: Construction of the function associated with an amodal completion

### Step 2: From level sets to a function

We now have a nested family of measurable sets  $(A_\lambda) \subset \Omega$  defined for almost every  $\lambda \in \mathbb{R}$ , actually for all  $\lambda \in \Lambda \cup (\mathbb{R} \setminus U_0(\partial\Omega))$ . Let us see how they can generate an essentially unique function  $u_\gamma$  defined on  $\Omega$ . Under the notations of Lemma 1 below, let  $D$  be the set of discontinuity points of  $(A_\lambda)$ . Let  $D' \subset \Lambda \cup (\mathbb{R} \setminus U_0(\partial\Omega))$  be countable and dense and define for every  $x \in \Omega$

$$u_\gamma^\Omega(x) := \sup\{\lambda \in D' : x \in A_\lambda\}.$$

We now prove that  $\{u_\gamma^\Omega \geq \lambda\} = A_\lambda$  (up to a Lebesgue-negligible set) for any  $\lambda \notin D$ ,  $\lambda \in \Lambda \cup (\mathbb{R} \setminus U_0(\partial\Omega))$ . By definition of  $u_\gamma^\Omega$ ,

$$A_\eta \subset \{u_\gamma^\Omega \geq \lambda\} \subset A_\nu$$

for any  $\eta, \nu \in D'$ ,  $\eta > \lambda > \nu$ . Choose sequences  $\eta_h \downarrow \lambda$  and  $\gamma_h \uparrow \lambda$  in  $D'$ . Then, by Lemma 1,

$$A_\lambda = \{u_\gamma^\Omega \geq \lambda\} \quad (\text{up to a Lebesgue negligible set}). \quad (10)$$

In particular,  $\{u_\gamma^\Omega \geq \lambda\}$  is measurable for any  $\lambda \notin D$ ,  $\lambda \in \Lambda \cup (\mathbb{R} \setminus U_0(\partial\Omega))$ . By approximation, the same property extends to any real number  $\lambda$  and  $u_\gamma^\Omega$  is measurable.

The uniqueness of  $u_\gamma^\Omega$  follows by a similar argument: if two functions  $u_1, u_2$  are such that  $\{u_1 \geq \lambda\} = \{u_2 \geq \lambda\}$  (up to a Lebesgue negligible set) for a dense set of  $\lambda$ 's, then  $u_1 = u_2$  almost everywhere in  $\Omega$ .

### Step 3: Properties of the new function

First remark that it follows from (8) and the finiteness of  $\mathcal{T}_\lambda$  that for every  $\lambda \in \Lambda$ ,  $\mathcal{H}^1(\partial A_\lambda) \leq \mathcal{H}^1(\partial\Omega) + \frac{1}{2}(\sum_{x \in \mathcal{T}_\lambda} \mathcal{H}^1(\gamma_x)) < \infty$ . Thus  $A_\lambda$  has finite perimeter and, by (9) and (10), also  $\{u_\gamma^\Omega \geq \lambda\}$  has finite perimeter and its essential boundary satisfies

$$\partial^* \{u_\gamma^\Omega \geq \lambda\} = \partial A_\lambda \subseteq \partial\Omega \cup \bigcup_{x \in \mathcal{T}_\lambda} \gamma_x \quad \text{up to a } \mathcal{H}^1\text{-negligible set.} \quad (11)$$

In particular,  $\mathcal{H}^1(\partial^* \{u_\gamma^\Omega \geq \lambda\}) = \mathcal{H}^1(\partial A_\lambda)$ . Since  $A_\lambda = \emptyset$  for all  $\lambda \leq \inf_{x \in \partial\Omega} U_0(x)$  and  $A_\lambda = \Omega$  for all  $\lambda \geq \sup_{x \in \partial\Omega} U_0(x)$  and because  $\mathcal{H}^1(U_0(\partial\Omega) \setminus \Lambda) = 0$  we can conclude that  $\{u_\gamma^\Omega \geq \lambda\}$  has finite perimeter in  $\Omega$  for almost every  $\lambda \in \mathbb{R}$ . Then,

$$\int_{-\infty}^{+\infty} \mathcal{H}^1(\partial^* \{u_\gamma^\Omega \geq \lambda\} \cap \Omega) d\lambda \leq \frac{1}{2} \int_{-\infty}^{+\infty} (\sum_{x \in \mathcal{T}_\lambda} \mathcal{H}^1(\gamma_x)) d\lambda \leq \frac{1}{2} \int_{\partial\Omega} E_p(\gamma_x) d\mu(x)$$

It follows from Lemma 2 that  $u_\gamma^\Omega \in \text{BV}(\Omega)$ . In addition, since  $U_0 \in \text{BV}(\mathbb{R}^2 \setminus \Omega)$  and  $\partial\Omega$  is smooth, the usual results on properties of the trace operator in BV (see for instance Corollary 3.89 in [3]) imply that the function  $u_\gamma$  defined by

$$u_\gamma(x) = \begin{cases} u_\gamma^\Omega(x) & \text{on } \Omega \\ U_0(x) & \text{on } \mathbb{R}^2 \setminus \Omega \end{cases}$$

is in  $\text{BV}(\mathbb{R}^2)$ . In addition, it is a direct consequence of 11 that for almost every  $\lambda \in \mathbb{R}$ ,

$$\partial^* \{u_\gamma \geq \lambda\} \cap \Omega \subset \bigcup_{x \in \mathcal{T}_\lambda} \gamma_x$$

### Step 4 : Case where the original amodal completion has no contact

In this situation, it follows from our construction in Step 1 that (8) rewrites

$$\partial A_\lambda \cap \Omega = \bigcup_{x \in \mathcal{T}_\lambda} \gamma_x \cap \Omega,$$

thus, for almost every  $\lambda \in \mathbb{R}$ ,

$$\partial^* \{u_\gamma \geq \lambda\} \cap \Omega = \bigcup_{x \in \mathcal{T}_\lambda} \gamma_x \cap \Omega,$$

up to a  $\mathcal{H}^1$ -negligible set. It follows that  $u_\gamma \in \mathcal{S}_p(\Omega)$  and, as a direct consequence of the definition of the energies,

$$F_p(u_\gamma) = E_p(\gamma)$$

□



**Lemma 1** For any monotone family of sets  $(X_\lambda)_{\lambda \in \mathbb{R}} \subset \mathbb{R}^2$ , there exists a finite or countable set  $D$  such that

$$\lim_{\mu \rightarrow \lambda} X_\mu = X_\lambda \quad \forall \lambda \in \mathbb{R} \setminus D,$$

where convergence means convergence in measure. We shall call  $D$  the set of discontinuity points of  $(X_\lambda)_{\lambda \in \mathbb{R}}$ .

PROOF: It is enough to notice that the map  $\lambda \mapsto |X_\lambda|$  is monotone, thus has at most countably many discontinuity points, and to choose  $D$  as the set of these discontinuity points.  $\square$

**Lemma 2** Let  $u : \Omega \rightarrow [-\infty, +\infty]$  be a Borel function such that  $u \not\equiv +\infty$  and  $u \not\equiv -\infty$  up to a Lebesgue negligible set. If  $\Omega$  is bounded, connected and with Lipschitz boundary then  $\lambda \mapsto \mathcal{H}^1(\partial^*\{u \geq \lambda\} \cap \Omega)$  is in  $L^1(\mathbb{R})$  if and only if  $u \in \text{BV}(\Omega)$ .

PROOF: See [2, Lemma 1]  $\square$

**Lemma 3** Let  $u \in \mathcal{S}_p(\Omega)$ . There exists an amodal completion  $\gamma_u$  naturally associated with  $u$  such that, in view of Theorem 1,  $u = u_{\gamma_u}$  almost everywhere in  $\mathbb{R}^2$  and

$$E_p(\gamma_u) = F_p(u).$$

PROOF: Recall from the definition of  $\mathcal{S}_p(\Omega)$  that for almost every  $\lambda \in U_0(\partial\Omega)$ ,  $\partial^*\{u \geq \lambda\} \cap \overline{\Omega}$  is a finite union of simple curves  $\gamma_i^\lambda$ ,  $i = 0, \dots, n_\lambda$  with good properties. For every  $x \in \mathcal{T}$  such that  $\partial^*\{u \geq U_0(x)\} \cap \overline{\Omega}$  satisfy this decomposition, let us define  $\gamma_{u,x}$  as the unique curve  $\gamma_i^\lambda$  that passes through  $x$ . Clearly,  $\{\gamma_{u,x}, x \in \mathcal{T}\}$  satisfies all the properties of an amodal completion. In addition, the map  $\gamma_u : x \in \mathcal{T} \mapsto \gamma_{u,x}$  maps  $\mathcal{T}$  into  $W^{2,p}([0, 1], \mathbb{R}^2)$ . It follows from the definition of  $\mathcal{S}_p$  that  $\gamma_u$  is an amodal completion on  $\Omega$ . Observe now that, up to a  $\mathcal{H}^1$ -negligible set,  $\partial^*(\{u \geq \lambda\} \cap \overline{\Omega}) = \cup_{i=1}^{n_\lambda} \gamma_i^\lambda \cup (\{u \geq \lambda\} \cap \partial\Omega)$ . Let  $(\Gamma_j^\lambda)_{j=1}^{k_\lambda}$  denote the associated family of closed curves as given by Step 1 in the proof of Theorem 1 and let  $A_\lambda$  the associated set. Then  $\partial A_\lambda = \partial^*(\{u \geq \lambda\} \cap \overline{\Omega})$  up to a  $\mathcal{H}^1$ -negligible set and it follows from the Gauss-Green formula that  $A_\lambda = \{u \geq \lambda\} \cap \Omega$  up to a Lebesgue-negligible set. Since we already know that  $A_\lambda = \{u_{\gamma_u} \geq \lambda\} \cap \Omega$  up to a Lebesgue-negligible set, the uniqueness of the representation implies that  $u = u_{\gamma_u}$  almost everywhere in  $\mathbb{R}^2$ . The claim about the energy is a direct consequence of the definitions of  $\mathcal{S}_p(\Omega)$  and  $F_p$ .  $\square$

## 4 Minimizing the amodal energy of an amodal completion

We recall our assumption that the data to interpolate within  $\Omega$  is the trace of an analytic function  $U_0$  such that  $F_p(U_0) < +\infty$ . We assume that  $(U_0, \Lambda)$  is an admissible occlusion. Thus, the set of T-junctions  $\mathcal{T}$  is not empty and we can define a canonical amodal completion associated with  $U_0$  in the following way: for every  $x \in \mathcal{T}$  we let  $\gamma_x$  denote the connected component of  $\{U_0 = \lambda\} \cap \overline{\Omega}$  containing  $x$  and we remark that, by  $(H_1)$ ,  $\gamma_x$  is an analytic curve. It is then easily seen that the map  $\gamma_0 : x \in \mathcal{T} \mapsto \gamma_x$  is an amodal completion and, by the change of variables formula,  $E_p(\gamma_0) = F_p(U_0) < \infty$ . Then the following result can be established.

**Theorem 2** *The problem*

$$\min\{E_p(\gamma) : \gamma \in \mathcal{D}_p\} \tag{P_1}$$

*has at least one solution  $\gamma \in \mathcal{D}_p$  that can be associated with a function  $u_\gamma \in \text{BV}(\mathbb{R}^2)$ .*

PROOF: The canonical amodal completion associated with  $U_0$  has obviously finite energy. Thus we may consider a minimizing sequence of amodal completions  $(\gamma^\ell)_{\ell \in \mathbb{N}}$  and assume without loss of generality that

$$\sup_{\ell \in \mathbb{N}} E_p(\gamma^\ell) = C < \infty,$$

so that the functions

$$f_\ell(x) = E_p(\gamma_x^\ell)$$

are uniformly bounded in  $L^1(\partial\Omega, \mu)$ .

**Step 1. Convergence of the energies  $\mathbf{f}_\ell(\mathbf{x}) = \mathbf{E}_p(\gamma_x^\ell)$**

Since  $U_0$  is assumed to be nonconstant on  $\partial\Omega$ , we can without loss of generality renormalize the measure  $\mu$  so that  $\mu(\partial\Omega) = 1$ . Since  $\mu$  has no atoms, any point on  $\partial\Omega$  can be associated with a unique value in  $[0, 1[$ . More precisely, given an origin  $x_0$  on  $\partial\Omega \cap \mathcal{T}$ , we associate with any  $x \in \mathcal{T}$  the unique  $n \in [0, 1[$  such that  $n = \mu([x_0, x])$  and denote  $f_\ell(n) := f_\ell(x)$ . Conversely, almost every  $n \in [0, 1[$  is associated with a unique  $x \in \mathcal{T}$  such that  $n = \mu([x_0, x])$  and we shall write  $x = \mu^{-1}(n)$  for simplicity.

Let us now consider for  $k, N \in \mathbb{N}$  the dyadic intervals on  $[0, 1[$ :

$$I_{N,k} = [k2^{-N}, (k+1)2^{-N}[$$

and define the functions

$$f_\ell^N : m \in [0, 1] \mapsto 2^N \int_{I_{N,k}} f_\ell(n) dn = 2^N \int_{\mu^{-1}(I_{N,k})} f_\ell(x) d\mu(x)$$

where  $I_{N,k}$  is the unique dyadic interval containing  $m$ . Remark that the functions  $f_\ell^N$  are constant on each interval  $I_{N,k}$  and, for every  $m \in [0, 1[$ ,  $|f_\ell^N(m)| \leq 2^N \int_0^1 f_\ell(n) dn \leq 2^N C$ . Using a diagonal extraction argument, we can find a subsequence of  $(f_\ell)$ , still denoted by  $(f_\ell)$ , such that

$$\forall (N, k), \quad f_\ell^N(m) \rightarrow f^N(m) \quad \text{for every } m \in I_{N,k}$$

For every  $N \in \mathbb{N}$ , the limit function  $f^N$  is positive and piecewise constant on the  $I_{N,k}$ 's. Moreover, remark that  $I_{N,k} = [k2^{-N}, (2k+1)2^{-N-1}[ \cup [(2k+1)2^{-N-1}, (k+1)2^{-N}[$  and

$$\begin{aligned} \forall m \in [k2^{-N}, (2k+1)2^{-N-1}[ & \quad f_\ell^{N+1}(m) = 2^{N+1} \int_{I_{N+1,2k}} f_\ell(n) dn \\ \forall m \in [(2k+1)2^{-N-1}, (k+1)2^{-N}[ & \quad f_\ell^{N+1}(m) = 2^{N+1} \int_{I_{N+1,2k+1}} f_\ell(n) dn \end{aligned}$$

thus

$$\int_{I_{N,k}} f_\ell^{N+1}(n) dn = \int_{I_{N,k}} f_\ell(n) dn$$

and finally, for every  $m \in I_{N,k}$ ,

$$f_\ell^N(m) = 2^N \int_{I_{N,k}} f_\ell^{N+1}(n) dn$$

By the Dominated Convergence Theorem, it follows that

$$f^N(m) = 2^N \int_{I_{N,k}} f^{N+1}(n) dn,$$

which proves that  $(f^N)_{N \in \mathbb{N}}$  is a martingale. In addition, remark that

$$\int_0^1 f_\ell^N(n) dn = \sum_k \int_{I_{N,k}} f_\ell(n) dn = \int_0^1 f_\ell(n) dn \leq C,$$

so that, by Fatou's Lemma

$$\int_0^1 f^N(n) dn \leq C < \infty.$$

Since, in addition,  $f^N \geq 0$ , it follows that  $(f^N)_{N \in \mathbb{N}}$  is a bounded positive martingale. By Doob's convergence theorem, there exists  $f \in L^1([0, 1])$  such that

$$f^N \rightarrow f \quad \text{almost everywhere}$$

## Step 2. Definition of a limit amodal completion

Let  $N, k \in \mathbb{N}$  and recall from above that the sequence  $(f_\ell^N)_{\ell \in \mathbb{N}}$  satisfies

$$\sup_{n \in I_{N,k}} \sup_{\ell \in \mathbb{N}} f_\ell^N(n) < \infty.$$

**Lemma 4** *Let  $I := I_{N,k}$  and  $A_\ell := \{x \in \mu^{-1}(I) \cap \mathcal{T} : f_\ell(x) < 2^N \int_I f_\ell(n) dn + \frac{1}{N}\}$ . Then there exists some  $\epsilon > 0$  such that  $\mu(A_\ell) \geq \epsilon$  for every  $\ell \in \mathbb{N}$ .*

PROOF: By definition and since  $\mu(\partial\Omega \setminus \mathcal{T}) = 0$ ,  $\mu^{-1}(I) \setminus A_\ell$  essentially coincides with  $\{x \in \mu^{-1}(I) \cap \mathcal{T} : f_\ell(x) \geq 2^N \int_I f_\ell(n) dn + \frac{1}{N}\}$ , hence

$$\int_{\mu^{-1}(I) \setminus A_\ell} f_\ell(x) d\mu(x) \geq \mu(\mu^{-1}(I) \setminus A_\ell) \left( 2^N \int_I f_\ell(n) dn + \frac{1}{N} \right)$$

Assume that for all  $\epsilon > 0$  there exists  $\ell \in \mathbb{N}$  such that  $\mu(A_\ell) < \epsilon$ . Then,

$$\int_I f_\ell(n) dn \geq (2^{-N} - \epsilon) \left( 2^N \int_I f_\ell(n) dn + \frac{1}{N} \right),$$

and therefore

$$2^N \int_I f_\ell(n) dn \geq \frac{1}{N} \left( \frac{1}{\epsilon 2^N} - 1 \right),$$

which gives a contradiction for  $\epsilon$  small enough since  $\sup_{\ell \in \mathbb{N}} f_\ell^N < +\infty$  on  $I$ . □

**Lemma 5** *Under the notations above, there exists a T-junction  $x \in \mathcal{T} \cap I$  such that, possibly passing to a subsequence of  $(A_\ell)_{\ell \in \mathbb{N}}$ ,*

$$x \in A_\ell, \quad \forall \ell \in \mathbb{N}$$

PROOF:  $\{\bigcup_{k \geq n} A_k\}_{n \in \mathbb{N}}$  is a decreasing family of sets such that, by Lemma 4,  $\mu(\bigcup_{k \geq n} A_k) \geq \epsilon$ ,  $\forall n \in \mathbb{N}$ , and therefore

$$\mu\left(\bigcap_{n \in \mathbb{N}} \bigcup_{k \geq n} A_k\right) \geq \epsilon,$$

from which the lemma follows.  $\square$

We are now in position to finish the proof of Theorem 2. Denoting by  $x$  the T-junction given by the previous lemma and choosing the appropriate subsequence, we have

$$E_p(\gamma_x^\ell) \equiv f_\ell(x) \leq 2^N \int_{I_{N,k}} f_\ell(n) dn + \frac{1}{N} \equiv f_\ell^N(x) + \frac{1}{N} < \infty.$$

By the weak compactness of the unit ball in  $W^{2,p}$  there exists a further subsequence, and a limit arc  $\gamma_x^{N,k} \in W^{2,p}$  such that

- $\gamma_x^\ell \rightharpoonup \gamma_x^{N,k}$  weakly in  $W^{2,p}([0,1], \mathbb{R}^2)$  (thus strongly in  $C^1$ ).
- $E_p(\gamma_x^{N,k}) \leq \liminf_{\ell \rightarrow \infty} f_\ell(x)$ , using (6), the lower semicontinuity of the  $W^{2,p}$  norm and the fact that  $\mathcal{L}(\gamma_x^{N,k}) = \lim_{\ell \rightarrow \infty} \mathcal{L}(\gamma_x^\ell)$

Since  $\mathcal{H}^0(\partial\{U_0 \geq \lambda\})$  is finite, the limit arc  $\gamma_x^{N,k}$  passes through another T-junction  $y \in \mathcal{T}_{U_0(x)}$ . Thus  $\gamma_x^{N,k}$  can be extended outside  $\Omega$  using arcs of  $\{y \in \mathbb{R}^2 \setminus \Omega : U_0(y) = U_0(x)\}$ . Let us prove that the extended curve, defined for example on  $[-\epsilon, 1 + \epsilon]$ , is globally of class  $W^{2,p}$ . We already know that it is  $W^{2,p}$  on each interval  $[-\epsilon, 0]$ ,  $[0, 1]$  and  $[1, 1 + \epsilon]$ . In addition, each arc  $\gamma_x^\ell$  extends outside  $\Omega$  into a globally  $C^1$  arc. Since  $U_0$  is analytic outside  $\Omega$  and since the convergence of  $(\gamma_x^\ell)$  holds also in the strong topology of  $C^1$ , we infer that  $\gamma_x^{N,k}$  is in  $C^1([0,1], \mathbb{R}^2)$  and admits a globally  $C^1$  extension. By the usual properties of Sobolev functions, it follows that  $\gamma_x^{N,k}$  is of class  $W^{2,p}$  in  $(-\epsilon, 1 + \epsilon)$ .

Since the  $I_{N,k}$ 's are countably many, we can again use a diagonal extraction to get a subsequence, still denoted as  $f_\ell$ , and for each  $(N, k)$  a limit arc of class  $W^{2,p}([0,1])$ , denoted as  $\gamma^{N,k}$ , such that

$$E_p(\gamma^{N,k}) \leq 2^N \int_{I_{N,k}} f^N(n) dn + \frac{1}{N} \quad (= 2^N \lim_{\ell \rightarrow \infty} \int_{I_{N,k}} f_\ell(n) dn + \frac{1}{N})$$

Moreover, the limit curves  $(\gamma^{N,k})_{N,k}$  do not cross by construction and can be extended outside  $\Omega$  into globally  $W^{2,p}$  curves.

Let us now see how a limit curve can be defined for any T-junction. Given  $x \in \mathcal{T}$ , there exists for every  $N \in \mathbb{N}$  some  $k_N$  such that  $\mu([x_0, x]) \in I_{N, k_N}$ . Considering the family of arcs  $(\gamma^{N, k_N})_{N,k}$  defined above, we have by definition

$$E_p(\gamma^{N, k_N}) \leq 2^N \int_{I_{N, k_N}} f^N(n) dn + \frac{1}{N} = f^N(x) + \frac{1}{N}$$

which converges - for  $\mu$ -almost every  $x$  - to  $f(x)$ . Using the weak compactness of  $W^{2,p}$ , there exists a subsequence, still denoted by  $(\gamma^{N,k_N})_{N \in \mathbb{N}}$ , that weakly converges in  $W^{2,p}$ , thus uniformly in  $C^1$ , to a limit arc  $\gamma_x$ . By the lower semicontinuity of the  $W^{2,p}$  norm and the fact that  $\mathcal{L}(\gamma_x) = \lim_{N \rightarrow \infty} \mathcal{L}(\gamma^{N,k_N})$ , it follows that

$$E_p(\gamma_x) \leq \liminf_{N \rightarrow \infty} E_p(\gamma^{N,k_N}) \leq \liminf_{N \rightarrow \infty} (f^N(x) + \frac{1}{N}) = f(x).$$

This procedure can be applied for  $\mu$ -almost every T-junction and thus we can define a limit amodal completion  $\gamma$ . In particular, two different arcs  $\gamma_{x_1}$  and  $\gamma_{x_2}$  cannot cross (but may intersect tangentially) since they are uniform limits of arcs  $(\gamma^{N,k})_{N,k}$  that do not cross by construction.

There are two technical points that must be checked in this construction process :

1) Given  $x \in \mathcal{T}_\lambda$  and its associated limit curve  $\gamma_x \in W^{2,p}(0,1)$ , we have  $\gamma_x(1) \in \mathcal{T}_\lambda$ . Thus  $\gamma_x$  can be extended outside  $\Omega$  using arcs of  $\{y \in \mathbb{R}^2 \setminus \Omega : U_0(y) = U_0(x)\}$  into a curve  $\gamma_x^\epsilon$  defined on  $[-\epsilon, 1 + \epsilon]$ . Following the same argument as above, we prove that  $\gamma_x^\epsilon$  is of class  $W^{2,p}$  on  $(-\epsilon, 1 + \epsilon)$ ;

2) We must control whether the curve  $\gamma_x$  passing through another T-junction  $y \in \mathcal{T}_{U_0(x)}$  coincides with  $\gamma_y$ . The answer is positive because there are finitely many T-junctions per level and because the convergence of the curves is meant in the strong topology of  $C^1$ .

Finally, we have built a limit amodal completion  $\gamma$  defined for  $\mu$ -almost every  $x \in \mathcal{T}$  and such that each curve  $\gamma_x$  is of class  $W^{2,p}$  and can be extended outside  $\Omega$  into a globally  $W^{2,p}$  curve whose restriction to  $\mathbb{R}^2 \setminus \Omega$  coincides with arcs of  $\{y \in \mathbb{R}^2 \setminus \Omega : U_0(y) = U_0(x)\}$ . Moreover,

$$E_p(\gamma(x)) \leq f(x), \quad \mu - \text{a.e. } x \in \mathcal{T}$$

It follows from Fatou's Lemma that

$$\begin{aligned} E_p(\gamma) &= \int_{\partial\Omega} E_p(\gamma(x)) d\mu(x) \leq \int_{\partial\Omega} f(x) d\mu(x) = \int_0^1 f(n) dn \\ &\leq \liminf_{N \rightarrow \infty} \int_0^1 f^N(n) dn \leq \liminf_{N \rightarrow \infty} \liminf_{\ell \rightarrow \infty} \int_0^1 f_\ell^N(n) dn = \liminf_{\ell \rightarrow \infty} \int_0^1 f_\ell(n) dn \end{aligned}$$

Thus

$$E_p(\gamma) \leq \liminf_{\ell \rightarrow \infty} E_p(\gamma^\ell) = \inf_{\tilde{\gamma} \in \mathcal{D}_p} E_p(\tilde{\gamma})$$

and we have proved that the limit amodal completion is optimal.  $\square$

**Remark 2** The previous theorem involves a definition of convergence in the class of amodal completions, namely,  $(\gamma_h)_{h \in \mathbb{N}} \rightarrow \gamma$  if

1. for each dyadic interval  $[k_N 2^{-N}, (k_N + 1) 2^{-N})$ , there exists an appropriate point  $x_{k_N, N} \in [k_N 2^{-N}, (k_N + 1) 2^{-N})$  such that  $\gamma_h(x_{k_N, N})$  converges weakly in  $W^{2,p}$  to  $\gamma(x_{k_N, N})$ ;
2. for  $\mu$ -almost every  $x \in \partial\Omega$ ,  $\gamma_x$  is the weak limit in  $W^{2,p}$  of a subsequence  $(\gamma(x_{k_M, M}))_{M \in \mathbb{N}}$  where  $x_{k_M, M} \rightarrow x$  as  $M \rightarrow \infty$ .

**Corollary 1** *Let  $(\gamma_h)_{h \in \mathbb{N}}$  be a sequence of amodal completions with uniformly bounded energies, i.e.*

$$\sup_{h \in \mathbb{N}} E_p(\gamma_h) < \infty.$$

*Then, possibly extracting a subsequence, there exists a limit amodal completion  $\gamma$  such that  $(\gamma_h)_{h \in \mathbb{N}}$  converges to  $\gamma$  in the sense of Remark 2 above.*

It is easily seen that the function  $u_\gamma$  associated with an amodal completion  $\gamma$ , as defined by Theorem 1, may have level lines that are not curves of  $\gamma$  (see figure 4). Consequently, the relationship between  $\overline{F_p}(u_\gamma)$  and  $E_p(\gamma)$  is not clear. The purpose of Lemma 7 is to provide a continuous function  $u_\epsilon \in \mathcal{S}_p$  whose level lines are very similar to the curves of  $\gamma$  and such that  $F_p(u_\epsilon)$  be arbitrarily close to  $E_p(\gamma)$ . We start with a lemma providing a way to separate the curves of an amodal completion.

**Lemma 6** *Let  $\gamma$  be an amodal completion with finite energy  $E_p(\gamma)$ . Then for every  $\eta > 0$  there is another amodal completion  $\gamma^\eta$  without contact such that  $|E_p(\gamma^\eta) - E_p(\gamma)| \leq \eta$  and for  $\mu$ -almost every  $x \in \partial\Omega$ ,*

$$\sup_{s \in [0,1]} |\gamma_x^\eta(s) - \gamma_x(s)| \leq \eta.$$

PROOF: The proof is tedious, but not deep. We consider a dense set  $x_n$  of T-junctions in  $\mathcal{T}$  such that the curves  $\gamma_n = \gamma_{x_n}$  have all finite energy  $E_p(\gamma_n) < \infty$ . The idea of the proof is to move all curves of the amodal completion smoothly and slightly in such a way that they all fall apart from  $\gamma_n$ . In other terms, we shall dilate slightly all curves  $\gamma_n$ . After its dilation,  $\gamma_n$  will have an open security region around it into which no other curve  $\gamma_x$  enters. Given two curves  $\gamma_x$  and  $\gamma_y$  with  $x \neq y$ , we can find a curve  $\gamma_n$  which separates them in wide sense. After the dilation,  $\gamma_x$  and  $\gamma_y$  will not touch anymore. In that way, any two distinct curves  $\gamma_x$  and  $\gamma_y$  will be separated by an open domain and have therefore a positive distance to each other. This argument needs some detail. Indeed, notice that a curve  $\gamma_n$  may meet the boundary and that it may meet itself. Thus, we must be careful to move the curve away from the boundary and to move it apart from itself at points where it is tangent to itself. The dilations of  $\gamma_n$  will be done by smooth diffeomorphisms close enough to identity, which will increase very little the energy of the curves.

### Step 1. Dividing all curves $\gamma_n$ into graphs

We start by covering the domain  $(0, L_n)$  of  $\gamma_n$  with a finite set of open intervals  $(s_i, t_i)$  such that

1.  $s_i < s_{i+1} < t_i < t_{i+1}$ ,  $s_0 = 0$ ,  $t_N = L_n$ ,
2.  $\gamma_n$  restricted to  $[s_i, t_i]$  is a graph
3. the restriction of  $\gamma_n$  to  $[s_i, t_i]$  meets  $\partial\Omega$  at most on one side.

We index by  $i \in \mathbb{N}$  all the pieces of curves of all  $\gamma_n$ .

### Step 2. Defining a diffeomorphism dilating locally $\gamma_n$ .

We can translate the second and third conditions in the following way. On the interval  $[s_i, t_i]$ , the curve  $\gamma_n$  is represented as a graph  $\Gamma_i = \{(x, f_i(x))\}$ ,  $x \in [0, x_i]$  in local euclidian coordinates  $(x, y)$ . The third condition implies that if  $\Gamma_i$  touches  $\partial\Omega$  at several points, then it is only from above or only from below. Assume that it is from above, the other case being similar (the case where  $\Gamma_i$

does not touch  $\partial\Omega$  can be treated using indifferently one or the other way). There exists a  $C^\infty$  function  $y = \psi_i(x)$  such that  $\psi_i(x) > f_i(x)$  on  $(0, x_i)$ ,  $\psi_i(0) = f_i(0)$ ,  $\psi_i(x_i) = f_i(x_i)$  and the open domain  $\mathcal{D}_i = \{(x, y), 0 < x < x_i, f_i(x) < y < \psi_i(x)\}$  is contained in  $\Omega$  (see Figure 8).




Figure 8: The operator  $F_i^\epsilon$  differs from the identity in  $D_i$ . It is designed to move the curves  $\gamma_m$  and  $\gamma_p$  apart from  $\gamma_n$  and to move  $\gamma_n$  itself apart from  $\partial\Omega$ .

We consider the diffeomorphism of  $\mathbb{R}^2$  defined by

$$F_i^\epsilon(x, y) = \begin{cases} (x, y + \epsilon M_i e^{-\frac{1}{(\psi_i(x)-y)^+}} \cdot e^{-\frac{1}{x^+} - \frac{1}{(x_i-x)^+}}) & \text{if } 0 \leq x \leq x_i, f_i(x) \leq y \leq \psi_i(x), \\ (x, y) & \text{otherwise} \end{cases}$$

with

$$M_i \leq \inf_{0 \leq x \leq x_i} \frac{1}{1 + \psi_i'^2(x) + |\psi_i''(x)|}.$$

Then we see that  $F_i^\epsilon$  is  $C^\infty$  and there exists a constant  $C > 0$  independent of  $i$  and independent of the curve  $\gamma_n$  such that :

- $DF_i = Id + \epsilon G_i$ , where  $G_i$  is  $C^\infty$  and uniformly bounded on  $[0, x_i]$  by  $C$ ;
- $D^2F_i^\epsilon = \epsilon H_i$ , where  $H_i$  is  $C^\infty$  and uniformly bounded on  $[0, x_i]$  by  $C$ ;

This follows immediately from the chain rule and the fact that  $s \rightarrow e^{-\frac{1}{s^+}}$  is a  $C^\infty$  function with all derivatives bounded on  $\mathbb{R}$ .

### Step 3. Using the diffeomorphism to move all curves apart from $\gamma_n$

We define the following operation, indexed by  $i \in \mathbb{N}$ . For every curve  $\gamma_x$  of the amodal completion, we consider all maximal intervals  $(s, t)$  of  $\gamma_x$  contained in the domain  $\overline{\mathcal{D}_i}$  and replace  $\gamma_x$  on  $(s, t)$  by the new curve  $F_i^\epsilon \circ \gamma_x$ . We also replace  $\gamma_n$  on  $(s_i, t_i)$  by  $F_i^{\frac{\epsilon}{2}} \circ \gamma_n$ . Now, the curve  $\gamma_n$  may meet itself and we have to fix a rule for moving the parts of  $\gamma_n$  which coincide with the restriction of  $\gamma_n$  to  $(s_i, t_i)$ . Let  $(s, t)$  be a maximal interval of  $\gamma_n$  contained in  $\overline{\mathcal{D}_i}$ . If  $\gamma_n((s, t))$  meets  $\mathcal{D}_i$ , we consider that this part of  $\gamma_n$  is ‘‘above’’ the restriction to  $(s_i, t_i)$  and we move this part of the curve as we moved the other  $\gamma_x$  (see Figure 9). If instead  $\gamma_n([s, t])$  coincides with  $\gamma_n([s_i, t_i])$ , we consider the maximal intervals  $[\sigma_i, \tau_i]$  and  $[\sigma, \tau]$  on which both curves coincide. Assume for instance

that  $\gamma_n(\sigma_i) = \gamma_n(\tau)$  (these pieces of curves can also have same orientation, i.e.  $\gamma_n(\sigma_i) = \gamma_n(\sigma)$ ). Consider a definition of the unit normal  $n(s)$  to  $\gamma_n$  such that on  $(s, t)$ ,  $n(s)$  has an acute angle with the coordinate axis  $(0, y)$ . The curve  $\gamma_n$  is a graph around  $\sigma_i$  in euclidian coordinates such that  $n(\sigma_i)$  gives the  $(0, y)$  axis (see Figure 10). If in these coordinates, the graph of  $\gamma_n$  around  $\sigma_i$  is above the graph of  $\gamma_n$  around  $\tau$ , we do not move the restriction of  $\gamma_n$  to  $(s, t)$ . If instead it is below, we move the restriction of  $\gamma_n$  to  $(s, t)$  as the other curves  $\gamma_x$ . An analogous procedure applies when  $\gamma_n(\sigma_i) = \gamma_n(\sigma)$ .

This specification had to be made in order to avoid creating a self-crossing of  $\gamma_n$ .




Figure 9:  $\gamma_n$  has autocontact on a maximal proper subset of  $\gamma_n([s_i, t_i])$ .




Figure 10: There exists an autocontact set that strictly contains  $\gamma_n([s_i, t_i])$ .

#### Step 4. Checking that the moving apart does not increase much the energy of the amodal completion

By construction, there exists  $C > 0$  such that  $\|DF_i^\varepsilon - Id\| \leq C\varepsilon$  and  $\|D^2F_i^\varepsilon\| \leq C\varepsilon$ . It is easily checked that the energy of a curve  $\gamma$  deformed by  $F_i^\varepsilon$  satisfies

$$(1 - D\varepsilon)E_p(\gamma_x) \leq E_p(F_i^\varepsilon(\gamma_x)) \leq (1 + D\varepsilon)E_p(\gamma_x)$$

for some constant  $D > C$  independent of  $\gamma_x$ . Thus, taking  $\eta$  such that  $D\varepsilon = \eta 2^{-i}$  and setting  $F_i = F_i^\varepsilon$ , we can ensure that the energy of the whole amodal completion, denoted by  $F_i(\gamma)$ , satisfies

$$|E_p(F_i(\gamma)) - E_p(\gamma)| \leq \eta 2^{-i}. \quad (12)$$



This also entails that for any pair of points  $z_1$  and  $z_2$  belonging to some curves  $\gamma_{x_1}$  and  $\gamma_{x_2}$ ,

$$(1 + \eta 2^{-i})|z_1 - z_2| \geq |F_i(z_1) - F_i(z_2)| \geq (1 - \eta 2^{-i})|z_1 - z_2|. \quad (13)$$

One moving apart operation therefore defines a new amodal completion with energy as close as we wish to the original and curves as close as we wish to the originals. The moving apart operation has then to be performed recursively at step  $i$  on the amodal completions resulting from the  $i - 1$  former operations. To formalize this, we set  $T_i = F_{i-1} \circ F_i \circ \dots \circ F_2 \circ F_1$  and  $T(z) = \lim_{i \rightarrow \infty} T_i(z)$ . So at the  $i$ -th step, all operations described in steps 1 to 5 are applied to the curves  $T_{i-1}(\gamma_x)$  ( $T_0 = Id$ ). From (13) follows that  $T$  is a bilipschitz map for  $\eta < \frac{1}{2}$ , so that open sets are mapped onto open sets.

**Step 5. The last remark shows that the moving apart operation isolates  $\gamma_n$  from all other curves and eliminates its self-contacts on  $(s_i, t_i)$ .**

Indeed, the fact that we move  $\gamma_n$  only halfway at step  $i$  implies that  $T_i(\gamma_n((s_i, t_i)))$  is contained in the open set  $O_i = \mathcal{D}_i \setminus F_i(D_i)$ , which contains no piece of no other curve of the amodal completion. Now, by the same argument as for  $T_i$ ,  $\tilde{T}_i = \lim_{k \rightarrow \infty} F_k \circ \dots \circ F_{i+1}$  also is a bilipschitz map. So the final position of  $\gamma_n(s_i, t_i)$  is the open domain  $\tilde{T}_i(O_i)$ . This being true for all  $i$ , we deduce that every curve  $T(\gamma_n)$  is contained (except its endpoints) in an open set  $\mathcal{D}_n$  which does not contain any other curve  $T(\gamma_x)$ .

**Step 6. Iteration of the moving apart operation.**

This iteration is defined for each curve  $\gamma_x$  by  $T_i(\gamma_x) = F_i \circ F_{i-1} \circ \dots \circ F_2 \circ F_1(\gamma_x)$ . By (13), the sequence of curves  $T_i(\gamma_x)$  converges uniformly to a curve  $T(\gamma_x)$  and by (12) and Fatou's lemma,  $|E_p(T(\gamma)) - E_p(\gamma)| \leq \eta$ .

**Step 7. Conclusion.**

Given two curves  $\gamma_x$  and  $\gamma_y$  of the amodal completion, there exists a curve  $\gamma_n$  which separates  $\gamma_x$  and  $\gamma_y$ , namely  $\gamma_x$  and  $\gamma_y$  do not belong to the same connected component of  $\Omega \setminus \gamma_n$ . Thus  $T(\gamma_x)$  and  $T(\gamma_y)$  are contained in two different connected components of  $\Omega \setminus \mathcal{D}_n$  and therefore stand at a positive distance from each other.

Let us now deal with curves  $\gamma_x$  which have at least one self-meeting. Call loops of  $\gamma_x$  the open connected components of  $\Omega \setminus \gamma_x$  whose boundary is fully contained in  $\gamma_x$ . If at least one loop of  $\gamma_x$  does not contain any piece of any other curve  $\gamma_n$  – which means that the previous procedure will let the loop unchanged – this is equivalent to say that it does not contain any piece of any other curve  $\gamma_x$ . Since loops have positive measure, we deduce that only a countable set of curves  $\gamma_x$  can have such empty loops. So we are allowed to add them up from the start to the curves  $\gamma_n$ . Thus, we may assume that all curves  $\gamma_x$  having a loop are such that the loop contains some piece of  $\gamma_n$ . This implies that the self-meeting points of  $\gamma_x$  also are self-meeting points for some  $\gamma_n$  and we conclude that the moving apart operation have moved them apart too.  $\square$

**Lemma 7** *Let  $\gamma \in \mathcal{D}_p$  be an amodal completion with finite energy and  $u_\gamma$  the associated function in BV. For every  $\epsilon > 0$  there exists a continuous function  $u_\epsilon \in \mathcal{S}_p$  such that  $|E_p(\gamma) - F_p(u_\epsilon)| \leq \epsilon$ . In addition,  $u_\epsilon$  tends to  $u_\gamma$  in  $L^1$  as  $\epsilon$  tends to zero.*

PROOF: In view of Lemma 6, for every  $\epsilon > 0$  there exists an amodal completion without contact  $\gamma_\epsilon$  such that  $|E_p(\gamma_\epsilon) - E_p(\gamma)| < \epsilon$ . By Theorem 1,  $\gamma_\epsilon$  can be associated with a function  $u_\epsilon \in \mathcal{S}_p$  and  $|E_p(\gamma) - F_p(u_\epsilon)| \leq \epsilon$ . In addition,  $u_\epsilon$  is continuous because its level lines are disjoint by construction.

From the construction procedure of Theorem 1 and the fact that the curves of  $\gamma_\epsilon$  converge uniformly to the curves of  $\gamma$ , we also deduce that  $u_\epsilon$  tends to  $u$  almost everywhere on  $\Omega$ . Remark now that, by construction, for every  $\epsilon > 0$  and for almost every  $x \in \Omega$ ,  $|u_\epsilon(x)| \leq \max_{y \in \partial\Omega} |U_0(y)|$ . We infer by the Dominated Convergence Theorem that  $u_\epsilon$  tends to  $u_\gamma$  in  $L^1$  as  $\epsilon$  tends to zero.  $\square$

**Lemma 8** *Let  $u \in C^2$  such that  $\mathcal{F}_p(u) < \infty$  and  $u$  coincide with  $U_0$  outside  $\Omega$ . Then there exists an amodal completion  $\gamma$  whose trace is contained in the topographic map of  $u$ , i.e. for almost every  $\lambda \in \Lambda$ ,  $\bigcup_{x \in \mathcal{T}_\lambda} (\gamma_x) \subset \{u = \lambda\} \cap \overline{\Omega}$ . Consequently,*

$$E_p(\gamma) \leq \mathcal{F}_p(u)$$

PROOF: This amodal completion will be constructed as a selection of level lines of  $u$  inside  $\Omega$ . By Sard Lemma we can find a set  $\tilde{\Lambda} \subset \Lambda$  of full measure such that  $\{u = \lambda\} \cap \Omega$  is a union of  $C^2$  curves. Of course we also have  $\mu(\{x \in \partial\Omega, u(x) \notin \tilde{\Lambda}\}) = 0$ . For every  $T$ -junction  $x \in \mathcal{T}$  such that  $u(x) \in \tilde{\Lambda}$ , the level line  $L_x$  of  $u$  passing by  $x$  is by definition of  $\tilde{\Lambda}$  a  $C^2$  Jordan curve and intersects  $\partial\Omega$  at some other  $T$ -junction  $y \in \mathcal{T}_{u(x)}$ . We take  $\gamma_x$  to be a  $C^2$  parameterization on  $[0, 1]$  of the arc of  $L_x$  between  $x$  and  $y$ . The map  $\gamma : x \in \mathcal{T} \cap u^{-1}(\tilde{\Lambda}) \mapsto \gamma_x$  is clearly an amodal completion whose trace is contained in the topographic map of  $u$ . The inequality  $E_p(\gamma) \leq \mathcal{F}_p(u)$  is then an obvious consequence of the coarea formula, as  $E_p(\gamma)$  is obtained from  $\mathcal{F}_p$  by a restriction to the levels of  $\tilde{\Lambda}$  and a selection of pieces of level lines at these levels.  $\square$

## 5 Comparison with the direct variational approach

This section is devoted to the proof that the problems  $(P_1)$  and  $(P_2)$  are equivalent. We do not know whether they also are equivalent with  $(P'_2)$ ; this would actually be true if one could prove that for  $u \in \mathcal{S}_p$ ,  $F_p(u) = \overline{\mathcal{F}_p}(u)$ . However two points remain unclear:

- we do not know whether any function  $u \in \mathcal{S}_p$  can be approximated by a sequence of smooth functions  $(u_h)$  such that  $F_p(u_h)$  tends to  $F_p(u)$  (this sounds reasonable);
- we do not know whether any  $u \in \mathcal{S}_p$  can be approximated by a sequence of smooth functions  $(u_h)$  such that  $\overline{\mathcal{F}_p}(u) = \lim_{h \rightarrow \infty} \mathcal{F}_p(u_h)$  and each  $u_h$  can be associated with a function  $\tilde{u}_h \in \mathcal{S}_p$  such that  $F_p(\tilde{u}_h) \leq \mathcal{F}_p(u_h)$  and  $(\tilde{u}_h)$  tends to  $u$  in  $L^1$ .

We start with the proof that  $(P_2)$  is well posed.

**Theorem 3** *The problem*

$$\min\{\overline{F}_p(u) : u = U_0 \text{ on } \mathbb{R}^2 \setminus \Omega\} \tag{P_2}$$

*has at least one solution  $u \in \text{BV}(\mathbb{R}^2)$  that coincides with  $U_0$  outside  $\Omega$ . In addition, for almost every  $\lambda \in u(\mathbb{R}^2)$ , there exists a finite family  $\Gamma^\lambda = \{\gamma_i^\lambda\}_{i \in I_\lambda}$  of regular curves of class  $W^{2,p}$  such that  $\partial^*\{u > \lambda\} \subset \bigcup_{i \in I_\lambda} (\gamma_i^\lambda)$  and any two curves of  $\Gamma^\lambda$  may intersect but only tangentially and without crossing each other.*

PROOF: Let  $(u_h)_{h \in \mathbb{N}}$  be a minimizing sequence. It follows from the definition of  $\overline{F}_p$  that  $(u_h)$  has uniformly bounded variation. Since every  $u_h$  coincides with  $U_0$  outside  $\Omega$ , we deduce from Lemma 2 that  $(u_h)$  is also uniformly bounded in  $L^1$ . Thus there exists a subsequence, still denoted

by  $(u_h)_{h \in \mathbb{N}}$ , and a limit function in BV such that  $u_h$  converges to  $u$  in  $L^1$ . It follows from the lower semicontinuity of relaxed functionals [17] that

$$\overline{F_p}(u) \leq \liminf_{h \rightarrow \infty} \overline{F_p}(u_h) = \inf\{\overline{F_p}(u) : u = U_0 \text{ on } \mathbb{R}^2 \setminus \Omega\}.$$

Thus  $u$  is a solution of  $(P_2)$ . In addition, by the usual extension theorems in BV[3],  $u \in \text{BV}(\mathbb{R}^2)$ .

By the properties of relaxed functionals, there exists a sequence  $(v_h)_{h \in \mathbb{N}}$  of functions in  $\mathcal{S}_p$  such that  $\overline{F_p}(u) = \lim_{h \rightarrow \infty} F_p(v_h)$ . By definition, for every  $h \in \mathbb{N}$  and for almost every  $t \in U_0(\partial\Omega)$ ,  $\partial^*\{v_h \geq t\} \cap \Omega$  essentially coincides with a finite union of curves of class  $W^{2,p}$ . In addition,

$$F_p(v_h) = \int_{-\infty}^{+\infty} \int_{\Omega \cap \partial^*\{v_h \geq t\}} (1 + |\kappa|^p) d\mathcal{H}^1.$$

By Fatou's lemma we get that

$$\int_{-\infty}^{+\infty} \liminf_{h \rightarrow \infty} \int_{\Omega \cap \partial^*\{v_h \geq t\}} (1 + |\kappa|^p) d\mathcal{H}^1 \leq \liminf_{h \rightarrow \infty} \int_{-\infty}^{+\infty} \int_{\Omega \cap \partial^*\{v_h \geq t\}} (1 + |\kappa|^p) d\mathcal{H}^1 < +\infty.$$

Thus for almost every  $t \in \mathbb{R}$ , there exists a subsequence (depending on  $t$ ), still denoted as  $(v_h)$ , such that  $\sup_{h \in \mathbb{N}} \int_{\Omega \cap \partial^*\{v_h \geq t\}} (1 + |\kappa|^p) d\mathcal{H}^1 < \infty$ . Since  $v_h \rightarrow u$  in  $L^1$ , the Cavalieri's principle implies that  $\{v_h \geq t\}$  converges for almost every  $t$  to  $\{u \geq t\}$  in  $L^1$ . The final claim of the Theorem is then a simple application of Theorem 7.1 in [6] (see also Corollary 1 in [4]).  $\square$

**Remark 3** An interesting consequence of the next result is that it provides an explicit integral formulation for  $\overline{F_p}(u)$  when  $u$  is a minimizer of  $(P_2)$ . Such a formulation could also be obtained, under a slightly different form, by combining the direct method developed in [4] with Theorem 8.6 in [7]. Indeed, by passing to a subsequence for which there is convergence in  $L^1$  of the characteristic functions of almost every level set, it can be easily proved that almost every limit set  $A_t$  has finitely many singularity points. Then Theorem 8.6 in [7] provides an explicit formula for the relaxed energy of  $A_t$ , from which an expression of  $\overline{F_p}$  is easily deduced.

**Theorem 4** *Problems  $(P_1)$  and  $(P_2)$  are equivalent, i.e.*

$$\min\{E_p(\gamma) : \gamma \in \mathcal{D}_p\} = \min\{\overline{F_p}(u) : u = U_0 \text{ on } \mathbb{R}^2 \setminus \Omega\}$$

*In addition, if  $\gamma \in \mathcal{D}_p$  is a minimizer of  $(P_1)$  then  $u_\gamma$  is a minimizer of  $(P_2)$  and  $\overline{F_p}(u_\gamma) = E_p(\gamma)$ . Conversely, if  $u$  is a minimizer of  $(P_2)$  then there exists an amodal completion  $\gamma_u$  such that  $\gamma_u$  is a minimizer of  $(P_1)$  and its associated function  $u_\gamma$  coincides with  $u$  almost everywhere. In addition,  $\overline{F_p}(u) = E_p(\gamma_u)$ .*

**PROOF:** Let  $u$  be a minimizer of  $(P_2)$  and consider a sequence  $(u_h)_{h \in \mathbb{N}}$  of functions in  $\mathcal{S}_p$  such that  $u_h$  tends to  $u$  in  $L^1$  and  $F_p(u_h)$  converges to  $\overline{F_p}(u)$ . According to Lemma 3, we can associate with each  $u_h$  an amodal completion  $\gamma_h$  such that  $F_p(u_h) = E_p(\gamma_h)$ . Since  $\sup_{h \in \mathbb{N}} E_p(\gamma_h) < \infty$ , there exists by Corollary 1 a limit amodal completion  $\gamma$  such that  $(\gamma_h)$  converges to  $\gamma$  (in the sense of Corollary 1) and  $E_p(\gamma) \leq \liminf_{h \rightarrow \infty} E_p(\gamma_h) = \overline{F_p}(u)$ . It follows that

$$\min\{E_p(\gamma) : \gamma \in \mathcal{D}_p\} \leq \min\{\overline{F_p}(u) : u = U_0 \text{ on } \mathbb{R}^2 \setminus \Omega\}.$$

Conversely, let  $\gamma$  be a minimizer of  $(P_1)$ . By Lemma 7, for each  $\epsilon > 0$  there exists a continuous  $u_\epsilon \in \mathcal{S}_p$  such that  $|E_p(\gamma) - F_p(u_\epsilon)| \leq \epsilon$  and, in addition,  $u_\epsilon$  tends to  $u_\gamma$  in  $L^1$  where  $u_\gamma$  is associated with  $\gamma$  through Theorem 1. It follows from the lower semicontinuity of  $\overline{F}_p$  that  $\overline{F}_p(u_\gamma) \leq E_p(\gamma)$  which proves that

$$\min\{E_p(\gamma) : \gamma \in \mathcal{D}_p\} \geq \min\{\overline{F}_p(u) : u = U_0 \text{ on } \mathbb{R}^2 \setminus \Omega\}$$

thus

$$\min\{E_p(\gamma) : \gamma \in \mathcal{D}_p\} = \min\{\overline{F}_p(u) : u = U_0 \text{ on } \mathbb{R}^2 \setminus \Omega\}.$$

In particular,  $\overline{F}_p(u_\gamma) = \min\{\overline{F}_p(u) : u = U_0 \text{ on } \mathbb{R}^2 \setminus \Omega\} = E_p(\gamma)$  and therefore  $u_\gamma$  is a minimizer of  $(P_2)$ .

Take now again a minimizer  $u$  of  $(P_2)$ . Like above we may find a sequence  $(u_h)$  of functions in  $\mathcal{S}_p$  and their associated amodal completions  $(\gamma_h)$  such that  $u_h \rightarrow u$  in  $L^1$ ,  $\overline{F}_p(u) = \lim_{h \rightarrow \infty} F_p(u_h)$  and  $(\gamma_h)$  converges (in the sense of Corollary 1) to a limit amodal completion  $\gamma$  such that  $E_p(\gamma) \leq \liminf_{h \rightarrow \infty} E_p(\gamma_h)$ . Let  $u_\gamma$  denote the function associated with  $\gamma$  according to Theorem 1. It follows from the construction procedure of Theorem 1 and the fact that all functions coincide with  $U_0$  on  $\partial\Omega$  that for almost every  $t \in \mathbb{R}$ ,  $\{u_h \geq t\}$  tends to  $\{u_\gamma \geq t\}$  in  $L^1$ . Besides, since  $(u_h)$  tends to  $u$  in  $L^1$ , by Cavalieri's principle and possibly taking a subsequence, we also know that for almost every  $t \in \mathbb{R}$ ,  $\{u_h \geq t\}$  tends to  $\{u \geq t\}$  in  $L^1$ . By the argument of Step 2 in the proof of Theorem 1,  $u$  and  $u_\gamma$  coincide almost everywhere on  $\mathbb{R}^2$ , and the theorem ensues. Remark in particular that, according to Theorem 1, for almost every  $t \in \Lambda$ ,

$$\partial^*\{u \geq t\} \cap \overline{\Omega} = \partial^*\{u_\gamma \geq t\} \cap \overline{\Omega} \subset \bigcup_{x \in \mathcal{I}_t} \gamma_u(x) \quad \text{up to a } \mathcal{H}^1\text{-negligible set.}$$

and

$$\overline{F}_p(u) = \overline{F}_p(u_\gamma) = E_p(\gamma_u).$$

□

## References

- [1] L. Alvarez, Y. Gousseau, and J.-M. Morel. The size of objects in natural and artificial images. *Advances in Imaging and Electron Physics*, 111:167–242, 1999.
- [2] L. Ambrosio, V. Caselles, S. Masnou, and J.-M. Morel. Connected components of sets of finite perimeter and applications to image processing. *J. Eur. Math. Soc.*, 3:39–92, 2001.
- [3] L. Ambrosio, N. Fusco, and D. Pallara. *Functions of Bounded Variation and Free Discontinuity Problems*. Oxford University Press, 2000.
- [4] L. Ambrosio and S. Masnou. A direct variational approach to a problem arising in image reconstruction. *Interfaces and Free Boundaries*, 5:63–81, 2003.
- [5] C. Ballester, M. Bertalmio, V. Caselles, G. Sapiro, and J. Verdera. Filling-in by joint interpolation of vector fields and gray levels. *IEEE Trans. On Image Processing*, 10(8):1200–1211, 2001.

- [6] G. Bellettini, G. Dal Maso, and M. Paolini. Semicontinuity and relaxation properties of a curvature depending functional in 2D. *Ann. Scuola Norm. Sup. Pisa Cl. Sci*, (4) 20, no 2:247–297, 1993.
- [7] G. Bellettini and L. Mugnai. Characterization and representation of the lower semicontinuous envelope of the elastica functional. *Ann. Inst. H. Poincaré, Anal. Non Lin.*, 21(6):839–880, 2004.
- [8] G. Bellettini and L. Mugnai. A varifolds representation of the relaxed elastica functional. submitted.
- [9] M. Bertalmio, A. Bertozzi, and G. Sapiro. Navier-Stokes, fluid dynamics, and image and video inpainting. In *Proc. IEEE Int. Conf. on Comp. Vision and Pattern Recog.*, Hawaiï, 2001.
- [10] M. Bertalmio, G. Sapiro, V. Caselles, and C. Ballester. Image inpainting. In *Proc. ACM Conf. Comp. Graphics (SIGGRAPH), New Orleans, USA*, pages 417–424, 2000.
- [11] M. Bertalmio, L. Vese, G. Sapiro, and S. Osher. Simultaneous structure and texture image inpainting. *IEEE Transactions on Image Processing*, 12(8):882–889, 2003.
- [12] R. Bornard, E. Lecan, L. Laborelli, and J.-H. Chenot. Missing data correction in still images and image sequences. In *Proc. 10th ACM Int. Conf. on Multimedia*, pages 355–361, 2002.
- [13] T.F. Chan, S.H. Kang, and J. Shen. Euler’s elastica and curvature based inpainting. *SIAM Journal of Applied Math.*, 63(2):564–592, 2002.
- [14] T.F. Chan and J. Shen. Mathematical models for local deterministic inpaintings. *SIAM Journal of Applied Math.*, 62(3):1019–1043, 2001.
- [15] T.F. Chan and J. Shen. Non-texture inpainting by curvature-driven diffusion (CDD). *Journal of Visual Comm. and Image Rep.*, 12(4):436–449, 2001.
- [16] A. Criminisi, P. Pérez, and K. Toyama. Object removal by exemplar-based inpainting. In *IEEE Int. Conf. Comp. Vision and Pattern Recog.*, volume 2, pages 721–728, 2003.
- [17] G. Dal Maso. *An Introduction to  $\Gamma$ -convergence*. Birkhäuser, 1993.
- [18] A. Efros and T. Leung. Texture synthesis by non-parametric sampling. In *Proc. Int. Conf. on Comp. Vision*, volume 2, pages 1033–1038, Kerkyra, Greece, 1999.
- [19] M. Elad, J.-L. Starck, D. Donoho, and P. Querre. Simultaneous cartoon and texture image inpainting using Morphological Component Analysis. *Applied and Comp. Harmonic Analysis*, 2005. to appear.
- [20] S. Esedoglu, S. Ruuth, and R. Tsai. Threshold dynamics for shape reconstruction and disocclusion. Technical report, UCLA CAM Report 05-22, April 2005. Accepted at IEEE Int. Conf. on Image Processing 2005.
- [21] S. Esedoglu and J. Shen. Digital image inpainting by the Mumford-Shah-Euler image model. *European J. Appl. Math.*, 13:353–370, 2002.

- [22] C. Fantoni and W. Gerbino. Contour interpolation by vector field combination. *Journal of Vision*, 3:281–303, 2003.
- [23] Y. Gousseau. *La distribution des formes dans les images naturelles*. PhD thesis, CEREMADE, Université Paris IX, 2000.
- [24] G. Kanizsa. *Organization in Vision: Essays on Gestalt Perception*. Präger, 1979.
- [25] S. Masnou. *Filtrage et Désocclusion d’Images par Méthodes d’Ensembles de Niveau*. PhD thesis, Université Paris-IX Dauphine, France, 1998.
- [26] S. Masnou. Disocclusion : a variational approach using level lines. *IEEE Trans. Image Processing*, 11:68–76, 2002.
- [27] S. Masnou. Euler spirals for image amodal completion. In preparation.
- [28] S. Masnou and J.-M. Morel. Level lines based disocclusion. In *5th IEEE Int. Conf. on Image Processing, Chicago, Illinois, October 4-7, 1998*.
- [29] G. Matheron. *Random Sets and Integral Geometry*. John Wiley, N.Y., 1975.
- [30] Y. Meyer. Oscillating patterns in image processing and nonlinear evolution equations. In Amer. Math. Soc., editor, *University Lecture Series*, volume Col. 22, 2001.
- [31] D. Mumford. Elastica and computer vision. In C.L. Bajaj, editor, *Algebraic Geometry and its Applications*, pages 491–506. Springer-Verlag, New-York, 1994.
- [32] M. Nitzberg and D. Mumford. The 2.1-D Sketch. In *Proc. 3rd Int. Conf. on Computer Vision*, pages 138–144, Osaka, Japan, 1990.
- [33] J. Petitot. Neurogeometry of V1 and Kanizsa contours. *Axiomathes*, 13:347–363, 2003.
- [34] L. Yaroslavsky and M. Eden. *Fundamentals of Digital Optics*. Birkhäuser, 1996.