

APPROXIMATION OF DYNAMIC AND QUASI-STATIC EVOLUTION PROBLEMS IN ELASTO-PLASTICITY BY CAP MODELS

JEAN-FRANÇOIS BABADJIAN AND MARIA GIOVANNA MORA

ABSTRACT. This work is devoted to the analysis of elasto-plasticity models arising in soil mechanics. Contrary to the typical models mainly used for metals, it is required here to take into account plastic dilatancy due to the sensitivity of granular materials to hydrostatic pressure. The yield criterion thus depends on the mean stress and the elasticity domain is unbounded and not invariant in the direction of hydrostatic matrices. In the mechanical literature, so-called *cap models* have been introduced, where the elasticity domain is cut in the direction of hydrostatic stresses by means of a strain-hardening yield surface, called a cap. The purpose of this article is to study the well-posedness of such models in dynamical and quasi-static regimes. An asymptotic analysis as the cap is moved to infinity is also performed, which enables one to recover solutions to the uncapped model of perfect elasto-plasticity.

1. INTRODUCTION

Models of elasto-plasticity have the capacity to predict the appearance of permanent deformations in a material when a critical stress is reached. From a microscopic point of view, these so-called plastic deformations are the result of atomic defects due to intercrystalline slips inside a lattice, called dislocations. It is experimentally observed that plastic flows occur on very thin zones called slip bands, on which there is strain localization: these zones are macroscopically interpreted as discontinuity surfaces of the displacement. For this reason, it has turned out to be convenient to approximate these models by regularized ones, *e.g.*, of viscosity or strain-hardening type. We refer to the monographs [18, 21] for an exhaustive presentation of elasto-plasticity models.

The mathematical models of plasticity are highly nonlinear and this makes difficult the search of solutions. However, the variational principles of Hodge-Prager for the stress rate, and of Greenberg for the velocity, enable one to formulate the problem in a more tractable way. In particular, the development of convex analysis and the interpretation of the model of perfect elasto-plasticity as the sweeping process of a moving convex set in [25] have permitted to prove existence and uniqueness of the stress history. Other mathematical results have been obtained in [14] by means of constructive theory of partial differential equations including Galerkin approximations, regularization, and penalization. The existence and uniqueness problem for the stress has been solved in a quite satisfactory way, while the evolution problem for the velocity (or the displacement) encountered additional difficulties connected to the regularity of the strain tensor. This problem was avoided in [19] by means of a weak formulation, which was however too weak to obtain full information on the strain. In the footsteps of these works, the quasi-static case was studied in [31, 32] and the dynamical case in [3, 22] by means of different types of visco-plastic regularizations (see also [2, 35]). The difficulty was connected to the definition of the correct functional setting for kinematically admissible displacement fields which can exhibit discontinuities. It has been overcome by the introduction in [24, 33] of the space BD of functions of bounded deformation (see [34] for

Date: August 24, 2012.

Key words and phrases. Elasto-plasticity, Functions of bounded deformation, Calculus of variations, Dynamic evolution, Quasi-static evolution, Convex analysis.

a comprehensive treatment on that subject). More recently, the quasi-static case was revisited in [7] within the general framework of variational evolutions of rate independent processes (see [23]).

To formulate more precisely the problem, let us consider a bounded open set $\Omega \subset \mathbb{R}^n$ (in dimension $n = 2$ or 3) which stands for the reference configuration of an elasto-plastic body. We will work in the framework of small strain elasto-plasticity where the natural kinematic variable is the displacement field $u : \Omega \times [0, T] \rightarrow \mathbb{R}^n$ (or the velocity field $v := \dot{u}$). We denote by $Eu := (Du + Du^T)/2$ the linearized strain tensor which takes its values in the set $\mathbb{M}_{sym}^{n \times n}$ of symmetric $n \times n$ matrices. In small strain elasto-plasticity, Eu decomposes additively in the following form:

$$Eu = e + p,$$

where $e : \Omega \times [0, T] \rightarrow \mathbb{M}_{sym}^{n \times n}$ is the elastic strain and $p : \Omega \times [0, T] \rightarrow \mathbb{M}_{sym}^{n \times n}$ the plastic strain. The elastic strain is related to the stress tensor $\sigma : \Omega \times [0, T] \rightarrow \mathbb{M}_{sym}^{n \times n}$ by means of Hooke's law $\sigma := \mathbb{C}e$, where \mathbb{C} is the symmetric fourth order elasticity tensor. In a dynamical framework and in the presence of an external body load $f : \Omega \times [0, T] \rightarrow \mathbb{R}^n$, the equation of motion writes

$$\ddot{u} - \operatorname{div} \sigma = f \quad \text{in } \Omega \times [0, T].$$

Plasticity is characterized by the existence of a yield zone beyond which permanent strains appear. The stress tensor is indeed constrained to belong to a fixed closed and convex subset K of $\mathbb{M}_{sym}^{n \times n}$:

$$\sigma \in K.$$

If σ lies inside the interior of K , the material behaves elastically, so that unloading will bring back the body into its initial configuration ($p = 0$). On the other hand, if σ reaches the boundary of K (called the yield surface), a plastic flow may develop, so that, after unloading, there will remain a non-trivial permanent plastic strain p . Its evolution is described by means of the flow rule and is expressed with the Prandtl-Reuss law

$$\dot{p} \in N_K(\sigma),$$

where $N_K(\sigma)$ is the normal cone to K at σ . From the theory of convex analysis, $N_K(\sigma) = \partial I_K(\sigma)$, *i.e.*, the subdifferential of the indicator function I_K of the set K ($I_K(\sigma) = 0$ if $\sigma \in K$, while $I_K(\sigma) = +\infty$ otherwise). Hence, from convex duality, the flow rule can be equivalently written as

$$\sigma : \dot{p} = \max_{\tau \in K} \tau : \dot{p} =: H(\dot{p}), \quad (1.1)$$

where $H : \mathbb{M}_{sym}^{n \times n} \rightarrow \mathbb{R}$ is the support function of K . This last formulation (1.1) is nothing but Hill's principle of maximum plastic work, and $H(\dot{p})$ denotes the plastic dissipation.

In general, the elasticity domain K is expressed by means of a yield function $F : \mathbb{M}_{sym}^{n \times n} \rightarrow \mathbb{R}$ as

$$K := \{\sigma \in \mathbb{M}_{sym}^{n \times n} : F(\sigma) \leq 0\}.$$

In this paper we assume that F is of the form

$$F(\sigma) := \alpha \sigma_m + \kappa(\sigma_D) - k,$$

where $\kappa : \mathbb{M}_D^{n \times n} \rightarrow [0, +\infty)$ is a convex and positively 1-homogeneous function with $\kappa(0) = 0$, and $\alpha > 0$ and $k > 0$ are positive constants related to the cohesion and the coefficient of internal friction of the material, respectively. Here, $\mathbb{M}_D^{n \times n} = \{\sigma \in \mathbb{M}_{sym}^{n \times n} : \operatorname{tr} \sigma = 0\}$ is the space of all deviatoric and symmetric $n \times n$ matrices,

$$\sigma_D := \sigma - \frac{\operatorname{tr} \sigma}{n} \operatorname{Id} \in \mathbb{M}_D^{n \times n} \quad \text{and} \quad \sigma_m := \frac{\operatorname{tr} \sigma}{n} \in \mathbb{R}$$

are respectively the deviatoric and spherical part of σ , so that $\sigma = \sigma_D + \sigma_m \operatorname{Id}$. Thus, the set K is actually a closed and convex cone with vertex on the axis of hydrostatic stresses. Note that if $\alpha = 0$, the yield function F does not depend on the mean stress σ_m and the set K is invariant in the direction of hydrostatic stresses. This is usually the case for most of metals and alloys for

which the influence of mean stress on yielding is generally negligible. This case has been studied in [7] in the quasi-static setting, and in [3, 22] in the dynamical one. A typical feature of these models is that, since the plastic strain is a deviatoric measure (recall that the displacement has bounded deformation), materials obeying this kind of laws do not develop plastic (or permanent) volumetric changes, and the displacement field admits only tangential discontinuities.

Here the function F do depend on the mean stress σ_m : in particular, the set K is not invariant and actually unbounded in the direction of hydrostatic matrices. It turns out that such yield criterions are necessary when it is desired to apply plasticity theory to soils, rocks, and concrete (see [13, 29]). Indeed, the essential property of such materials is that they are composed of many small particles. Consequently, permanent deformations and plastic slips occur when these particles slide over one another, and thus, as in ductile metals, failure occur primarily in shear. However, a strong difference with metals is that the shear strength is strongly influenced by the compressive normal stress acting on the shear plane, and therefore by the hydrostatic pressure. The physical reason of this phenomenon is connected to the fact that the void between the particles is composed of water and air. When the material is loaded in compression, the void ratio decreases in an irreversible way, leading to a permanent volume change. Therefore, the intergranular interaction is governed by a Coulomb type law of friction, where the shear and normal stresses achieve a critical combination on the shearing plane, depending on the angle of internal friction and the cohesion of the grains. In conclusion, the sensitivity on hydrostatic pressure as well as plastic dilatancy are typical features of this kind of granular materials.

Several well known models are recovered here. For instance, the *Drucker-Prager model* corresponds to

$$\kappa(\sigma_D) = |\sigma_D|,$$

while the *Mohr-Coulomb model* corresponds to

$$\kappa(\sigma_D) = \max_{i,j} \{(\sigma_D)_i - (\sigma_D)_j\},$$

where $(\sigma_D)_i, i = 1, \dots, n$, are the eigenvalues of σ_D . Note that when $\alpha = 0$, the Drucker-Prager criterion reduces to the Von Mises criterion, while the Mohr-Coulomb criterion reduces to that of Tresca.

All these models are called perfectly plastic, referring to the fact that the yield surface is fixed and does not move during the evolution. For more sophisticated models with a work-hardening material, the yield function may depend on an additional internal variable describing the position of the yield surface. Typical hardening rules are the isotropic hardening, representing a global uniform expansion of the elastic domain in all directions with no change in shape, and the kinematic hardening, representing a translation of the yield surface in stress space by shifting its reference point.

We are interested in studying the following model of dynamical evolution in perfect elasto-plasticity. Let $f : \Omega \times [0, T] \rightarrow \mathbb{R}^n$ be a given body force and $w : \partial\Omega \times [0, T] \rightarrow \mathbb{R}^n$ be a boundary displacement. We consider an initial datum $(u_0, e_0, p_0) : \Omega \rightarrow \mathbb{R}^n \times \mathbb{M}_{sym}^{n \times n} \times \mathbb{M}_{sym}^{n \times n}$ satisfying $Eu_0 = e_0 + p_0$ in Ω , $u_0 = w(0)$ on $\partial\Omega$, and $\sigma_0 := \mathbb{C}e_0 \in K$, and an initial velocity $v_0 : \Omega \rightarrow \mathbb{R}^n$ satisfying $v_0 = \dot{w}(0)$ on $\partial\Omega$. We look for a triplet $(u, e, p) : \Omega \times [0, T] \rightarrow \mathbb{R}^n \times \mathbb{M}_{sym}^{n \times n} \times \mathbb{M}_{sym}^{n \times n}$ with the properties

$$\begin{cases} Eu = e + p \text{ in } \Omega \times [0, T], & u = w \text{ on } \partial\Omega \times [0, T], \\ \sigma = \mathbb{C}e, & \sigma \in K \text{ in } \Omega \times [0, T], \\ \ddot{u} - \operatorname{div}\sigma = f \text{ in } \Omega \times [0, T], \\ \dot{p} \in N_K(\sigma) \text{ in } \Omega \times [0, T], \\ (u(0), e(0), p(0)) = (u_0, e_0, p_0), & \dot{u}(0) = v_0 \text{ in } \Omega. \end{cases} \quad (1.2)$$

For some geomaterials, it turns out that this kind of pressure-dependent models overestimate the yield stress and inadequately predict plastic dilatancy, which exceeds what is observed experimentally. In order to remedy these defects, a modified model has been introduced in [12], where the cone K is cut in the direction of hydrostatic stresses through the use of a strain-hardening yield surface or cap. In [27, 28] a Drucker-Prager cap model with a hardening law on the cap surface has been studied: if the stress reaches the cap, it is pushed forward in such a way that, if the stress gets the same position at some subsequent time, it will then be an interior point of the new elasticity domain. In our framework this corresponds to introduce an auxiliary variable ξ , related to the position of the cap, and to consider the yield functions $F_i : \mathbb{M}_{sym}^{n \times n} \times \mathbb{R} \rightarrow \mathbb{R}$, for $i = 1, 2, 3$, defined by

$$\begin{cases} F_1(\sigma, \xi) := F(\sigma), \\ F_2(\sigma, \xi) := \lambda\xi - \sigma_m, \\ F_3(\sigma, \xi) := \xi, \end{cases}$$

where $\lambda \geq 1$. These functions are clearly convex, and we define the closed convex set $K_\lambda \subset \mathbb{M}_{sym}^{n \times n} \times \mathbb{R}$ by

$$K_\lambda := \{(\sigma, \xi) \in \mathbb{M}_{sym}^{n \times n} \times \mathbb{R} : F_i(\sigma, \xi) \leq 0 \text{ for } i = 1, 2, 3\}.$$

The hardening cap model reads as follows: find a quadruplet $(u, e, p, \xi) : \Omega \times [0, T] \rightarrow \mathbb{R}^n \times \mathbb{M}_{sym}^{n \times n} \times \mathbb{M}_{sym}^{n \times n} \times \mathbb{R}$ satisfying

$$\begin{cases} Eu = e + p \text{ in } \Omega \times [0, T], & u = w \text{ on } \partial\Omega \times [0, T], \\ \sigma = \mathbb{C}e, \xi = -z, & (\sigma, \xi) \in K_\lambda \text{ in } \Omega \times [0, T], \\ \ddot{u} - \operatorname{div} \sigma = f \text{ in } \Omega \times [0, T], \\ (\dot{p}, \dot{z}) \in N_{K_\lambda}(\sigma, \xi) \text{ in } \Omega \times [0, T], \\ (u(0), e(0), p(0), \xi(0)) = (u_0, e_0, p_0, \xi_0), & \dot{u}(0) = v_0 \text{ in } \Omega. \end{cases} \quad (1.3)$$

Following [22, 3], the resolution of this dynamical problem can be performed by means of a vanishing viscosity method (see Theorems 3.1 and 4.1). In Theorem 5.1 we then prove existence and uniqueness of solutions for the uncapped dynamical problem (1.2), by showing convergence of the solution of (1.3) to a solution of (1.2), as $\lambda \rightarrow \infty$.

When the evolution is assumed to be slow, inertia terms (as thus the acceleration \ddot{u}) can be neglected, and the equations of motion in (1.2) and (1.3) become an equation of quasi-static equilibrium

$$-\operatorname{div} \sigma = f \quad \text{in } \Omega \times [0, T].$$

From a mathematical point of view, it may seem easier to deal with the quasi-static model instead of the dynamical one. Surprisingly, this observation turns out to be wrong. This is related to regularity issues of the stress and the displacement. Indeed it is known that the displacement can be discontinuous, and that the right functional setting to treat this problem is the space $BD(\Omega)$ of functions of bounded deformation. On the other hand, since the set K is bounded in no direction, the best integrability one can hope for the stress is $\sigma \in L^2(\Omega; \mathbb{M}_{sym}^{n \times n})$. Therefore, the flow rule (or equivalently Hills's principle of maximum plastic work) has to be written in a measure theoretic sense. Indeed, in (1.1) it is possible to define $H(\dot{p})$ using the theory of convex functions of a measure [17, 9, 10], while, according to [20, 16], it is also possible to give a sense to the stress/plastic strain duality product $\sigma : \dot{p}$ as a distribution (and even as a measure) by means of an integration by parts formula (see Definition 2.3 and formula (2.7) below). However, this definition makes sense provided σ and \dot{u} have enough space integrability. Indeed, taking smooth enough data, the natural regularities are either

$$\sigma \in L^n(\Omega; \mathbb{M}_{sym}^{n \times n}), \quad \dot{u} \in L^{n/(n-1)}(\Omega; \mathbb{R}^n),$$

or

$$\sigma \in L^2(\Omega; \mathbb{M}_{sym}^{n \times n}), \quad \dot{u} \in L^2(\Omega; \mathbb{R}^n). \tag{1.4}$$

In the dynamic problem the control of the kinetic energy gives us a natural $L^2(\Omega; \mathbb{R}^n)$ bound for the velocity \dot{u} , so that the conditions in (1.4) are always fulfilled and the stress/strain duality is well defined in any dimension. The quasi-static case is unfortunately less manageable. Indeed, except for planar elasto-plasticity ($n = 2$), where the previous alternatives are clearly equivalent, in higher dimension we only have $\sigma \in L^2(\Omega; \mathbb{M}_{sym}^{n \times n})$ and $\dot{u} \in L^{n/(n-1)}(\Omega; \mathbb{R}^n)$ by Sobolev embedding. Consequently, higher regularity results would be needed, either for the stress or the velocity. To the best of our knowledge, such results are not available in the literature. However, we can still give a meaning to the flow rule in the quasi-static setting by writing it in terms of an energy equality, which reduces to the usual flow rule when the solutions are smooth enough. Existence of solutions in this sense is proved in Theorems 6.3 and 6.5.

This obstruction on the dimension was already present in related works on the subject. In [26] the authors study a Hencky plasticity problem with a Mohr-Coulomb yield function¹ and consider a formulation within the framework of a minimax problem. As usual in elasto-plasticity, the associated Lagrangian is defined on a non-reflexive Banach space, so that existence of saddle points is not ensured. Consequently, the Lagrangian needs to be relaxed and this turns out to be possible only if there exists a statically and plastically admissible stress σ in $L^n(\Omega; \mathbb{M}_{sym}^{n \times n})$ (see [26, Lemma 3.2]). In the two-dimensional case, this condition is clearly satisfied provided the intersection between statically and plastically admissible stresses is not empty (since the stress is always at least squared integrable), while in the three-dimensional case the condition is in general not fulfilled. In the footsteps of this work, in [30] the author continues the study of the previous Mohr-Coulomb model by deriving a $H_{loc}^1(\Omega; \mathbb{M}_{sym}^{n \times n})$ regularity property for the stress. Note that such estimates are standard for yield functions independent of the mean stress (see [4, 11]). In order to get such a regularity property in the Mohr-Coulomb case, the author employs a visco-plastic regularization of the constitutive law, similar to that we used in Section 3. He shows that the $H_{loc}^1(\Omega; \mathbb{M}_{sym}^{n \times n})$ norm of the visco-plastic stress can be bounded in terms of the $L^2(\Omega; \mathbb{R}^n)$ norm of the visco-plastic displacement (see formula (3.33) in [30]). Unfortunately, this estimate is uniform (with respect to the viscosity parameter) only in dimension $n = 2$.

Let us now comment on our choice of boundary conditions: we only impose Dirichlet boundary conditions, which correspond to a hard device applied to the whole boundary. The case of a Neumann condition (even on a portion of the boundary) seems to be difficult to carry out, and the reason is again connected to regularity issues. Indeed, here $\sigma \in L^2(\Omega; \mathbb{M}_{sym}^{n \times n})$ with $\text{div} \sigma \in L^2(\Omega; \mathbb{R}^n)$ (respectively, $\text{div} \sigma \in L^n(\Omega; \mathbb{R}^n)$ in the quasi-static case) so that the normal stress $\sigma \nu$ is *a priori* defined as an element of $H^{-1/2}(\partial\Omega; \mathbb{R}^n)$. On the other hand, a trace theorem in $BD(\Omega)$ asserts that kinematically admissible displacements u have a trace (still denoted by u) in $L^1(\partial\Omega; \mathbb{R}^n)$. Consequently, the displacement and the stress are not in duality on the boundary. In [7] and [3] this problem was avoided owing to a result of [20]: since the set K is bounded in the direction of deviatoric matrices, only the tangential part of the normal stress is relevant on the boundary, which turns out to belong to $L^\infty(\partial\Omega; \mathbb{R}^n)$. Consequently, in that case, (the tangential part of) $\sigma \nu$ and u are in duality on the boundary and this makes possible to take into account traction loads on a portion of $\partial\Omega$.

As a consequence, the only forces applied to our system are body loads, and, as usual in plasticity, we must ensure that the space of statically and plastically admissible stresses is not empty. In the quasi-static case, as observed in [25, 19], a stronger hypothesis is needed, namely a safe-load condition: the forces must derive from a potential χ well contained in the set K (see (6.1) and (6.2)). This safety condition, which ensures that the body is not in a free flow, is necessary, from

¹In our terminology, the model studied in [26] actually corresponds to a Drucker-Prager model.

a mathematical point of view, in order to obtain an *a priori* estimate on the plastic strain rate. In the dynamical case we observe that this safe-load condition is no longer necessary thanks to the presence of the kinetic energy which is of higher order than the work of external forces.

The paper is organized as follows. In Section 2 we collect the main notation and introduce the different energies involved in our model; in particular, we define the dissipation functionals as convex functionals of measures and in terms of the stress/strain duality. In Section 3 we prove existence and uniqueness of solutions to a visco-plastic regularization of the dynamical cap model, similar to that of [22]. Our approach uses an implicit finite difference approximation of the underlying hyperbolic system. In Section 4 we perform a vanishing viscosity analysis in order to get existence and uniqueness of solutions of the dynamical elasto-plastic cap model. Section 5 is devoted to the asymptotic analysis of the previous cap model as the position of the cap is sent to infinity. In particular, we show existence and uniqueness of solutions to the dynamical elasto-plastic uncapped model. Eventually, in Section 6 we review the quasi-static case: in the framework of planar elasto-plasticity, we show existence of solutions to the quasi-static cap model, as well as its convergence when the cap is sent to infinity. In higher dimension existence is proved for a weaker formulation of the problem, where the flow rule is replaced by an energy equality.

2. THE MATHEMATICAL SETTING

Throughout the paper, Ω is a bounded connected open set in \mathbb{R}^n with Lipschitz boundary. The Lebesgue measure in \mathbb{R}^n and the $(n-1)$ -dimensional Hausdorff measure are denoted by \mathcal{L}^n and \mathcal{H}^{n-1} , respectively.

We use standard notation for Lebesgue and Sobolev spaces. In particular, for $1 \leq p \leq \infty$, the $L^p(\Omega)$ -norms of the various quantities are denoted by $\|\cdot\|_p$.

The notation \odot stands for the symmetrized tensor product between vectors in \mathbb{R}^n .

Given a locally compact set $E \subset \mathbb{R}^n$ and a Euclidean space X , we denote by $\mathcal{M}(E; X)$ (or simply $\mathcal{M}(E)$ if $X = \mathbb{R}$) the space of bounded Radon measures on E with values in X , endowed with the norm $\|\mu\|_1 := |\mu|(E)$, where $|\mu| \in \mathcal{M}(E)$ is the total variation of the measure μ . The Riesz Representation Theorem ensures that $\mathcal{M}(E; X)$ can be identified with the dual of $\mathcal{C}_0(E; X)$, the space of continuous functions $\varphi : E \rightarrow X$ vanishing on the boundary of E , *i.e.*, such that $\{|\varphi| \geq \varepsilon\}$ is compact for any $\varepsilon > 0$. For $\mu \in \mathcal{M}(E; X)$ we consider the Lebesgue decomposition $\mu = \mu^a + \mu^s$, where μ^a is absolutely continuous and μ^s is singular with respect to the Lebesgue measure \mathcal{L}^n . Moreover, if ν is a non-negative Radon measure over E , we denote by $\frac{d\mu}{d\nu}$ the Radon-Nikodym derivative of μ with respect to ν .

Finally, $BD(\Omega)$ stands for the space of functions of bounded deformation in Ω , *i.e.*, $u \in BD(\Omega)$ if $u \in L^1(\Omega; \mathbb{R}^n)$ and $Eu \in \mathcal{M}(\Omega; \mathbb{M}_{sym}^{n \times n})$, where $Eu := (Du + Du^T)/2$ and Du is the distributional derivative of u . We refer to [34] for general properties of this space.

2.1. The elastic energy. Let \mathbb{C} be a fourth order tensor satisfying the usual symmetry conditions

$$\mathbb{C}_{ijkl} = \mathbb{C}_{klij} = \mathbb{C}_{jikl} \quad \text{for all } i, j, k, l \in \{1, \dots, n\}.$$

We assume that there exist constants $0 < \alpha_{\mathbb{C}} \leq \beta_{\mathbb{C}} < \infty$ such that

$$\alpha_{\mathbb{C}}|e|^2 \leq \mathbb{C}e : e \leq \beta_{\mathbb{C}}|e|^2 \quad \text{for all } e \in \mathbb{M}_{sym}^{n \times n}. \quad (2.1)$$

We define the elastic energy, for all $e \in L^2(\Omega; \mathbb{M}_{sym}^{n \times n})$, by

$$\mathcal{Q}(e) := \frac{1}{2} \int_{\Omega} \mathbb{C}e(x) : e(x) dx.$$

2.2. The dissipation energies. Let $\kappa : \mathbb{M}_D^{n \times n} \rightarrow [0, +\infty)$ be a convex and positively 1-homogeneous function with $\kappa(0) = 0$, and let

$$K := \{\sigma \in \mathbb{M}_{sym}^{n \times n} : \alpha \sigma_m + \kappa(\sigma_D) - k \leq 0\},$$

where $\alpha > 0$ and $k > 0$ are given constants. For every $\lambda \geq 1$ let

$$K_\lambda := \{(\sigma, \xi) \in K \times \mathbb{R}^- : \lambda \xi - \sigma_m \leq 0\}. \quad (2.2)$$

We define the support functions $H : \mathbb{M}_{sym}^{n \times n} \rightarrow [0, +\infty]$ and $H_\lambda : \mathbb{M}_{sym}^{n \times n} \times \mathbb{R} \rightarrow [0, +\infty]$ of K and K_λ by

$$H(p) := \sup_{\sigma \in K} \sigma : p \quad \text{and} \quad H_\lambda(p, z) := \sup_{(\sigma, \xi) \in K_\lambda} \{\sigma : p + \xi z\}.$$

Since K and K_λ are closed and convex, then H and H_λ are convex, lower semicontinuous, and positively 1-homogeneous.

The following result states some relations between H and H_λ .

Lemma 2.1. *For any $(p, z) \in \mathbb{M}_{sym}^{n \times n} \times \mathbb{R}$, we have*

$$\sup_{\lambda \geq 1} H_\lambda(p, z) = H(p) + I_{\mathbb{R}^+}(z),$$

where $I_{\mathbb{R}^+}$ is the indicator function of \mathbb{R}^+ , i.e., $I_{\mathbb{R}^+}(z) = 0$ if $z \geq 0$, while $I_{\mathbb{R}^+}(z) = +\infty$ otherwise.

Proof. By definition of K_λ , if $1 \leq \lambda_1 \leq \lambda_2$, then $K_{\lambda_1} \subset K_{\lambda_2}$ and consequently $H_{\lambda_1} \leq H_{\lambda_2}$. Defining $K_\infty := \cup_{\lambda \geq 1} K_\lambda$, we have that for every $(p, z) \in \mathbb{M}_{sym}^{n \times n} \times \mathbb{R}$ and $\lambda \geq 1$,

$$H_\lambda(p, z) \leq H_\infty(p, z) := \sup_{(\sigma, \xi) \in K_\infty} \{\sigma : p + \xi z\} = H(p) + I_{\mathbb{R}^+}(z),$$

where the last equality follows from the fact that $\overline{K}_\infty = K \times \mathbb{R}^-$. Hence we deduce that for any $(p, z) \in \mathbb{M}_{sym}^{n \times n} \times \mathbb{R}$,

$$\sup_{\lambda \geq 1} H_\lambda(p, z) \leq H(p) + I_{\mathbb{R}^+}(z).$$

To prove the converse inequality, assume that $\sup_{\lambda} H_\lambda(p, z) < \infty$. If $(\sigma, \xi) \in K_\infty$, then $(\sigma, \xi) \in K_\lambda$ for some λ , and thus

$$\sup_{\lambda \geq 1} H_\lambda(p, z) \geq \sigma : p + \xi z.$$

Maximizing with respect to $(\sigma, \xi) \in K_\infty$ in the right-handside of the previous inequality yields

$$\sup_{\lambda \geq 1} H_\lambda(p, z) \geq H(p) + I_{\mathbb{R}^+}(z),$$

and the proof is complete. \square

The functions H and H_λ enjoy a nice coercivity property. Indeed, since $\kappa(0) = 0$, then 0 is an interior point of K , and thus, there exists $\alpha_H > 0$ such that $B(0, \alpha_H) \subset K$, from which we deduce that

$$\alpha_H |p| \leq H(p) \quad \text{for all } p \in \mathbb{M}_{sym}^{n \times n}.$$

Moreover, since $B(0, \alpha_H) \times (-\infty, -\frac{\alpha_H}{\sqrt{n}}] \subset K_1 \subset K_\lambda$ for all $\lambda \geq 1$, then

$$\alpha_H |p| - \frac{\alpha_H}{\sqrt{n}} z \leq H_\lambda(p, z) \quad \text{for all } (p, z) \in \mathbb{M}_{sym}^{n \times n} \times \mathbb{R}^+, \quad (2.3)$$

and

$$z < 0 \quad \text{implies that} \quad H_\lambda(p, z) = +\infty. \quad (2.4)$$

The dissipated energy is then defined, for all $(p, z) \in L^2(\Omega; \mathbb{M}_{sym}^{n \times n}) \times L^2(\Omega)$, by

$$\mathcal{H}_\lambda(p, z) := \int_{\Omega} H_\lambda(p(x), z(x)) \, dx.$$

As a consequence of the previous properties of H_λ , we infer that \mathcal{H}_λ is sequentially weakly lower semicontinuous in $L^2(\Omega; \mathbb{M}_{sym}^{n \times n}) \times L^2(\Omega)$. It will also be useful to extend the definition of \mathcal{H}_λ when $p \in \mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n})$. According to [17], we define the non-negative Borel measures

$$H(p)(B) := \int_B H \left(\frac{dp}{d|p|}(x) \right) d|p|(x)$$

and

$$H_\lambda(p, z)(B) := \int_B H_\lambda \left(\frac{dp}{d\mathcal{L}^n}(x), z(x) \right) dx + \int_B H_\lambda \left(\frac{dp}{d|p^s|}(x), 0 \right) d|p^s|(x)$$

for any Borel set $B \subset \bar{\Omega}$. In general, by [17] the measures $H(p)$ and $H_\lambda(p, z)$ are not even locally finite. However, if further $H(p)$ and $H_\lambda(p, z)$ have finite mass, *i.e.*, if $H(p)$ and $H_\lambda(p, z)$ are bounded Radon measures, we can define the functionals

$$\mathcal{H}(p) := H(p)(\bar{\Omega}) \quad \text{and} \quad \mathcal{H}_\lambda(p, z) := H_\lambda(p, z)(\bar{\Omega}).$$

In that case, the results of [9, 10] apply and these convex functions of measures can be expressed by means of duality formulas. To this aim, let us extend p and z by zero on $\Omega' \setminus \bar{\Omega}$, where $\Omega' \subset \mathbb{R}^n$ is a bounded smooth open set containing $\bar{\Omega}$. As a consequence, $H(p)$ and $H_\lambda(p, z)$ are in $\mathcal{M}(\Omega')$ and thanks to [10, Theorem 2.1–(ii)], we get that

$$\begin{aligned} \int_{\Omega'} \varphi d[H(p)] &= \sup \left\{ \int_{\Omega'} \varphi \sigma : dp \quad : \sigma \in \mathcal{C}_c^\infty(\Omega'; K) \right\}, \\ \int_{\Omega'} \varphi d[H_\lambda(p, z)] &= \sup \left\{ \int_{\Omega'} \varphi \sigma : dp + \int_{\Omega'} \xi z \varphi dx \quad : (\sigma, \xi) \in \mathcal{C}_c^\infty(\Omega'; K_\lambda) \right\}, \end{aligned} \quad (2.5)$$

for any $\varphi \in \mathcal{C}_c(\Omega')$ with $\varphi \geq 0$, and in particular

$$\begin{aligned} \mathcal{H}(p) &= \sup \left\{ \int_{\bar{\Omega}} \sigma : dp \quad : \sigma \in \mathcal{C}^\infty(\bar{\Omega}; K) \right\}, \\ \mathcal{H}_\lambda(p, z) &= \sup \left\{ \int_{\bar{\Omega}} \sigma : dp + \int_{\bar{\Omega}} \xi z dx \quad : (\sigma, \xi) \in \mathcal{C}^\infty(\bar{\Omega}; K_\lambda) \right\}. \end{aligned} \quad (2.6)$$

Note also that Reshetnyak Theorem (see [1, Theorem 2.38]) still holds here, so that \mathcal{H} and \mathcal{H}_λ are sequentially weakly* lower semicontinuous in $\mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n})$ and $\mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n}) \times L^2(\Omega)$, respectively.

2.3. The total dissipation. Let $(p, z) : [0, T] \rightarrow \mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n}) \times L^2(\Omega)$. The \mathcal{H}_λ -variation of (p, z) on a time interval $[a, b]$, which will play the role of the total dissipation, is defined as

$$\begin{aligned} \mathcal{D}_\lambda(p, z; [a, b]) &:= \sup \left\{ \sum_{j=1}^N \mathcal{H}_\lambda(p(t_j) - p(t_{j-1}), z(t_j) - z(t_{j-1})) : \right. \\ &\quad \left. a = t_0 \leq t_1 \leq \dots \leq t_N = b, N \in \mathbb{N} \right\}. \end{aligned}$$

Since \mathcal{H}_λ is sequentially weakly* lower semicontinuous in $\mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n}) \times L^2(\Omega)$, we deduce that $\mathcal{D}_\lambda(\cdot, \cdot; [a, b])$ is sequentially weakly* lower semicontinuous in $\mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n}) \times L^2(\Omega)$ as well.

The following result enables one to write the total dissipation \mathcal{D}_λ as a time integral when p and z are regular enough with respect to time. This property follows from an adaptation of [7, Theorem 7.1] (see also [5, Appendix]). Indeed, a careful inspection of the proof of that result shows that it is enough to have the functional \mathcal{H}_λ lower semicontinuous, which is ensured in our case by Reshetnyak Theorem. In particular, the fact that the function H_λ can take the value $+\infty$ does not affect the validity of the result.

Proposition 2.2. *Assume that $p \in AC([0, T]; \mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n}))$, $z \in AC([0, T]; L^2(\Omega))$, and*

$$\mathcal{D}_\lambda(p, z; [0, T]) < +\infty.$$

Then, for a.e. $t \in [0, T]$, there exist $(\dot{p}(t), \dot{z}(t)) \in \mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n}) \times L^2(\Omega)$ such that

$$\begin{cases} \frac{p(s) - p(t)}{s - t} \rightharpoonup \dot{p}(t) \text{ weakly}^* \text{ in } \mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n}), \\ \frac{z(s) - z(t)}{s - t} \rightarrow \dot{z}(t) \text{ strongly in } L^2(\Omega), \end{cases} \quad \text{as } s \rightarrow t.$$

Moreover, the function $t \mapsto \mathcal{H}_\lambda(\dot{p}(t), \dot{z}(t))$ is measurable and for all $0 \leq a \leq b \leq T$,

$$\mathcal{D}_\lambda(p, z; [a, b]) = \int_a^b \mathcal{H}_\lambda(\dot{p}(t), \dot{z}(t)) dt.$$

2.4. Duality between the stress and the plastic strain. The duality pairing between stresses and plastic strains is *a priori* not well defined, since the former are only squared Lebesgue integrable, while the latter are measures. This is clearly an obstacle if one wishes to express in a pointwise sense Hill's principle of maximum plastic work (1.1). Using an integration by parts formula as in [20, 16], it is actually possible to give a sense to this duality pairing as a distribution, and even as a measure for solutions of the elasto-plasticity model.

Definition 2.3. Let $\sigma \in L^2(\Omega; \mathbb{M}_{sym}^{n \times n})$ with $\operatorname{div} \sigma \in L^2(\Omega; \mathbb{R}^n)$, and let $p \in \mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n})$ be such that there exists a triple $(u, e, w) \in (BD(\Omega) \cap L^2(\Omega; \mathbb{R}^n)) \times L^2(\Omega; \mathbb{M}_{sym}^{n \times n}) \times H^1(\Omega; \mathbb{R}^n)$ satisfying $Eu = e + p$ in Ω and $p = (w - u) \odot \nu \mathcal{H}^{n-1}$ on $\partial\Omega$, where ν is the outer unit normal to $\partial\Omega$. We define the distribution $[\sigma : p]$ on \mathbb{R}^n as

$$\langle [\sigma : p], \varphi \rangle = \int_\Omega \varphi(w - u) \cdot \operatorname{div} \sigma dx + \int_\Omega \sigma : [(w - u) \odot \nabla \varphi] dx + \int_\Omega \sigma : (Ew - e) \varphi dx$$

for every $\varphi \in C_c^\infty(\mathbb{R}^n)$.

It is easy to check that the definition of $[\sigma : p]$ is independent of the choice of (u, e, w) , so that the distribution $[\sigma : p]$ is well defined. Moreover, since $[\sigma : p]$ is a distribution supported in $\bar{\Omega}$, we can define the duality product $\langle \sigma, p \rangle$ as

$$\langle \sigma, p \rangle := \langle [\sigma : p], 1 \rangle = \int_\Omega (w - u) \cdot \operatorname{div} \sigma dx + \int_\Omega \sigma : (Ew - e) dx. \quad (2.7)$$

Remark 2.4. Note that, contrary to [20, 16], here the stress σ may not be in $L^\infty(\Omega; \mathbb{M}_{sym}^{n \times n})$; therefore, it is not clear how to prove at this stage that the distribution $[\sigma : p]$ is actually a measure. However, this property will be obtained afterwards for solutions of the plasticity problems (see Theorems 4.1, 5.1, 6.4, and 6.6).

Using this notion of stress/strain duality, the duality formulas (2.5) and (2.6) can be now extended to less regular statically and plastically admissible stresses. This property rests on a density result, together with the fact that, when the stress is smooth enough, the stress/strain duality reduces to the usual duality between continuous functions and measures. Indeed, according to the integration by parts formula in $BD(\Omega)$, if $\sigma \in C^1(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n})$, we have

$$\langle [\sigma : p], \varphi \rangle = \int_{\bar{\Omega}} \varphi \sigma : dp \quad \text{for all } \varphi \in C_c^\infty(\mathbb{R}^n), \quad (2.8)$$

and

$$\langle \sigma, p \rangle = \int_{\bar{\Omega}} \sigma : dp. \quad (2.9)$$

Moreover, the following approximation result is an immediate adaptation of [7, Lemma 2.3].

Lemma 2.5. *Let $1 \leq p < \infty$ and $(\sigma, \xi) \in L^p(\Omega; \mathbb{M}_{sym}^{n \times n}) \times L^2(\Omega)$ be such that $\operatorname{div} \sigma \in L^p(\Omega; \mathbb{R}^n)$ and $(\sigma(x), \xi(x)) \in K_\lambda$ for a.e. $x \in \Omega$. There exists a sequence $(\sigma_k, \xi_k) \subset C^\infty(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n}) \times C^\infty(\bar{\Omega})$ such that $(\sigma_k, \xi_k) \rightarrow (\sigma, \xi)$ strongly in $L^p(\Omega; \mathbb{M}_{sym}^{n \times n}) \times L^2(\Omega)$, $\operatorname{div} \sigma_k \rightarrow \operatorname{div} \sigma$ strongly in $L^p(\Omega; \mathbb{R}^n)$, and $(\sigma_k(x), \xi_k(x)) \in K_\lambda$ for all $x \in \Omega$ and all $k \in \mathbb{N}$.*

If $p \in \mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n})$ with $\mathcal{H}(p) < +\infty$, then, using the duality formulas (2.5) and (2.6), together with (2.8), (2.9), and the approximation result [7, Lemma 2.3], we infer that for all $\varphi \in C_c^\infty(\Omega')$ with $\varphi \geq 0$,

$$\int_{\Omega'} \varphi d[H(p)] = \sup \left\{ \langle [\sigma : p], \varphi \rangle : \sigma \in L^2(\Omega; \mathbb{M}_{sym}^{n \times n}) \text{ with } \operatorname{div} \sigma \in L^2(\Omega; \mathbb{R}^n) \right. \\ \left. \text{and } \sigma(x) \in K \text{ for a.e. } x \in \Omega \right\}, \quad (2.10)$$

and in particular,

$$\mathcal{H}(p) = \sup \left\{ \langle \sigma, p \rangle : \sigma \in L^2(\Omega; \mathbb{M}_{sym}^{n \times n}) \text{ with } \operatorname{div} \sigma \in L^2(\Omega; \mathbb{R}^n) \right. \\ \left. \text{and } \sigma(x) \in K \text{ for a.e. } x \in \Omega \right\}. \quad (2.11)$$

Analogously, if $p \in \mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n})$, $z \in L^2(\Omega)$, and $\mathcal{H}_\lambda(p, z) < +\infty$, then, by (2.5), (2.6), (2.8), (2.9), and Lemma 2.5, we infer that for all $\varphi \in C_c^\infty(\Omega')$ with $\varphi \geq 0$,

$$\int_{\Omega'} \varphi d[H_\lambda(p, z)] = \sup \left\{ \langle [\sigma : p], \varphi \rangle + \int_{\Omega} \xi z \varphi dx : (\sigma, \xi) \in L^2(\Omega; \mathbb{M}_{sym}^{n \times n}) \times L^2(\Omega) \text{ with} \right. \\ \left. \operatorname{div} \sigma \in L^2(\Omega; \mathbb{R}^n) \text{ and } (\sigma(x), \xi(x)) \in K_\lambda \text{ for a.e. } x \in \Omega \right\}, \quad (2.12)$$

and in particular,

$$\mathcal{H}_\lambda(p, z) = \sup \left\{ \langle \sigma, p \rangle + \int_{\Omega} \xi z dx : (\sigma, \xi) \in L^2(\Omega; \mathbb{M}_{sym}^{n \times n}) \times L^2(\Omega) \text{ with} \right. \\ \left. \operatorname{div} \sigma \in L^2(\Omega; \mathbb{R}^n) \text{ and } (\sigma(x), \xi(x)) \in K_\lambda \text{ for a.e. } x \in \Omega \right\}. \quad (2.13)$$

Remark 2.6. Note that, if p is associated to a displacement $u \in BD(\Omega) \setminus L^2(\Omega; \mathbb{R}^n)$, we can still define the distribution $[\sigma : p]$ for $\sigma \in L^n(\Omega; \mathbb{M}_{sym}^{n \times n})$ with $\operatorname{div} \sigma \in L^n(\Omega; \mathbb{R}^n)$, because of the embedding of $BD(\Omega)$ into $L^{n/(n-1)}(\Omega; \mathbb{R}^n)$. In that case, formulas (2.10)–(2.13) hold with supremum taken over all $\sigma \in L^n(\Omega; \mathbb{M}_{sym}^{n \times n})$ satisfying $\operatorname{div} \sigma \in L^n(\Omega; \mathbb{R}^n)$. Nevertheless, the definition of the duality will be a source of difficulties when dealing with the quasi-static case in dimension higher than two, because the velocity and/or the stress may miss the required integrability (see Theorems 6.4 and 6.6).

3. THE DYNAMICAL VISCO-PLASTIC CAP MODEL

The main result of this section is an existence result for a visco-plastic dynamical cap model. The kind of viscosity we use is not related to a regularization of the flow rule of Perzyna or Norton-Hoff type as in [31, 32, 35], but rather connected to the constitutive law. We assume that the (visco-plastic) stress $\tilde{\sigma} = \mathbb{C}e + \varepsilon E\dot{u}$ is the sum of two terms. The first part $\sigma := \mathbb{C}e$ is the stress that originates from the elastic reaction to the deformation, while the second part $\varepsilon E\dot{u}$ is a damping term due to viscosity ($\varepsilon > 0$ is a viscosity coefficient). The model described below (Theorem 3.1) is similar to that studied in [22]. In that reference, existence and uniqueness were proved by means of a Galerkin method, while here, we employ a time discretization procedure of the underlying hyperbolic system. However, our proofs are very close to those of [22].

Let us define the set of visco-plastic kinematically admissible fields in the following way: given a boundary displacement $\hat{w} \in H^1(\Omega; \mathbb{R}^n)$,

$$\mathcal{A}_{\text{vp}}(\hat{w}) := \{(v, \eta, q) \in H^1(\Omega; \mathbb{R}^n) \times L^2(\Omega; \mathbb{M}_{\text{sym}}^{n \times n}) \times L^2(\Omega; \mathbb{M}_{\text{sym}}^{n \times n}) : \\ Ev = \eta + q \text{ a.e. in } \Omega, v = \hat{w} \mathcal{H}^{n-1}\text{-a.e. on } \partial\Omega\}.$$

We fix $\lambda \geq 1$ and we consider a body load $f \in AC([0, T]; L^2(\Omega; \mathbb{R}^n))$ and a boundary displacement which is the trace on $\partial\Omega \times (0, T)$ of a function $w \in H^2([0, T]; H^1(\Omega; \mathbb{R}^n)) \cap H^3([0, T]; L^2(\Omega; \mathbb{R}^n))$. Moreover, let $(u_0, e_0, p_0, z_0) \in \mathcal{A}_{\text{vp}}(w(0)) \times L^2(\Omega)$ and $v_0 \in H^1(\Omega; \mathbb{R}^n)$ be initial data satisfying

$$v_0 = \dot{w}(0) \mathcal{H}^{n-1}\text{-a.e. on } \partial\Omega$$

and

$$(\sigma_0, \xi_0) := (\mathbb{C}e_0, -z_0) \in K_\lambda, \quad -\text{div}\sigma_0 = f(0) \quad \text{a.e. in } \Omega. \quad (3.1)$$

Theorem 3.1. *Let $\varepsilon > 0$. There exist unique*

$$\begin{cases} u_\varepsilon \in W^{1,\infty}([0, T]; L^2(\Omega; \mathbb{R}^n)) \cap H^1([0, T]; H^1(\Omega; \mathbb{R}^n)), \text{ with } \dot{u}_\varepsilon \in L^2(0, T; H^{-1}(\Omega; \mathbb{R}^n)), \\ \sigma_\varepsilon, e_\varepsilon \in H^1([0, T]; L^2(\Omega; \mathbb{M}_{\text{sym}}^{n \times n})), \\ p_\varepsilon \in H^1([0, T]; L^2(\Omega; \mathbb{M}_{\text{sym}}^{n \times n})), \\ \xi_\varepsilon, z_\varepsilon \in H^1([0, T]; L^2(\Omega)), \end{cases}$$

with the following properties: for all $t \in [0, T]$,

$$\begin{cases} Eu_\varepsilon(t) = e_\varepsilon(t) + p_\varepsilon(t) \text{ a.e. in } \Omega, & u_\varepsilon(t) = w(t) \mathcal{H}^{n-1}\text{-a.e. on } \partial\Omega, \\ \sigma_\varepsilon(t) = \mathbb{C}e_\varepsilon(t), & \xi_\varepsilon(t) = -z_\varepsilon(t), \\ (\sigma_\varepsilon(t), \xi_\varepsilon(t)) \in K_\lambda \text{ a.e. in } \Omega, \end{cases} \quad (3.2)$$

and

$$\begin{cases} \ddot{u}_\varepsilon - \text{div}(\sigma_\varepsilon + \varepsilon E\dot{u}_\varepsilon) = f & \text{in } L^2(0, T; H^{-1}(\Omega; \mathbb{R}^n)), \\ (u_\varepsilon(0), e_\varepsilon(0), p_\varepsilon(0), z_\varepsilon(0)) = (u_0, e_0, p_0, z_0), & \dot{u}_\varepsilon(0) = v_0. \end{cases} \quad (3.3)$$

Moreover, for a.e. $t \in [0, T]$,

$$\dot{z}_\varepsilon(t) \geq 0 \quad \text{and} \quad (\dot{p}_\varepsilon(t), \dot{z}_\varepsilon(t)) \in N_{K_\lambda}(\sigma_\varepsilon(t), \xi_\varepsilon(t)) \quad \text{a.e. in } \Omega. \quad (3.4)$$

If further $v_0 \in H^2(\Omega; \mathbb{R}^n)$, then

$$\begin{cases} u_\varepsilon \in W^{2,\infty}([0, T]; L^2(\Omega; \mathbb{R}^n)) \cap H^2([0, T]; H^1(\Omega; \mathbb{R}^n)), \\ \sigma_\varepsilon, e_\varepsilon \in W^{1,\infty}([0, T]; L^2(\Omega; \mathbb{M}_{\text{sym}}^{n \times n})), \\ \xi_\varepsilon, z_\varepsilon \in W^{1,\infty}([0, T]; L^2(\Omega)), \end{cases}$$

and the equation of motion holds a.e. in $\Omega \times (0, T)$.

Remark 3.2. Note that (3.2) and (3.4) ensure that the map $t \mapsto \xi_\varepsilon(t)$ is non-increasing. Since ξ_ε is an internal variable describing the position of the cap (see (2.2)), this property means that the cap is moving outwards as time proceeds. This is exactly the strain-hardening phenomenon we wanted to highlight on the evolution of the cap surface.

3.1. Time discretization. The proof of Theorem 3.1 relies on a time discretization procedure. Let us consider a partition of the time interval $[0, T]$ into N_k sub-intervals of equal length δ_k :

$$0 = t_k^0 < t_k^1 < \dots < t_k^{N_k} = T, \quad \text{with} \quad \delta_k := \frac{T}{N_k} = t_k^i - t_k^{i-1} \rightarrow 0.$$

We define the discrete body loads by $f_k^i := f(t_k^i)$ for all $i \in \{0, \dots, N_k\}$ and the discrete boundary values $\{w_k^i\}_{0 \leq i \leq N_k}$ by

$$w_k^0 := w(0), \quad w_k^1 = w(0) + \delta_k \dot{w}(0), \quad \text{and} \quad w_k^i = w(t_k^i) \text{ for all } i \in \{2, \dots, N_k\}.$$

Then we define inductively

$$(u_k^0, e_k^0, p_k^0, z_k^0) := (u_0, e_0, p_0, z_0), \quad (u_k^1, e_k^1, p_k^1, z_k^1) := (u_0, e_0, p_0, z_0) + \delta_k(v_0, 0, Ev_0, 0), \quad (3.5)$$

and, for all $i \in \{2, \dots, N_k\}$, we define $(u_k^i, e_k^i, p_k^i, z_k^i)$ as the solution of the following minimum problem

$$\min_{(v, \eta, q, \zeta) \in \mathcal{A}_{\text{vp}}(w_k^i) \times L^2(\Omega)} \left\{ \mathcal{Q}(\eta) + \frac{\varepsilon}{2\delta_k} \|Ev - Eu_k^{i-1}\|_2^2 + \frac{1}{2\delta_k^2} \|v - 2u_k^{i-1} + u_k^{i-2}\|_2^2 + \frac{1}{2} \|\zeta\|_2^2 + \mathcal{H}_\lambda(q - p_k^{i-1}, \zeta - z_k^{i-1}) - \int_\Omega f_k^i \cdot v \, dx \right\}. \quad (3.6)$$

Korn's inequality together with the sequential weak lower semicontinuity in $L^2(\Omega; \mathbb{M}_{sym}^{n \times n}) \times L^2(\Omega)$ of \mathcal{H}_λ imply that the previous minimum problem admits a solution denoted by $(u_k^i, e_k^i, p_k^i, z_k^i) \in \mathcal{A}_{\text{vp}}(w_k^i) \times L^2(\Omega)$, which is unique by strict convexity of the functional. In particular, since $\mathcal{H}_\lambda(p_k^i - p_k^{i-1}, z_k^i - z_k^{i-1}) < \infty$, we deduce from (2.4) that

$$z_k^i \geq z_k^{i-1} \quad \text{a.e. in } \Omega. \quad (3.7)$$

We now derive the Euler-Lagrange equation of the previous minimum problem.

Proposition 3.3. *Let $(u_k^i, e_k^i, p_k^i, z_k^i) \in \mathcal{A}_{\text{vp}}(w_k^i) \times L^2(\Omega)$ be defined by (3.5) and (3.6). Then, for all $i \in \{0, \dots, N_k\}$,*

$$\begin{cases} Eu_k^i = e_k^i + p_k^i \text{ a.e. in } \Omega, & u_k^i = w_k^i \text{ } \mathcal{H}^{n-1}\text{-a.e. on } \partial\Omega, \\ \sigma_k^i := \mathbb{C}e_k^i, \xi_k^i := -z_k^i, & (\sigma_k^i, \xi_k^i) \in K_\lambda \text{ a.e. in } \Omega, \end{cases}$$

and for all $i \in \{2, \dots, N_k\}$,

$$\begin{cases} \frac{u_k^i - 2u_k^{i-1} + u_k^{i-2}}{\delta_k^2} - \operatorname{div} \left(\sigma_k^i + \varepsilon \frac{Eu_k^i - Eu_k^{i-1}}{\delta_k} \right) = f_k^i \text{ a.e. in } \Omega, \\ (p_k^i - p_k^{i-1}, z_k^i - z_k^{i-1}) \in N_{K_\lambda}(\sigma_k^i, \xi_k^i) \text{ a.e. in } \Omega. \end{cases} \quad (3.8)$$

Proof. The first condition is a consequence of the fact that $(u_k^i, e_k^i, p_k^i, z_k^i) \in \mathcal{A}_{\text{vp}}(w_k^i) \times L^2(\Omega)$ for all $i \geq 0$. Let us define $\sigma_k^i := \mathbb{C}e_k^i$ and $\xi_k^i := -z_k^i$. Clearly from (3.1) and (3.5) we have $(\sigma_k^0, \xi_k^0) \in K_\lambda$ and $(\sigma_k^1, \xi_k^1) \in K_\lambda$.

We now consider $i \geq 2$. For any $(v, \eta, q, \zeta) \in \mathcal{A}_{\text{vp}}(0) \times L^2(\Omega)$ and $t > 0$, the quadruplet $(u_k^i, e_k^i, p_k^i, z_k^i) + t(v, \eta, q, \zeta) \in \mathcal{A}_{\text{vp}}(w_k^i) \times L^2(\Omega)$ is admissible for the minimum problem (3.6). Hence choosing it as a competitor, dividing the resulting inequality by $t > 0$, and letting $t \rightarrow 0^+$ yield

$$\begin{aligned} 0 \leq \int_\Omega \sigma_k^i : \eta \, dx + \varepsilon \int_\Omega \frac{Eu_k^i - Eu_k^{i-1}}{\delta_k} : Ev \, dx + \int_\Omega \frac{u_k^i - 2u_k^{i-1} + u_k^{i-2}}{\delta_k^2} \cdot v \, dx + \int_\Omega z_k^i \zeta \, dx \\ + \mathcal{H}_\lambda(p_k^i - p_k^{i-1} + q, z_k^i - z_k^{i-1} + \zeta) - \mathcal{H}_\lambda(p_k^i - p_k^{i-1}, z_k^i - z_k^{i-1}) - \int_\Omega f_k^i \cdot v \, dx, \end{aligned}$$

where we used the convexity of \mathcal{H}_λ . Taking in particular $(v, \eta, q, \zeta) = \pm(\varphi, E\varphi, 0, 0)$ where $\varphi \in C_c^\infty(\Omega; \mathbb{R}^n)$, we infer that

$$\int_\Omega \left(\sigma_k^i + \varepsilon \frac{Eu_k^i - Eu_k^{i-1}}{\delta_k} \right) : E\varphi \, dx + \int_\Omega \frac{u_k^i - 2u_k^{i-1} + u_k^{i-2}}{\delta_k^2} \cdot \varphi \, dx = \int_\Omega f_k^i \cdot \varphi \, dx,$$

leading to

$$\frac{u_k^i - 2u_k^{i-1} + u_k^{i-2}}{\delta_k^2} - \operatorname{div} \left(\sigma_k^i + \varepsilon \frac{Eu_k^i - Eu_k^{i-1}}{\delta_k} \right) = f_k^i \quad \text{a.e. in } \Omega.$$

Next choosing $(v, \eta, q, \zeta) = (0, -\hat{q} + p_k^i - p_k^{i-1}, \hat{q} - p_k^i + p_k^{i-1}, \hat{\zeta} - z_k^i + z_k^{i-1})$ where $\hat{q} \in L^2(\Omega; \mathbb{M}_{sym}^{n \times n})$ and $\hat{\zeta} \in L^2(\Omega)$ implies

$$\mathcal{H}_\lambda(\hat{q}, \hat{\zeta}) \geq \mathcal{H}_\lambda(p_k^i - p_k^{i-1}, z_k^i - z_k^{i-1}) + \int_\Omega \sigma_k^i : (\hat{q} - (p_k^i - p_k^{i-1})) \, dx + \int_\Omega \xi_k^i (\hat{\zeta} - (z_k^i - z_k^{i-1})) \, dx.$$

Localizing this inequality yields

$$\begin{aligned} H_\lambda(\hat{q}, \hat{\zeta}) &\geq H_\lambda(p_k^i(x) - p_k^{i-1}(x), z_k^i(x) - z_k^{i-1}(x)) \\ &\quad + \sigma_k^i(x) : (\hat{q} - (p_k^i(x) - p_k^{i-1}(x))) + \xi_k^i(x) (\hat{\zeta} - (z_k^i(x) - z_k^{i-1}(x))), \end{aligned}$$

for all $(\hat{q}, \hat{\zeta}) \in \mathbb{M}_{sym}^{n \times n} \times \mathbb{R}$ and for a.e. $x \in \Omega$. This implies that $(\sigma_k^i, \xi_k^i) \in K_\lambda$ and $(p_k^i - p_k^{i-1}, z_k^i - z_k^{i-1}) \in N_{K_\lambda}(\sigma_k^i, \xi_k^i)$ a.e. in Ω . \square

3.2. A priori estimates and compactness. Thanks to the Euler-Lagrange equation, we derive *a priori* estimates. Let us define three types of interpolations. We start with piecewise constant left-continuous interpolations defined by

$$\begin{aligned} u_k(0) &:= u_0, & w_k(0) &:= w(0), & f_k(0) &:= f(0), & e_k(0) &:= e_0, \\ \sigma_k(0) &:= \sigma_0, & p_k(0) &:= p_0, & \xi_k(0) &:= \xi_0, & z_k(0) &:= z_0, \end{aligned}$$

and, for all $t \in (t_k^{i-1}, t_k^i]$ and $i \in \{1, \dots, N_k\}$,

$$\begin{aligned} u_k(t) &:= u_k^i, & w_k(t) &:= w_k^i, & f_k(t) &:= f_k^i, & e_k(t) &:= e_k^i, \\ \sigma_k(t) &:= \sigma_k^i, & p_k(t) &:= p_k^i, & \xi_k(t) &:= \xi_k^i, & z_k(t) &:= z_k^i. \end{aligned}$$

We will also consider the piecewise affine interpolations given by

$$\begin{aligned} \hat{u}_k(0) &:= u_0, & \hat{w}_k(0) &:= w(0), & \hat{e}_k(0) &:= e_0, & \hat{\sigma}_k(0) &:= \sigma_0, \\ \hat{p}_k(0) &:= p_0, & \hat{\xi}_k(0) &:= \xi_0, & \hat{z}_k(0) &:= z_0, \end{aligned}$$

and, for every $t \in (t_k^{i-1}, t_k^i]$ and $i \in \{1, \dots, N_k\}$,

$$\begin{aligned} \hat{u}_k(t) &:= u_k^{i-1} + \frac{t - t_k^{i-1}}{\delta_k} (u_k^i - u_k^{i-1}), & \hat{w}_k(t) &:= w_k^{i-1} + \frac{t - t_k^{i-1}}{\delta_k} (w_k^i - w_k^{i-1}), \\ \hat{e}_k(t) &:= e_k^{i-1} + \frac{t - t_k^{i-1}}{\delta_k} (e_k^i - e_k^{i-1}), & \hat{\sigma}_k(t) &:= \sigma_k^{i-1} + \frac{t - t_k^{i-1}}{\delta_k} (\sigma_k^i - \sigma_k^{i-1}), \\ \hat{\xi}_k(t) &:= \xi_k^{i-1} + \frac{t - t_k^{i-1}}{\delta_k} (\xi_k^i - \xi_k^{i-1}), & \hat{z}_k(t) &:= z_k^{i-1} + \frac{t - t_k^{i-1}}{\delta_k} (z_k^i - z_k^{i-1}), \\ \hat{p}_k(t) &:= p_k^{i-1} + \frac{t - t_k^{i-1}}{\delta_k} (p_k^i - p_k^{i-1}). \end{aligned}$$

Finally, let \tilde{u}_k , \tilde{w}_k , and \tilde{w}_k be quadratic interpolations of $\{u_k^i\}_{0 \leq i \leq N_k}$ and $\{w_k^i\}_{0 \leq i \leq N_k}$ satisfying $\tilde{u}_k(t_k^i) = u_k^i$, $\tilde{w}_k(t_k^i) = w_k^i$ for all $i \in \{0, \dots, N_k\}$, and

$$\ddot{\tilde{u}}_k(t) = \frac{u_k^i - 2u_k^{i-1} + u_k^{i-2}}{\delta_k^2}, \quad \ddot{\tilde{w}}_k(t) = \frac{w_k^i - 2w_k^{i-1} + w_k^{i-2}}{\delta_k^2}, \quad \ddot{\tilde{w}}_k(t) = \frac{w_k^{i+1} - 2w_k^i + w_k^{i-1}}{\delta_k^2},$$

for all $t \in (t_k^{i-1}, t_k^i]$, $i \in \{1, \dots, N_k\}$, where we set $u_k^{-1} := u_0$, $w_k^{-1} := w(0)$, and $w_k^{N_k+1} := w(T)$.

Observe that for all $t \in [0, T]$,

$$\begin{cases} Eu_k(t) = e_k(t) + p_k(t) \text{ a.e. in } \Omega, & u_k(t) = w_k(t) \text{ } \mathcal{H}^{n-1}\text{-a.e. on } \partial\Omega, \\ E\hat{u}_k(t) = \hat{e}_k(t) + \hat{p}_k(t) \text{ a.e. in } \Omega, & \hat{u}_k(t) = \hat{w}_k(t) \text{ } \mathcal{H}^{n-1}\text{-a.e. on } \partial\Omega, \\ \sigma_k(t) = \mathbb{C}e_k(t), \xi_k(t) = -z_k(t), & (\sigma_k(t), \xi_k(t)) \in K_\lambda \text{ a.e. in } \Omega, \\ \hat{\sigma}_k(t) = \mathbb{C}\hat{e}_k(t), \hat{\xi}_k(t) = -\hat{z}_k(t), & (\hat{\sigma}_k(t), \hat{\xi}_k(t)) \in K_\lambda \text{ a.e. in } \Omega, \end{cases} \quad (3.9)$$

and for a.e. $t \in [\bar{\delta}_k, T]$,

$$\begin{cases} \ddot{\bar{u}}_k(t) - \operatorname{div}(\sigma_k(t) + \varepsilon E\dot{\bar{u}}_k(t)) = f_k(t) \text{ a.e. in } \Omega, \\ (\dot{\bar{p}}_k(t), \dot{\bar{z}}_k(t)) \in N_{K_\lambda}(\sigma_k(t), \xi_k(t)) \text{ a.e. in } \Omega. \end{cases} \quad (3.10)$$

Moreover (3.7) ensures that for all $0 \leq s \leq t \leq T$,

$$z_k(s) \leq z_k(t), \quad \hat{z}_k(s) \leq \hat{z}_k(t), \quad \text{and} \quad \hat{z}_k(t) \geq 0 \quad \text{a.e. in } \Omega. \quad (3.11)$$

We now derive some *a priori* estimates.

Proposition 3.4. *There exists a constant $C > 0$ (independent of k , ε , and λ) such that*

$$\begin{aligned} & \|z_k\|_{L^\infty(0,T;L^2(\Omega))} + \|e_k\|_{L^\infty(0,T;L^2(\Omega;M_{sym}^{n \times n}))} + \|\hat{u}_k\|_{L^\infty(0,T;L^2(\Omega;\mathbb{R}^n))} + \sqrt{\varepsilon} \|E\dot{\bar{u}}_k\|_{L^2(0,T;L^2(\Omega;M_{sym}^{n \times n}))} \\ & \leq C (\|v_0\|_2 + \|Ev_0\|_2 + \|e_0\|_2 + \|z_0\|_2 + \|\ddot{w}\|_{L^1(0,T;H^1(\Omega;\mathbb{R}^n))} + \|f\|_{L^1(0,T;L^2(\Omega;\mathbb{R}^n))}). \end{aligned} \quad (3.12)$$

Moreover, there exists a constant $C_\varepsilon > 0$ (independent of k) such that

$$\begin{aligned} & \|\hat{u}_k\|_{L^\infty(0,T;H^1(\Omega;\mathbb{R}^n))} + \|\hat{z}_k\|_{L^2(0,T;L^2(\Omega))} \\ & \quad + \|\hat{e}_k\|_{L^2(0,T;L^2(\Omega;M_{sym}^{n \times n}))} + \|\hat{p}_k\|_{L^2(0,T;L^2(\Omega;M_{sym}^{n \times n}))} \leq C_\varepsilon. \end{aligned} \quad (3.13)$$

Proof. Let $i \geq 2$, let us multiply the first line of (3.8) by $u_k^i - u_k^{i-1}$ and integrate by parts over Ω . Using the kinematic compatibility $Eu_k^i - Eu_k^{i-1} = (e_k^i - e_k^{i-1}) + (p_k^i - p_k^{i-1})$ yields

$$\begin{aligned} & \frac{1}{2} \left\| \frac{u_k^i - u_k^{i-1}}{\delta_k} \right\|_2^2 - \frac{1}{2} \left\| \frac{u_k^{i-1} - u_k^{i-2}}{\delta_k} \right\|_2^2 + \frac{1}{2} \left\| \frac{u_k^i - 2u_k^{i-1} + u_k^{i-2}}{\delta_k} \right\|_2^2 + \int_\Omega \sigma_k^i : (p_k^i - p_k^{i-1}) \, dx \\ & \quad + \mathcal{Q}(e_k^i) - \mathcal{Q}(e_k^{i-1}) + \mathcal{Q}(e_k^i - e_k^{i-1}) + \varepsilon \delta_k \left\| \frac{Eu_k^i - Eu_k^{i-1}}{\delta_k} \right\|_2^2 \\ & = \int_\Omega \left(\sigma_k^i + \varepsilon \frac{Eu_k^i - Eu_k^{i-1}}{\delta_k} \right) : (Eu_k^i - Eu_k^{i-1}) \, dx + \int_\Omega \frac{u_k^i - 2u_k^{i-1} + u_k^{i-2}}{\delta_k^2} \cdot (w_k^i - w_k^{i-1}) \, dx \\ & \quad + \int_\Omega f_k^i \cdot (u_k^i - u_k^{i-1}) \, dx - \int_\Omega f_k^i \cdot (w_k^i - w_k^{i-1}) \, dx. \end{aligned}$$

But according to the discrete flow rule in (3.8) and convex duality, we have $(\sigma_k^i, \xi_k^i) \in \partial H_\lambda(p_k^i - p_k^{i-1}, z_k^i - z_k^{i-1})$ a.e. in Ω , and thus

$$\begin{aligned} & \int_\Omega \sigma_k^i : (p_k^i - p_k^{i-1}) \, dx = \mathcal{H}_\lambda(p_k^i - p_k^{i-1}, z_k^i - z_k^{i-1}) - \int_\Omega \xi_k^i (z_k^i - z_k^{i-1}) \, dx \\ & = \mathcal{H}_\lambda(p_k^i - p_k^{i-1}, z_k^i - z_k^{i-1}) + \frac{1}{2} \|z_k^i\|_2^2 - \frac{1}{2} \|z_k^{i-1}\|_2^2 + \frac{1}{2} \|z_k^i - z_k^{i-1}\|_2^2, \end{aligned}$$

where we used the fact that $\xi_k^i = -z_k^i$. Hence

$$\begin{aligned}
 & \frac{1}{2} \left\| \frac{u_k^i - u_k^{i-1}}{\delta_k} \right\|_2^2 - \frac{1}{2} \left\| \frac{u_k^{i-1} - u_k^{i-2}}{\delta_k} \right\|_2^2 + \mathcal{H}_\lambda(p_k^i - p_k^{i-1}, z_k^i - z_k^{i-1}) \\
 & \quad + \frac{1}{2} \|z_k^i\|_2^2 - \frac{1}{2} \|z_k^{i-1}\|_2^2 + \mathcal{Q}(e_k^i) - \mathcal{Q}(e_k^{i-1}) + \varepsilon \delta_k \left\| \frac{Eu_k^i - Eu_k^{i-1}}{\delta_k} \right\|_2^2 \\
 & \leq \int_\Omega \left(\sigma_k^i + \varepsilon \frac{Eu_k^i - Eu_k^{i-1}}{\delta_k} \right) : (Ew_k^i - Ew_k^{i-1}) dx + \int_\Omega \frac{u_k^i - 2u_k^{i-1} + u_k^{i-2}}{\delta_k^2} \cdot (w_k^i - w_k^{i-1}) dx \\
 & \quad + \int_\Omega f_k^i \cdot (u_k^i - u_k^{i-1}) dx - \int_\Omega f_k^i \cdot (w_k^i - w_k^{i-1}) dx.
 \end{aligned}$$

Summing up for $i = 2$ to j , and using a discrete integration by parts leads to

$$\begin{aligned}
 & \frac{1}{2} \left\| \frac{u_k^j - u_k^{j-1}}{\delta_k} \right\|_2^2 - \frac{1}{2} \|v_0\|_2^2 + \sum_{i=2}^j \mathcal{H}_\lambda(p_k^i - p_k^{i-1}, z_k^i - z_k^{i-1}) \\
 & \quad + \frac{1}{2} \|z_k^j\|_2^2 - \frac{1}{2} \|z_0\|_2^2 + \mathcal{Q}(e_k^j) - \mathcal{Q}(e_0) + \varepsilon \sum_{i=2}^j \delta_k \left\| \frac{Eu_k^i - Eu_k^{i-1}}{\delta_k} \right\|_2^2 \\
 & \leq \sum_{i=2}^j \int_\Omega \left(\sigma_k^i + \varepsilon \frac{Eu_k^i - Eu_k^{i-1}}{\delta_k} \right) : (Ew_k^i - Ew_k^{i-1}) dx \\
 & \quad - \sum_{i=2}^{j-1} \int_\Omega (u_k^i - u_k^{i-1}) \cdot \frac{w_k^{i+1} - 2w_k^i + w_k^{i-1}}{\delta_k^2} dx - \int_\Omega v_0 \cdot \frac{w_k^2 - w_k^1}{\delta_k} dx \\
 & \quad + \int_\Omega \frac{u_k^j - u_k^{j-1}}{\delta_k} \cdot \frac{w_k^j - w_k^{j-1}}{\delta_k} dx + \sum_{i=2}^j \int_\Omega f_k^i \cdot (u_k^i - u_k^{i-1}) dx - \sum_{i=2}^j \int_\Omega f_k^i \cdot (w_k^i - w_k^{i-1}) dx.
 \end{aligned}$$

Consequently, if $t \in (t_k^{j-1}, t_k^j]$, we get from the positive 1-homogeneity of H_λ and Hölder's inequality

$$\begin{aligned}
 & \frac{1}{2} \|\hat{u}_k(t)\|_2^2 - \frac{1}{2} \|v_0\|_2^2 + \int_{\delta_k}^{t_k^j} \mathcal{H}_\lambda(\dot{p}_k(s), \dot{z}_k(s)) ds \\
 & \quad + \frac{1}{2} \|z_k(t)\|_2^2 - \frac{1}{2} \|z_0\|_2^2 + \mathcal{Q}(e_k(t)) - \mathcal{Q}(e_0) + \varepsilon \int_{\delta_k}^{t_k^j} \|E\hat{u}_k(s)\|_2^2 ds \\
 & \leq \left(\sqrt{T} \|\sigma_k\|_{L^\infty(0,T;L^2(\Omega; \mathbb{M}_{sym}^n \times \mathbb{M}_{sym}^n))} + \varepsilon \|E\hat{u}_k\|_{L^2(0,T;L^2(\Omega; \mathbb{M}_{sym}^n \times \mathbb{M}_{sym}^n))} \right) \|E\hat{w}_k\|_{L^2(0,T;L^2(\Omega; \mathbb{M}_{sym}^n \times \mathbb{M}_{sym}^n))} \\
 & \quad + \left(\|\ddot{w}_k\|_{L^1(0,T;L^2(\Omega; \mathbb{R}^n))} + \|\dot{w}_k\|_{L^\infty(0,T;L^2(\Omega; \mathbb{R}^n))} \right) \|\hat{u}_k\|_{L^\infty(0,T;L^2(\Omega; \mathbb{R}^n))} + \|v_0\|_2 \|\dot{w}_k\|_{L^\infty(0,T;L^2(\Omega; \mathbb{R}^n))} \\
 & \quad + \left(\|\hat{u}_k\|_{L^\infty(0,T;L^2(\Omega; \mathbb{R}^n))} + \|\hat{w}_k\|_{L^\infty(0,T;L^2(\Omega; \mathbb{R}^n))} \right) \|f_k\|_{L^1(0,T;L^2(\Omega; \mathbb{R}^n))},
 \end{aligned}$$

from which (3.12) follows.

On the other hand, since

$$\hat{u}_k(t) = u_0 + \int_0^t \hat{u}_k(s) ds \quad \text{and} \quad E\hat{u}_k(t) = Eu_0 + \int_0^t E\hat{u}_k(s) ds,$$

we deduce that

$$\|\hat{u}_k\|_{L^\infty(0,T;H^1(\Omega; \mathbb{R}^n))} \leq C_\varepsilon,$$

for some constant $C_\varepsilon > 0$ depending on ε . Moreover, according to the kinematic compatibility (3.9), we have $E\dot{u}_k(t) = \dot{e}_k(t) + \dot{p}_k(t)$ for all $t \in [0, T]$. Multiplying this equality by $\dot{\sigma}_k(t)$ and integrating over Ω yields, from the coercivity condition (2.1),

$$\alpha_{\mathbb{C}} \|\dot{e}_k(t)\|_2^2 \leq \int_{\Omega} \dot{\sigma}_k(t) : \dot{e}_k(t) dx = \int_{\Omega} \dot{\sigma}_k(t) : E\dot{u}_k(t) dx - \int_{\Omega} \dot{\sigma}_k(t) : \dot{p}_k(t) dx.$$

By the discrete flow rule in (3.8), for all $i \geq 2$ we have

$$\begin{aligned} \int_{\Omega} \sigma_k^i : (p_k^i - p_k^{i-1}) dx + \int_{\Omega} \xi_k^i (z_k^i - z_k^{i-1}) dx &= \mathcal{H}_\lambda(p_k^i - p_k^{i-1}, z_k^i - z_k^{i-1}) \\ &\geq \int_{\Omega} \sigma_k^{i-1} : (p_k^i - p_k^{i-1}) dx + \int_{\Omega} \xi_k^{i-1} (z_k^i - z_k^{i-1}) dx, \end{aligned}$$

since $(\sigma_k^{i-1}, \xi_k^{i-1}) \in K_\lambda$. Hence, if $t \in (t_k^{i-1}, t_k^i]$, we have that

$$\begin{aligned} \int_{\Omega} \dot{\sigma}_k(t) : \dot{p}_k(t) dx &= \frac{1}{\delta_k^2} \int_{\Omega} (\sigma_k^i - \sigma_k^{i-1}) : (p_k^i - p_k^{i-1}) dx \\ &\geq -\frac{1}{\delta_k^2} \int_{\Omega} (\xi_k^i - \xi_k^{i-1}) (z_k^i - z_k^{i-1}) dx = \left\| \frac{z_k^i - z_k^{i-1}}{\delta_k} \right\|_2^2 = \|\dot{z}_k(t)\|_2^2. \end{aligned}$$

Consequently, we derive that for all $t \in [\delta_k, T]$,

$$\|\dot{z}_k(t)\|_2^2 + \alpha_{\mathbb{C}} \|\dot{e}_k(t)\|_2^2 \leq \beta_{\mathbb{C}} \|\dot{e}_k(t)\|_2 \|E\dot{u}_k(t)\|_2$$

and by (3.12), we deduce that

$$\|\dot{z}_k\|_{L^2(\delta_k, T; L^2(\Omega))} + \|\dot{e}_k\|_{L^2(\delta_k, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))} \leq C_\varepsilon$$

for some constant $C_\varepsilon > 0$ independent of k . Moreover, by the relation $\dot{p}_k = E\dot{u}_k - \dot{e}_k$, we also have that

$$\|\dot{p}_k\|_{L^2(\delta_k, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))} \leq C_\varepsilon.$$

Finally since by (3.5) we have $\dot{e}_k(t) = 0$, $\dot{p}_k(t) = E v_0$, and $\dot{z}_k(t) = 0$ for all $t \in [0, \delta_k]$, the proof of the proposition is complete. \square

From the previous *a priori* estimates, we now deduce compactness results. Indeed, as a consequence of (3.12), (3.13), and Korn's inequality, we can extract a subsequence (not relabeled), and find

$$u \in H^1([0, T]; H^1(\Omega; \mathbb{R}^n)) \cap W^{1, \infty}([0, T]; L^2(\Omega; \mathbb{R}^n))$$

such that

$$\begin{cases} \hat{u}_k \rightharpoonup u \text{ weakly in } H^1([0, T]; H^1(\Omega; \mathbb{R}^n)), \\ \hat{u}_k \rightharpoonup^* u \text{ weakly* in } L^\infty([0, T]; L^2(\Omega; \mathbb{R}^n)). \end{cases}$$

Moreover, since for all $t \in [0, T]$,

$$\|\hat{u}_k(t) - u_k(t)\|_{H^1(\Omega; \mathbb{R}^n)} \leq 2\delta_k \|\hat{u}_k(t)\|_{H^1(\Omega; \mathbb{R}^n)},$$

we infer that

$$u_k \rightharpoonup u \text{ weakly in } L^2(0, T; H^1(\Omega; \mathbb{R}^n)).$$

Note that the previous weak convergences of the sequence (\hat{u}_k) implies, by Ascoli-Arzelà Theorem, that $\hat{u}_k(t) \rightharpoonup u(t)$ weakly in $H^1(\Omega; \mathbb{R}^n)$ for all $t \in [0, T]$. But since $\hat{u}_k(t) = \hat{w}_k(t)$ \mathcal{H}^{n-1} -a.e. on $\partial\Omega$, and $\hat{w}_k(t) \rightarrow w(t)$ strongly in $H^1(\Omega; \mathbb{R}^n)$ (by the absolute continuity of $t \mapsto w(t)$ in $H^1(\Omega; \mathbb{R}^n)$), we infer by the continuity of the trace that $u(t) = w(t)$ \mathcal{H}^{n-1} -a.e. on $\partial\Omega$. Moreover, since $\hat{u}_k(0) = u_0$, we deduce that $u(0) = u_0$.

Using again (3.13), we get that

$$\begin{cases} \hat{e}_k \rightharpoonup e \text{ weakly in } H^1([0, T]; L^2(\Omega; \mathbb{M}_{sym}^{n \times n})), \\ \hat{p}_k \rightharpoonup p \text{ weakly in } H^1([0, T]; L^2(\Omega; \mathbb{M}_{sym}^{n \times n})), \\ \hat{z}_k \rightharpoonup z \text{ weakly in } H^1([0, T]; L^2(\Omega)), \end{cases} \quad (3.14)$$

for some $e, p \in H^1([0, T]; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))$, and some $z \in H^1([0, T]; L^2(\Omega))$. Applying again Ascoli-Arzelà Theorem, we obtain that for all $t \in [0, T]$,

$$\begin{cases} \hat{e}_k(t) \rightharpoonup e(t) \text{ weakly in } L^2(\Omega; \mathbb{M}_{sym}^{n \times n}), \\ \hat{p}_k(t) \rightharpoonup p(t) \text{ weakly in } L^2(\Omega; \mathbb{M}_{sym}^{n \times n}), \\ \hat{z}_k(t) \rightharpoonup z(t) \text{ weakly in } L^2(\Omega). \end{cases}$$

In particular, since $(\hat{e}_k(0), \hat{p}_k(0), \hat{z}_k(0)) = (e_0, p_0, z_0)$, we deduce that

$$(e(0), p(0), z(0)) = (e_0, p_0, z_0).$$

Moreover, for all $t \in [0, T]$, we have

$$(u(t), e(t), p(t)) \in \mathcal{A}_{vp}(w(t)),$$

and by (3.11), we get that for all $0 \leq s \leq t \leq T$,

$$z(s) \leq z(t) \quad \text{a.e. in } \Omega.$$

Then, for a.e. $t \in [0, T]$, we have

$$(\dot{u}(t), \dot{e}(t), \dot{p}(t)) \in \mathcal{A}_{vp}(\dot{w}(t)), \quad \dot{z}(t) \geq 0. \quad (3.15)$$

On the other hand, since for all $t \in [0, T]$,

$$\begin{cases} \|\hat{e}_k(t) - e_k(t)\|_{L^2(\Omega; \mathbb{M}_{sym}^{n \times n})} \leq 2\delta_k \|\dot{\hat{e}}_k(t)\|_{L^2(\Omega; \mathbb{M}_{sym}^{n \times n})}, \\ \|\hat{p}_k(t) - p_k(t)\|_{L^2(\Omega; \mathbb{M}_{sym}^{n \times n})} \leq 2\delta_k \|\dot{\hat{p}}_k(t)\|_{L^2(\Omega; \mathbb{M}_{sym}^{n \times n})}, \\ \|\hat{z}_k(t) - z_k(t)\|_{L^2(\Omega)} \leq 2\delta_k \|\dot{\hat{z}}_k(t)\|_{L^2(\Omega)}, \end{cases} \quad (3.16)$$

we also have that

$$\begin{cases} e_k \rightharpoonup e \text{ weakly* in } L^\infty(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n})), \\ p_k \rightharpoonup p \text{ weakly in } L^2(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n})), \\ z_k \rightharpoonup z \text{ weakly* in } L^\infty(0, T; L^2(\Omega)). \end{cases}$$

Finally since, for all $t \in [0, T]$, $(\hat{\sigma}_k(t), \hat{\xi}_k(t)) = (\mathbb{C}\hat{e}_k(t), -\hat{z}_k(t)) \rightharpoonup (\mathbb{C}e(t), -z(t)) =: (\sigma(t), \xi(t))$ weakly in $L^2(\Omega; \mathbb{M}_{sym}^{n \times n}) \times L^2(\Omega)$, and, from (3.9), $(\hat{\sigma}_k(t), \hat{\xi}_k(t)) \in K_\lambda$ a.e. in Ω , where K_λ is a closed and convex set, we infer that for all $t \in [0, T]$

$$(\sigma(t), \xi(t)) \in K_\lambda \quad \text{a.e. in } \Omega. \quad (3.17)$$

3.3. Weak formulation of the equation of motion. At this step, we do not have enough time regularity on the velocity \dot{u} to write the initial condition $\dot{u}(0) = v_0$. As usual in hyperbolic equations, this condition will be expressed by giving a weak formulation of the equation of motion.

Proposition 3.5. *For every $\varphi \in C_c^\infty(\Omega \times [0, T]; \mathbb{R}^n)$,*

$$-\int_0^T \int_\Omega \dot{u} \cdot \dot{\varphi} \, dx \, dt + \int_0^T \int_\Omega (\sigma + \varepsilon E\dot{u}) : E\varphi \, dx \, dt = \int_0^T \int_\Omega f \cdot \varphi \, dx \, dt + \int_\Omega v_0 \varphi(0) \, dx.$$

Proof. Let $\varphi \in \mathcal{C}_c^\infty(\Omega \times [0, T]; \mathbb{R}^n)$ and define the right-continuous piecewise constant and piecewise affine interpolations by

$$\begin{cases} \varphi_k(t) := \varphi(t_k^{i-1}), \\ \hat{\varphi}_k(t) := \varphi(t_k^{i-1}) + \frac{t - t_k^i}{\delta_k} (\varphi(t_k^i) - \varphi(t_k^{i-1})), \end{cases} \quad \text{for all } t \in [t_k^{i-1}, t_k^i], \quad i \in \{1, \dots, N_k\}.$$

Let us multiply the first equation of (3.8) by $\varphi(t_k^{i-1})$ and integrate by parts over Ω . Since, for k large enough, we have

$$\begin{aligned} \sum_{i=2}^{N_k} \delta_k \int_{\Omega} \frac{u_k^i - 2u_k^{i-1} + u_k^{i-2}}{\delta_k^2} \cdot \varphi(t_k^{i-1}) \, dx \\ = - \sum_{i=1}^{N_k} \delta_k \int_{\Omega} \frac{u_k^i - u_k^{i-1}}{\delta_k} \cdot \frac{\varphi(t_k^i) - \varphi(t_k^{i-1})}{\delta_k} \, dx - \int_{\Omega} v_0 \varphi(0) \, dx, \end{aligned}$$

we deduce that

$$\begin{aligned} - \sum_{i=1}^{N_k} \delta_k \int_{\Omega} \frac{u_k^i - u_k^{i-1}}{\delta_k} \cdot \frac{\varphi(t_k^i) - \varphi(t_k^{i-1})}{\delta_k} \, dx + \sum_{i=2}^{N_k} \delta_k \int_{\Omega} \left(\sigma_k^i + \varepsilon \left(\frac{Eu_k^i - Eu_k^{i-1}}{\delta_k} \right) \right) : E\varphi(t_k^{i-1}) \, dx \\ = \sum_{i=2}^{N_k} \delta_k \int_{\Omega} f_k^i \cdot \varphi(t_k^{i-1}) \, dx + \int_{\Omega} v_0 \varphi(0) \, dx, \end{aligned}$$

hence,

$$- \int_0^T \int_{\Omega} \dot{u}_k \cdot \dot{\varphi}_k \, dx \, dt + \int_0^T \int_{\Omega} (\sigma_k + \varepsilon E \dot{u}_k) : E\varphi_k \, dx \, dt = \int_0^T \int_{\Omega} f_k \cdot \varphi_k \, dx \, dt + \int_{\Omega} v_0 \varphi(0) \, dx.$$

Note that $\varphi_k \rightarrow \varphi$ strongly in $L^2(0, T; H^1(\Omega; \mathbb{R}^n))$, and $\dot{\varphi}_k \rightarrow \dot{\varphi}$ strongly in $L^2(0, T; L^2(\Omega; \mathbb{R}^n))$. Thus since, by the absolute continuity of $t \mapsto f(t)$ in $L^2(\Omega; \mathbb{R}^n)$ we have $f_k \rightarrow f$ strongly in $L^2(0, T; L^2(\Omega; \mathbb{R}^n))$, we get the desired result by passing to the limit as $k \rightarrow \infty$ in the previous expression, and by using the weak convergences established for the sequences (\hat{u}_k) and (σ_k) . \square

Remark 3.6. As a consequence of Proposition 3.5, we have $\ddot{u} \in L^2(0, T; H^{-1}(\Omega; \mathbb{R}^n))$. Hence since $\dot{u} - \dot{w} \in L^2(0, T; H_0^1(\Omega; \mathbb{R}^n))$ and $\ddot{u} - \ddot{w} \in L^2(0, T; H^{-1}(\Omega; \mathbb{R}^n))$, we deduce from [15, Section 5.9, Theorem 3] that $\dot{u} \in \mathcal{C}([0, T]; L^2(\Omega; \mathbb{R}^n))$. Therefore,

$$\ddot{u} - \operatorname{div}(\sigma + \varepsilon E \dot{u}) = f \quad \text{in } L^2(0, T; H^{-1}(\Omega; \mathbb{R}^n)),$$

and the initial condition is expressed in the standard way

$$\dot{u}(0) = v_0. \tag{3.18}$$

3.4. Flow rule. We now derive the flow rule. To this end we need to improve the weak convergences established so far for the elastic strain and the hardening variable, into strong convergences.

Lemma 3.7. *The following strong convergences hold:*

$$\begin{cases} e_k, \hat{e}_k \rightarrow e \text{ strongly in } L^2(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n})), \\ z_k, \hat{z}_k \rightarrow z \text{ strongly in } L^2(0, T; L^2(\Omega)). \end{cases}$$

Proof. For every $t \in (0, T]$ and for every k , we set $[t]_k := t_k^i$ where $i \in \{1, \dots, N_k\}$ is such that $t \in (t_k^{i-1}, t_k^i]$. According to Remark 3.6, we have for all $\varphi \in L^2(0, T; H_0^1(\Omega; \mathbb{R}^n))$,

$$\int_0^T \langle \dot{u}(s), \varphi(s) \rangle \, ds + \int_0^T \int_{\Omega} (\sigma + \varepsilon E \dot{u}) : E\varphi \, dx \, ds = \int_0^T \int_{\Omega} f \cdot \varphi \, dx \, ds,$$

where the brackets $\langle \cdot, \cdot \rangle$ denote the duality pairing between $H_0^1(\Omega; \mathbb{R}^n)$ and $H^{-1}(\Omega; \mathbb{R}^n)$. Hence, taking $\varphi = (\hat{u}_k - \hat{w}_k)\chi_{[\delta_k, [t]_k]}$ as test function in the above relation and in the first equation of (3.10), and subtracting the resulting expressions yield

$$\begin{aligned} \int_{\delta_k}^{[t]_k} \langle \ddot{\hat{u}}_k(s) - \ddot{u}(s), \hat{u}_k(s) - \hat{w}_k(s) \rangle ds + \int_{\delta_k}^{[t]_k} \int_{\Omega} ((\sigma_k + \varepsilon E \hat{u}_k) - (\sigma + \varepsilon E \dot{u})) : (E \hat{u}_k - E \hat{w}_k) dx ds \\ = \int_{\delta_k}^{[t]_k} \int_{\Omega} (f_k - f) \cdot (\hat{u}_k - \hat{w}_k) dx ds. \end{aligned}$$

According to (3.12), the sequence (\hat{u}_k) is uniformly bounded in $L^\infty(0, T; L^2(\Omega; \mathbb{R}^n))$. Hence, since $f_k \rightarrow f$ strongly in $L^1(0, T; L^2(\Omega; \mathbb{R}^n))$, the integral in the right-handside of the previous equality tends to zero as $k \rightarrow \infty$. Moreover, the initial condition (3.5) and Proposition 3.4 ensure that

$$\begin{aligned} \lim_{k \rightarrow \infty} \left(\int_{\delta_k}^{[t]_k} \langle \ddot{\hat{u}}_k(s) - \ddot{u}(s), \hat{u}_k(s) - \hat{w}_k(s) \rangle ds \right. \\ \left. + \int_0^{[t]_k} \int_{\Omega} ((\sigma_k + \varepsilon E \hat{u}_k) - (\sigma + \varepsilon E \dot{u})) : (E \hat{u}_k - E \hat{w}_k) dx ds \right) = 0. \end{aligned}$$

According to (3.10) and (3.12), the sequence $(\|\ddot{\hat{u}}_k\|_{L^2(\delta_k, T; H^{-1}(\Omega; \mathbb{R}^n))})_{k \in \mathbb{N}}$ is uniformly bounded with respect to k . Thus since $\sigma_k \rightharpoonup \sigma$ and $E \hat{u}_k \rightharpoonup E \dot{u}$ weakly in $L^2(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))$, and $E \hat{w}_k \rightarrow E \dot{w}$ strongly in $L^2(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))$, we infer by dominated convergence that

$$\begin{aligned} \lim_{k \rightarrow \infty} \int_0^T \left(\int_{\delta_k}^{[t]_k} \langle \ddot{\hat{u}}_k(s) - \ddot{u}(s), \hat{u}_k(s) - \hat{w}_k(s) \rangle ds + \int_0^{[t]_k} \int_{\Omega} (\sigma_k - \sigma) : E \hat{u}_k dx ds \right. \\ \left. + \varepsilon \int_0^{[t]_k} \int_{\Omega} |E \hat{u}_k - E \dot{u}|^2 dx ds \right) dt = 0. \quad (3.19) \end{aligned}$$

We now estimate the first two integrals of (3.19). Let us start with

$$I_k^1 := \int_0^T \int_0^{[t]_k} \int_{\Omega} (\sigma_k - \sigma) : E \hat{u}_k dx ds dt.$$

Indeed, since $E \hat{u}_k \rightharpoonup E \dot{u}$ weakly in $L^2(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))$, then

$$\limsup_{k \rightarrow \infty} I_k^1 = \limsup_{k \rightarrow \infty} \int_0^T \int_0^{[t]_k} \int_{\Omega} \sigma_k : (E \hat{u}_k - E \dot{u}) dx ds dt,$$

and, by kinematic compatibility, $E \hat{u}_k - E \dot{u} = (\hat{e}_k - \dot{e}) + (\hat{p}_k - \dot{p})$, we infer that

$$\limsup_{k \rightarrow \infty} I_k^1 = \limsup_{k \rightarrow \infty} \left(\int_0^T \int_0^{[t]_k} \int_{\Omega} \sigma_k : (\hat{e}_k - \dot{e}) dx ds dt + \int_0^T \int_0^{[t]_k} \int_{\Omega} \sigma_k : (\hat{p}_k - \dot{p}) dx ds dt \right).$$

Using the fact that $\dot{\hat{e}}_k \rightharpoonup \dot{e}$ and $\sigma_k \rightharpoonup \sigma$ weakly in $L^2(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))$, and that $\sigma_k - \hat{\sigma}_k \rightarrow 0$ strongly in $L^2(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))$, we obtain thanks to the bounds (3.13)

$$\begin{aligned} & \limsup_{k \rightarrow \infty} I_k^1 \\ &= \limsup_{k \rightarrow \infty} \left(\int_0^T \int_0^{[t]_k} \int_{\Omega} (\hat{\sigma}_k - \sigma) : (\dot{\hat{e}}_k - \dot{e}) \, dx \, ds \, dt + \int_0^T \int_0^{[t]_k} \int_{\Omega} (\sigma_k : \dot{\hat{p}}_k - \sigma : \dot{p}) \, dx \, ds \, dt \right) \\ &= \limsup_{k \rightarrow \infty} \left(\int_0^T \mathcal{Q}(\hat{e}_k([t]_k) - e([t]_k)) \, dt + \int_0^T \int_0^{[t]_k} \int_{\Omega} (\sigma_k : \dot{\hat{p}}_k - \sigma : \dot{p}) \, dx \, ds \, dt \right), \end{aligned}$$

because $\hat{e}_k(0) - e(0) = 0$. But since, by (3.10), $(\dot{\hat{p}}_k, \dot{\hat{z}}_k) \in N_{K_\lambda}(\sigma_k, \xi_k)$ a.e. in $\Omega \times (\delta_k, T)$, we deduce that

$$\begin{aligned} \int_0^T \int_{\delta_k}^{[t]_k} \int_{\Omega} (\sigma_k : \dot{\hat{p}}_k + \xi_k \dot{\hat{z}}_k) \, dx \, ds \, dt &= \int_0^T \int_{\delta_k}^{[t]_k} \mathcal{H}_\lambda(\dot{\hat{p}}_k(s), \dot{\hat{z}}_k(s)) \, ds \, dt \\ &\geq \int_0^T \int_{\delta_k}^{[t]_k} \int_{\Omega} (\sigma : \dot{\hat{p}}_k + \xi \dot{\hat{z}}_k) \, dx \, ds \, dt, \end{aligned}$$

where we used that $(\sigma, \xi) \in K_\lambda$ a.e. in $\Omega \times (0, T)$ by (3.17). Hence, recalling that $\xi = -z$ and $\xi_k = -z_k$, and using the initial condition (3.5), leads to

$$\begin{aligned} & \limsup_{k \rightarrow \infty} I_k^1 \\ &\geq \limsup_{k \rightarrow \infty} \left(\int_0^T \mathcal{Q}(\hat{e}_k([t]_k) - e([t]_k)) \, dt + \int_0^T \int_0^{[t]_k} \int_{\Omega} (\sigma : (\dot{\hat{p}}_k - \dot{p}) + \dot{\hat{z}}_k(z_k - z)) \, dx \, ds \, dt \right) \\ &= \limsup_{k \rightarrow \infty} \left(\int_0^T \mathcal{Q}(\hat{e}_k([t]_k) - e([t]_k)) \, dt + \int_0^T \int_0^{[t]_k} \int_{\Omega} \dot{\hat{z}}_k(z_k - z) \, dx \, ds \, dt \right), \end{aligned}$$

where we used the fact that $\dot{\hat{p}}_k \rightharpoonup \dot{p}$ weakly in $L^2(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))$. Now since $z_k \rightharpoonup z$ weakly in $L^2(0, T; L^2(\Omega))$, and $z_k - \hat{z}_k \rightarrow 0$ strongly in $L^2(0, T; L^2(\Omega))$, we obtain using the bounds (3.13) that

$$\begin{aligned} \limsup_{k \rightarrow \infty} I_k^1 &\geq \limsup_{k \rightarrow \infty} \left(\int_0^T \mathcal{Q}(\hat{e}_k([t]_k) - e([t]_k)) \, dt + \int_0^T \int_0^{[t]_k} \int_{\Omega} (\dot{\hat{z}}_k - \dot{z})(\hat{z}_k - z) \, dx \, ds \, dt \right) \\ &= \limsup_{k \rightarrow \infty} \left(\int_0^T \mathcal{Q}(\hat{e}_k([t]_k) - e([t]_k)) \, dt + \frac{1}{2} \int_0^T \|\hat{z}_k([t]_k) - z([t]_k)\|_2^2 \, dt \right), \end{aligned}$$

where we used the fact that $\hat{z}_k(0) - z(0) = 0$. Since $\hat{e}_k([t]_k) = e_k(t)$, $\hat{z}_k([t]_k) = z_k(t)$ and $e \in H^1([0, T]; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))$, $z \in H^1([0, T]; L^2(\Omega))$, we conclude that

$$\limsup_{k \rightarrow \infty} I_k^1 \geq \limsup_{k \rightarrow \infty} \left(\int_0^T \mathcal{Q}(e_k(t) - e(t)) \, dt + \frac{1}{2} \int_0^T \|z_k(t) - z(t)\|_2^2 \, dt \right). \quad (3.20)$$

We now estimate

$$I_k^2 := \int_0^T \int_{\delta_k}^{[t]_k} \langle \ddot{u}_k(s) - \ddot{u}(s), \dot{u}_k(s) - \dot{u}(s) \rangle \, ds \, dt.$$

Let us further split the previous integral as

$$I_k^2 = \int_0^T \int_{\delta_k}^{[t]_k} \langle \ddot{w}_k(s) - \ddot{u}(s), \dot{u}_k(s) - \dot{u}(s) \rangle \, ds \, dt + \int_0^T \int_{\delta_k}^{[t]_k} \langle \ddot{u}_k(s) - \ddot{w}_k(s), \dot{u}_k(s) - \dot{u}(s) \rangle \, ds \, dt.$$

Since $\ddot{w}_k \chi_{[\delta_k, [t]_k]} \rightarrow \ddot{w} \chi_{[0, t]}$ strongly in $L^2(0, T; L^2(\Omega; \mathbb{R}^n))$, while $\dot{u}_k - \dot{w}_k \rightharpoonup \dot{u} - \dot{w}$ weakly in $L^2(0, T; H_0^1(\Omega; \mathbb{R}^n))$, we get that

$$\begin{aligned} \lim_{k \rightarrow \infty} \int_0^T \int_{\delta_k}^{[t]_k} \langle \ddot{w}_k(s) - \ddot{u}(s), \dot{u}_k(s) - \dot{w}_k(s) \rangle ds dt &= \int_0^T \int_0^t \langle \ddot{w}(s) - \ddot{u}(s), \dot{u}(s) - \dot{w}(s) \rangle ds dt \\ &= \int_0^T \left(\frac{\|\dot{u}(0) - \dot{w}(0)\|_2^2}{2} - \frac{\|\dot{u}(t) - \dot{w}(t)\|_2^2}{2} \right) dt, \end{aligned} \quad (3.21)$$

by [15, Section 5.9, Theorem 3] and Remark 3.6. On the other hand, if $s \in (t_k^{i-1}, t_k^i]$ (for some $i \in \{2, \dots, N_k\}$), then

$$\begin{aligned} &\langle \ddot{u}_k(s) - \ddot{w}_k(s), \dot{u}_k(s) - \dot{w}_k(s) \rangle \\ &= \int_{\Omega} \left(\frac{u_k^i - 2u_k^{i-1} + u_k^{i-2}}{\delta_k^2} - \frac{w_k^i - 2w_k^{i-1} + w_k^{i-2}}{\delta_k^2} \right) \cdot \left(\frac{u_k^i - u_k^{i-1}}{\delta_k} - \frac{w_k^i - w_k^{i-1}}{\delta_k} \right) dx \\ &\geq \frac{1}{2\delta_k} \left\| \frac{u_k^i - u_k^{i-1}}{\delta_k} - \frac{w_k^i - w_k^{i-1}}{\delta_k} \right\|_2^2 - \frac{1}{2\delta_k} \left\| \frac{u_k^{i-1} - u_k^{i-2}}{\delta_k} - \frac{w_k^{i-1} - w_k^{i-2}}{\delta_k} \right\|_2^2. \end{aligned}$$

Hence, summing up for $i = 2$ to j , where $j \in \{2, \dots, N_k\}$ is such that $t_k^j = [t]_k$, leads to

$$\begin{aligned} &\int_{\delta_k}^{[t]_k} \langle \ddot{u}_k(s) - \ddot{w}_k(s), \dot{u}_k(s) - \dot{w}_k(s) \rangle ds \\ &\geq \frac{1}{2} \left\| \frac{u_k^j - u_k^{j-1}}{\delta_k} - \frac{w_k^j - w_k^{j-1}}{\delta_k} \right\|_2^2 - \frac{1}{2} \left\| \frac{u_k^1 - u_k^0}{\delta_k} - \frac{w_k^1 - w_k^0}{\delta_k} \right\|_2^2 \\ &= \frac{\|\dot{u}_k(t) - \dot{w}_k(t)\|_2^2}{2} - \frac{\|v_0 - \dot{w}(0)\|_2^2}{2}, \end{aligned}$$

where we used (3.5). Integrating this last inequality between 0 and T , and taking the liminf as $k \rightarrow \infty$ yield

$$\begin{aligned} \liminf_{k \rightarrow \infty} \int_0^T \int_{\delta_k}^{[t]_k} \langle \ddot{u}_k(s) - \ddot{w}_k(s), \dot{u}_k(s) - \dot{w}_k(s) \rangle ds dt \\ \geq \int_0^T \left(\frac{\|\dot{u}(t) - \dot{w}(t)\|_2^2}{2} - \frac{\|v_0 - \dot{w}(0)\|_2^2}{2} \right) dt. \end{aligned} \quad (3.22)$$

Gathering (3.21) and (3.22) together with the velocity initial condition (3.18) leads to

$$\liminf_{k \rightarrow \infty} I_k^2 \geq 0. \quad (3.23)$$

Finally, in view of (3.19), (3.20), and (3.23), we obtain that

$$\lim_{k \rightarrow \infty} \int_0^T (\|e_k(t) - e(t)\|_2^2 + \|z_k(t) - z(t)\|_2^2) dt = 0.$$

Eventually the strong convergences of the sequences (\hat{e}_k) and (\hat{z}_k) follow from (3.16) and (3.13). \square

We are now in position to derive the flow rule.

Corollary 3.8. *For a.e. $t \in [0, T]$,*

$$(\dot{p}(t), \dot{z}(t)) \in N_{K_\lambda}(\sigma(t), \xi(t)) \quad \text{a.e. in } \Omega.$$

Proof. According to (3.10) we have that for all $t \in [\delta_k, T]$ and a.e. $x \in \Omega$,

$$(\dot{p}_k(t), \dot{z}_k(t)) \in N_{K_\lambda}(\sigma_k(t), \xi_k(t)).$$

Thus for all $(\hat{\sigma}, \hat{\xi}) \in L^2(\Omega \times (0, T); K_\lambda)$ we have that

$$\int_{\delta_k}^T \int_{\Omega} ((\sigma_k - \hat{\sigma}) : \dot{p}_k + (\xi_k - \hat{\xi}) \dot{z}_k) dx dt \geq 0.$$

By Lemma 3.7 we have that $\sigma_k \rightarrow \sigma$ strongly in $L^2(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))$ and $\xi_k \rightarrow \xi$ strongly in $L^2(0, T; L^2(\Omega))$, while (3.14) ensures that $\dot{p}_k \rightharpoonup \dot{p}$ weakly in $L^2(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))$ and $\dot{z}_k \rightharpoonup \dot{z}$ weakly in $L^2(0, T; L^2(\Omega))$. Therefore, passing to the limit in the previous inequality yields

$$\int_0^T \int_{\Omega} ((\sigma - \hat{\sigma}) : \dot{p} + (\xi - \hat{\xi}) \dot{z}) dx dt \geq 0,$$

and the result follows by a standard localization argument. \square

Remark 3.9. Since $N_{K_\lambda} = \partial I_{K_\lambda}$ and $(\sigma(t), \xi(t)) \in K_\lambda$ a.e. in Ω for every $t \in [0, T]$, we deduce by convex duality that the flow rule is equivalent to each one of the following formulations:

(i) for a.e. $x \in \Omega$ and a.e. $t \in [0, T]$,

$$(\dot{p}(t), \dot{z}(t)) \in \partial I_{K_\lambda}(\sigma(t), \xi(t));$$

(ii) for a.e. $x \in \Omega$ and a.e. $t \in [0, T]$,

$$(\sigma(t), \xi(t)) \in \partial H_\lambda(\dot{p}(t), \dot{z}(t));$$

(iii) for a.e. $x \in \Omega$ and a.e. $t \in [0, T]$,

$$H_\lambda(\dot{p}(t), \dot{z}(t)) = \sigma(t) : \dot{p}(t) + \xi(t) \dot{z}(t);$$

(iv) for a.e. $x \in \Omega$ and a.e. $t \in [0, T]$,

$$\sigma(t) : \dot{p}(t) + \xi(t) \dot{z}(t) \geq \tau : \dot{p}(t) + \eta : \dot{z}(t) \quad \text{for every } (\tau, \eta) \in K_\lambda.$$

Note that condition (iv) is precisely Hill's principle of maximum plastic work.

3.5. Uniqueness of the solution. So far we have established the existence of solutions to the dynamical visco-plastic cap model described in Theorem 3.1. We now show the uniqueness.

Let us consider two solutions $(u_1, e_1, p_1, z_1, \sigma_1, \xi_1)$ and $(u_2, e_2, p_2, z_2, \sigma_2, \xi_2)$. Subtracting the equations of motions leads to

$$\ddot{u}_1 - \ddot{u}_2 - \operatorname{div}((\sigma_1 + \varepsilon E \dot{u}_1) - (\sigma_2 + \varepsilon E \dot{u}_2)) = 0 \quad \text{in } L^2(0, T; H^{-1}(\Omega; \mathbb{R}^n)),$$

and since $\dot{u}_1 - \dot{u}_2 \in L^2(0, T; H_0^1(\Omega; \mathbb{R}^n))$, we infer that

$$\begin{aligned} \int_0^t \langle \ddot{u}_1(s) - \ddot{u}_2(s), \dot{u}_1(s) - \dot{u}_2(s) \rangle ds + \int_0^t \int_{\Omega} (\sigma_1 - \sigma_2) : (E \dot{u}_1 - E \dot{u}_2) dx ds \\ + \varepsilon \int_0^t \int_{\Omega} |E \dot{u}_1 - E \dot{u}_2|^2 dx ds = 0. \end{aligned} \quad (3.24)$$

Since $\ddot{u}_1 - \ddot{u}_2 \in L^2(0, T; H^{-1}(\Omega; \mathbb{R}^n))$, we get from [15, Section 5.9, Theorem 3] that

$$\int_0^t \langle \ddot{u}_1(s) - \ddot{u}_2(s), \dot{u}_1(s) - \dot{u}_2(s) \rangle ds = \frac{\|\dot{u}_1(t) - \dot{u}_2(t)\|_2^2}{2} \quad (3.25)$$

since $\dot{u}_1(0) = \dot{u}_2(0) = v_0$. On the other hand, since by kinematic compatibility $E\dot{u}_1 - E\dot{u}_2 = (\dot{e}_1 - \dot{e}_2) + (\dot{p}_1 - \dot{p}_2)$, then

$$\begin{aligned} & \int_0^t \int_{\Omega} (\sigma_1 - \sigma_2) : (E\dot{u}_1 - E\dot{u}_2) dx ds \\ &= \int_0^t \int_{\Omega} (\sigma_1 - \sigma_2) : (\dot{e}_1 - \dot{e}_2) dx ds + \int_0^t \int_{\Omega} (\sigma_1 - \sigma_2) : (\dot{p}_1 - \dot{p}_2) dx ds \\ &= \mathcal{Q}(e_1(t) - e_2(t)) + \int_0^t \int_{\Omega} (\sigma_1 - \sigma_2) : (\dot{p}_1 - \dot{p}_2) dx ds, \end{aligned} \quad (3.26)$$

since $e_1(0) = e_2(0) = e_0$. In order to estimate the last integral we use the flow rule as well as the fact that $(\sigma_1, \xi_1), (\sigma_2, \xi_2) \in K_\lambda$ a.e. in $\Omega \times (0, T)$. Indeed,

$$\begin{aligned} \int_0^t \int_{\Omega} (\sigma_1 - \sigma_2) : \dot{p}_1 dx ds &\geq - \int_0^t \int_{\Omega} (\xi_1 - \xi_2) \dot{z}_1 dx ds = \int_0^t \int_{\Omega} (z_1 - z_2) \dot{z}_1 dx ds, \\ \int_0^t \int_{\Omega} (\sigma_2 - \sigma_1) : \dot{p}_2 dx ds &\geq - \int_0^t \int_{\Omega} (\xi_2 - \xi_1) \dot{z}_2 dx ds = \int_0^t \int_{\Omega} (z_2 - z_1) \dot{z}_2 dx ds, \end{aligned}$$

and summing up, we deduce that

$$\int_0^t \int_{\Omega} (\sigma_1 - \sigma_2) : (\dot{p}_1 - \dot{p}_2) dx ds \geq \int_0^t \int_{\Omega} (z_1 - z_2) (\dot{z}_1 - \dot{z}_2) dx ds = \frac{\|z_1(t) - z_2(t)\|_2^2}{2}, \quad (3.27)$$

since $z_1(0) = z_2(0) = z_0$. Gathering (3.24)–(3.27) yields $e_1 = e_2$ (hence $\sigma_1 = \sigma_2$), $z_1 = z_2$ (hence $\xi_1 = \xi_2$) and $\dot{u}_1 = \dot{u}_2$. But since $u_1(0) = u_2(0) = u_0$, we deduce that $u_1 = u_2$ and finally that $p_1 = p_2$.

Remark 3.10. Thanks to the uniqueness of the solution, there is actually no need to extract subsequences in all weak and strong convergences obtained so far.

3.6. A posteriori estimates. The object of this subsection is to prove some time regularity properties of the velocity \dot{u} , the stress σ and the hardening cap variable ξ with uniform estimates with respect to ε and λ .

Proposition 3.11. *Assume that $v_0 \in H^2(\Omega; \mathbb{R}^n)$. Then, there exists a constant $C > 0$ (independent of ε and λ) such that*

$$\begin{aligned} & \sup_{t \in [0, T]} \|\dot{u}(t)\|_2^2 + \varepsilon \int_0^T \int_{\Omega} |E\ddot{u}|^2 dx ds + \sup_{t \in [0, T]} \|\dot{\sigma}(t)\|_2^2 + \sup_{t \in [0, T]} \|\dot{z}(t)\|_2^2 \\ & \leq C \left(\varepsilon^2 \|\Delta v_0\|_2^2 + \|Ev_0\|_2^2 + \|\dot{f}\|_{L^1(0, T; L^2(\Omega; \mathbb{R}^n))}^2 \right. \\ & \quad \left. + \|\ddot{w}\|_{L^1(0, T; L^2(\Omega; \mathbb{R}^n))}^2 + \|E\ddot{w}\|_{L^2(0, T; L^2(\Omega; \mathbb{M}_{sym}^n))}^2 \right). \end{aligned} \quad (3.28)$$

Proof. We will use the difference quotient method. Let us extend continuously the fields for $s < 0$ by setting

$$\begin{cases} u(s) = u_0 + sv_0, & w(s) = w(0) + s\dot{w}(0), & f(s) = f(0), \\ e(s) = e_0, & \sigma(s) = \sigma_0, \\ p(s) = p_0, \\ z(s) = z_0, & \xi(s) = \xi_0. \end{cases} \quad (3.29)$$

In the following we will use the notation

$$\partial_t^h g(s) := \frac{g(s) - g(s-h)}{h}.$$

Let $t \in [0, T]$ and $h < t$. Using the equation of motion, we have for all $\varphi \in L^2(0, T+h; H_0^1(\Omega; \mathbb{R}^n))$,

$$\begin{aligned} \int_0^T \langle \ddot{u}(s), \varphi(s) \rangle ds + \int_0^T \int_{\Omega} (\sigma(s) + \varepsilon E \dot{u}(s)) : E \varphi(s) dx ds &= \int_0^T \int_{\Omega} f(s) \cdot \varphi(s) dx ds, \\ \int_h^{T+h} \langle \ddot{u}(s-h), \varphi(s) \rangle ds + \int_h^{T+h} \int_{\Omega} (\sigma(s-h) + \varepsilon E \dot{u}(s-h)) : E \varphi(s) dx ds \\ &= \int_h^{T+h} \int_{\Omega} f(s-h) \cdot \varphi(s) dx ds. \end{aligned}$$

Taking the difference of the two previous equalities with the test function $\varphi = \chi_{[0,t]} \partial_t^h(\dot{u} - \dot{w})$ yields

$$\begin{aligned} \int_0^t \langle \partial_t^h \ddot{u}(s), \partial_t^h(\dot{u} - \dot{w})(s) \rangle ds + \int_0^t \int_{\Omega} \partial_t^h(\sigma + \varepsilon E \dot{u})(s) : E(\partial_t^h(\dot{u} - \dot{w}))(s) dx ds \\ - \int_0^t \int_{\Omega} \partial_t^h f(s) \cdot \partial_t^h(\dot{u} - \dot{w}) dx ds \\ = \frac{1}{h} \int_0^h \int_{\Omega} [f(0) \cdot \partial_t^h(\dot{u} - \dot{w})(s) - (\sigma_0 + \varepsilon E v_0) : E(\partial_t^h(\dot{u} - \dot{w}))(s)] dx ds \\ = \frac{1}{h} \int_0^h \int_{\Omega} \varepsilon \Delta v_0 \cdot \partial_t^h(\dot{u} - \dot{w})(s) dx ds, \end{aligned}$$

where we used the initial condition $-\operatorname{div} \sigma_0 = f(0)$ a.e. in Ω . Hence thanks to the Cauchy-Schwarz inequality,

$$\begin{aligned} \int_0^t \langle \partial_t^h \ddot{u}(s), \partial_t^h(\dot{u} - \dot{w})(s) \rangle ds + \int_0^t \int_{\Omega} \partial_t^h(\sigma + \varepsilon E \dot{u})(s) : E(\partial_t^h(\dot{u} - \dot{w}))(s) dx ds \\ \leq (\varepsilon \|\Delta v_0\|_2 + \|\partial_t^h f\|_{L^1(0,T;L^2(\Omega;\mathbb{R}^n))}) \sup_{s \in [0,T]} \|\partial_t^h(\dot{u} - \dot{w})(s)\|_2. \quad (3.30) \end{aligned}$$

Next using the kinematic compatibility for the rates $E \dot{u} = \dot{e} + \dot{p}$ a.e. on $\Omega \times [0, T]$, we get for all $\tau \in L^2(0, T+h; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))$,

$$\begin{aligned} \int_0^T \int_{\Omega} E \dot{u}(s) : \tau(s) dx ds &= \int_0^T \int_{\Omega} \dot{e}(s) : \tau(s) dx ds + \int_0^T \int_{\Omega} \dot{p}(s) : \tau(s) dx ds, \\ \int_h^{T+h} \int_{\Omega} E \dot{u}(s-h) : \tau(s) dx ds &= \int_h^{T+h} \int_{\Omega} \dot{e}(s-h) : \tau(s) dx ds + \int_h^{T+h} \int_{\Omega} \dot{p}(s-h) : \tau(s) dx ds. \end{aligned}$$

Taking the difference of the two previous relations with the test function $\tau = \chi_{[0,t]}(\partial_t^h \sigma)$ yields

$$\begin{aligned} \int_0^t \int_{\Omega} E \partial_t^h \dot{u}(s) : \partial_t^h \sigma(s) dx ds - \int_0^t \int_{\Omega} \partial_t^h \dot{e}(s) : \partial_t^h \sigma(s) dx ds - \int_0^t \int_{\Omega} \partial_t^h \dot{p}(s) : \partial_t^h \sigma(s) dx ds \\ = -\frac{1}{h} \int_0^h \int_{\Omega} E v_0 : \partial_t^h \sigma(s) dx ds \leq \|E v_0\|_2 \sup_{s \in [0,h]} \|\partial_t^h \sigma(s)\|_2. \quad (3.31) \end{aligned}$$

According to the flow rule, since for a.e. $s \in (0, T)$,

$$(\dot{p}(s), \dot{z}(s)) \in N_{K_{\lambda}}(\sigma(s), \xi(s)), \quad (\dot{p}(s-h), \dot{z}(s-h)) \in N_{K_{\lambda}}(\sigma(s-h), \xi(s-h)),$$

then

$$\begin{aligned}
 & \int_0^t \int_{\Omega} \partial_t^h \dot{p}(s) : \partial_t^h \sigma(s) \, dx \, ds \\
 &= \frac{1}{h^2} \int_0^t \int_{\Omega} \dot{p}(s) : (\sigma(s) - \sigma(s-h)) \, dx \, ds + \frac{1}{h^2} \int_0^t \int_{\Omega} \dot{p}(s-h) : (\sigma(s-h) - \sigma(s)) \, dx \, ds \\
 &\geq -\frac{1}{h^2} \int_0^t \int_{\Omega} \dot{z}(s)(\xi(s) - \xi(s-h)) \, dx \, ds - \frac{1}{h^2} \int_0^t \int_{\Omega} \dot{z}(s-h)(\xi(s-h) - \xi(s)) \, dx \, ds \\
 &= \frac{1}{h^2} \int_0^t \int_{\Omega} \dot{z}(s)(z(s) - z(s-h)) \, dx \, ds + \frac{1}{h^2} \int_0^t \int_{\Omega} \dot{z}(s-h)(z(s-h) - z(s)) \, dx \, ds \\
 &= \int_0^t \int_{\Omega} \partial_t^h \dot{z}(s) \partial_t^h z(s) \, dx \, ds. \quad (3.32)
 \end{aligned}$$

Gathering (3.30)–(3.32), we obtain that

$$\begin{aligned}
 & \int_0^t \langle \partial_t^h (\ddot{u} - \ddot{w})(s), \partial_t^h (\dot{u} - \dot{w})(s) \rangle \, ds + \varepsilon \int_0^t \int_{\Omega} |\partial_t^h (E\dot{u})|^2 \, dx \, ds \\
 & \quad + \int_0^t \int_{\Omega} \partial_t^h \dot{e}(s) : \partial_t^h \sigma(s) \, dx \, ds + \int_0^t \int_{\Omega} \partial_t^h \dot{z}(s) \partial_t^h z(s) \, dx \, ds \\
 & \leq (\varepsilon \|\Delta v_0\|_2 + \|\partial_t^h f\|_{L^1(0,T;L^2(\Omega;\mathbb{R}^n))} + \|\partial_t^h \ddot{w}\|_{L^1(0,T;L^2(\Omega;\mathbb{R}^n))}) \sup_{s \in [0,T]} \|\partial_t^h (\dot{u} - \dot{w})(s)\|_2 \\
 & \quad + \left(\|Ev_0\|_2 + \|E(\partial_t^h \dot{w})\|_{L^1(0,T;L^2(\Omega;\mathbb{M}_{sym}^{n \times n}))} \right) \sup_{s \in [0,T]} \|\partial_t^h \sigma(s)\|_2 \\
 & \quad + \varepsilon \|E(\partial_t^h \dot{w})\|_{L^2(0,T;L^2(\Omega;\mathbb{M}_{sym}^{n \times n}))} \|E(\partial_t^h \dot{u})\|_{L^2(0,T;L^2(\Omega;\mathbb{M}_{sym}^{n \times n}))},
 \end{aligned}$$

and thus, since $\partial_t^h (\dot{u} - \dot{w}) \in L^2(0, T; H_0^1(\Omega; \mathbb{R}^n))$ and $\partial_t^h (\ddot{u} - \ddot{w}) \in L^2(0, T; H^{-1}(\Omega; \mathbb{R}^n))$, we get from [15, Section 5.9, Theorem 3] that

$$\begin{aligned}
 & \frac{\|\partial_t^h (\dot{u} - \dot{w})(t)\|_2^2}{2} + \varepsilon \int_0^t \int_{\Omega} |\partial_t^h (E\dot{u})|^2 \, dx \, ds + \mathcal{Q}(\partial_t^h e(t)) + \frac{\|\partial_t^h z(t)\|_2^2}{2} \\
 & \leq (\varepsilon \|\Delta v_0\|_2 + \|\partial_t^h f\|_{L^1(0,T;L^2(\Omega;\mathbb{R}^n))} + \|\partial_t^h \ddot{w}\|_{L^1(0,T;L^2(\Omega;\mathbb{R}^n))}) \sup_{s \in [0,T]} \|\partial_t^h (\dot{u} - \dot{w})(s)\|_2 \\
 & \quad + \left(\|Ev_0\|_2 + \|E(\partial_t^h \dot{w})\|_{L^1(0,T;L^2(\Omega;\mathbb{M}_{sym}^{n \times n}))} \right) \sup_{s \in [0,T]} \|\partial_t^h \sigma(s)\|_2 \\
 & \quad + \varepsilon \|E(\partial_t^h \dot{w})\|_{L^2(0,T;L^2(\Omega;\mathbb{M}_{sym}^{n \times n}))} \|E(\partial_t^h \dot{u})\|_{L^2(0,T;L^2(\Omega;\mathbb{M}_{sym}^{n \times n}))} \\
 & \quad + \frac{\|\partial_t^h (\dot{u} - \dot{w})(0)\|_2^2}{2} + \mathcal{Q}(\partial_t^h e(0)) + \frac{\|\partial_t^h z(0)\|_2^2}{2}.
 \end{aligned}$$

Hence, applying Young's inequality, and according to the choice of the extensions (3.29), we obtain

$$\begin{aligned}
 & \sup_{t \in [0,T]} \|\partial_t^h (\dot{u} - \dot{w})(t)\|_2^2 + \varepsilon \int_0^T \int_{\Omega} |\partial_t^h (E\dot{u})|^2 \, dx \, ds + \sup_{t \in [0,T]} \|\partial_t^h \sigma(t)\|_2^2 + \sup_{t \in [0,T]} \|\partial_t^h z(t)\|_2^2 \\
 & \leq C \left(\varepsilon^2 \|\Delta v_0\|_2^2 + \|\partial_t^h f\|_{L^1(0,T;L^2(\Omega;\mathbb{R}^n))}^2 + \|\partial_t^h \ddot{w}\|_{L^1(0,T;L^2(\Omega;\mathbb{R}^n))}^2 + \|Ev_0\|_2^2 \right. \\
 & \quad \left. + \|E(\partial_t^h \dot{w})\|_{L^2(0,T;L^2(\Omega;\mathbb{M}_{sym}^{n \times n}))}^2 \right),
 \end{aligned}$$

for some constant $C > 0$ independent of ε , λ and h . Letting $h \rightarrow 0$ leads to

$$\begin{aligned} & \sup_{t \in [0, T]} \|\ddot{u}(t)\|_2^2 + \varepsilon \int_0^T \int_{\Omega} |E\ddot{u}|^2 dx ds + \sup_{t \in [0, T]} \|\dot{\sigma}(t)\|_2^2 + \sup_{t \in [0, T]} \|\dot{z}(t)\|_2^2 \\ & \leq C \left(\varepsilon^2 \|\Delta v_0\|_2^2 + \|f\|_{L^1(0, T; L^2(\Omega; \mathbb{R}^n))}^2 + \|E v_0\|_2^2 + \|\ddot{w}\|_{L^1(0, T; L^2(\Omega; \mathbb{R}^n))}^2 + \|E\ddot{w}\|_{L^2(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))}^2 \right), \end{aligned}$$

which is (3.28). \square

4. THE DYNAMICAL ELASTO-PLASTIC CAP MODEL

In this section, we pass to the limit as the viscosity parameter ε tends to zero in order to recover a solution for the dynamical elasto-plastic cap model (1.3) from the visco-plastic evolutions obtained in Theorem 3.1. In this case, due to a lack of coercivity in reflexive Sobolev spaces, the space of kinematically admissible fields needs to be relaxed in the following way: given a boundary displacement $\hat{w} \in H^1(\Omega; \mathbb{R}^n)$, we set

$$\begin{aligned} \mathcal{A}_{\text{dyn}}(\hat{w}) := & \left\{ (v, \eta, q) \in (BD(\Omega) \cap L^2(\Omega; \mathbb{R}^n)) \times L^2(\Omega; \mathbb{M}_{sym}^{n \times n}) \times \mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n}) : \right. \\ & \left. E v = \eta + q \text{ in } \Omega, \quad q = (\hat{w} - v) \odot \nu \mathcal{H}^{n-1} \text{ on } \partial\Omega \right\}. \end{aligned}$$

Indeed, it may happen that a kinematically admissible displacement v does not match the prescribed boundary value \hat{w} on some portion of the boundary (of positive \mathcal{H}^{n-1} measure). In that case, on this portion of the boundary a plastic strain q must develop, compatible with the fact that q is the jump part of the measure $E v$.

As in the previous section, we fix $\lambda \geq 1$ and consider a body load $f \in AC([0, T]; L^2(\Omega; \mathbb{R}^n))$ and a boundary displacement which is the trace on $\partial\Omega \times (0, T)$ of a function $w \in H^2([0, T]; H^1(\Omega; \mathbb{R}^n)) \cap H^3([0, T]; L^2(\Omega; \mathbb{R}^n))$. We also consider initial data $(u_0, e_0, p_0, z_0) \in \mathcal{A}_{\text{dyn}}(w(0)) \times L^2(\Omega)$ and $v_0 \in H^1(\Omega; \mathbb{R}^n)$ satisfying

$$v_0 = \dot{w}(0) \quad \mathcal{H}^{n-1}\text{-a.e. on } \partial\Omega,$$

and

$$(\sigma_0, \xi_0) := (\mathbb{C}e_0, -z_0) \in K_\lambda, \quad -\text{div}\sigma_0 = f(0) \quad \text{a.e. in } \Omega.$$

The main result of this section is the following existence and uniqueness result for a dynamical elasto-plastic cap model, obtained as a vanishing viscosity limit of the visco-plastic cap evolutions constructed in Theorem 3.1.

Theorem 4.1. *There exist unique*

$$\begin{cases} u \in AC([0, T]; BD(\Omega)) \cap W^{2, \infty}([0, T]; L^2(\Omega; \mathbb{R}^n)), \\ \sigma, e \in W^{1, \infty}([0, T]; L^2(\Omega; \mathbb{M}_{sym}^{n \times n})), \\ p \in AC([0, T]; \mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n})), \\ \xi, z \in W^{1, \infty}([0, T]; L^2(\Omega)), \end{cases}$$

with the following properties: for all $t \in [0, T]$,

$$\begin{cases} E u(t) = e(t) + p(t) \text{ in } \Omega, & p(t) = (w(t) - u(t)) \odot \nu \mathcal{H}^{n-1} \text{ on } \partial\Omega, \\ \sigma(t) = \mathbb{C}e(t), & \xi(t) = -z(t), \\ (\sigma(t), \xi(t)) \in K_\lambda \text{ a.e. in } \Omega, \end{cases} \quad (4.1)$$

and

$$\begin{cases} \ddot{u} - \text{div}\sigma = f \quad \text{a.e. in } \Omega \times (0, T), \\ (u(0), e(0), p(0), z(0)) = (u_0, e_0, p_0, z_0), \quad \dot{u}(0) = v_0. \end{cases} \quad (4.2)$$

Moreover, for a.e. $t \in [0, T]$,

$$\dot{z}(t) \geq 0 \quad \text{a.e. in } \Omega \quad (4.3)$$

and the distribution $[\sigma(t) : \dot{p}(t)]$ is a measure in $\mathcal{M}(\bar{\Omega})$ satisfying

$$H_\lambda(\dot{p}(t), \dot{z}(t)) = [\sigma(t) : \dot{p}(t)] + \xi(t)\dot{z}(t) \quad \text{in } \mathcal{M}(\bar{\Omega}). \quad (4.4)$$

Remark 4.2. Within the proof of Theorem 4.1, we will also prove the following bound: there exists a constant $C > 0$ (independent of λ) such that

$$\begin{aligned} & \|\ddot{u}\|_{L^\infty(0,T;L^2(\Omega;\mathbb{R}^n))} + \|\dot{e}\|_{L^\infty(0,T;L^2(\Omega;\mathbb{M}_{sym}^{n \times n}))} + \|\dot{z}\|_{L^\infty(0,T;L^2(\Omega))} \\ & \leq C \left(\|Ev_0\|_2 + \|\dot{f}\|_{L^1(0,T;L^2(\Omega;\mathbb{R}^n))} + \|\ddot{w}\|_{L^1(0,T;L^2(\Omega;\mathbb{R}^n))} + \|E\dot{w}\|_{L^2(0,T;L^2(\Omega;\mathbb{M}_{sym}^{n \times n}))} \right). \end{aligned} \quad (4.5)$$

The remaining of the section is devoted to the proof of Theorem 4.1.

4.1. A priori estimates and compactness. In order to apply the result of Theorem 3.1, we need to regularize the initial data. According to [8, Lemma 5.1], there exists a sequence $(u_{0,\varepsilon}) \subset H^1(\Omega; \mathbb{R}^n)$ such that $u_{0,\varepsilon} = w(0)$ \mathcal{H}^{n-1} -a.e. on $\partial\Omega$, $u_{0,\varepsilon} \rightarrow u_0$ strongly in $L^1(\Omega; \mathbb{R}^n)$, and $Eu_{0,\varepsilon} \rightarrow Eu_0$ weakly* in $\mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n})$. Setting $p_{0,\varepsilon} = Eu_{0,\varepsilon} - e_0$, we get that $(u_{0,\varepsilon}, e_0, p_{0,\varepsilon}) \in \mathcal{A}_{vp}(w(0))$. Moreover, using a standard approximation argument we can find a sequence $(v_{0,\varepsilon}) \subset H^2(\Omega; \mathbb{R}^n)$ such that $v_{0,\varepsilon} \rightarrow v_0$ strongly in $H^1(\Omega; \mathbb{R}^n)$ and $\varepsilon \Delta v_{0,\varepsilon} \rightarrow 0$ strongly in $L^2(\Omega; \mathbb{R}^n)$. For every $\varepsilon > 0$ let $u_\varepsilon, e_\varepsilon, \sigma_\varepsilon, p_\varepsilon, z_\varepsilon, \xi_\varepsilon$ be the unique solution of (3.2)–(3.4) with initial data $(u_{0,\varepsilon}, e_0, p_{0,\varepsilon}, z_0, v_{0,\varepsilon})$, whose existence is guaranteed by Theorem 3.1.

We first show that an energy equality holds.

Proposition 4.3. For every $t \in [0, T]$,

$$\begin{aligned} \mathcal{Q}(e_\varepsilon(t)) + \frac{1}{2}\|z_\varepsilon(t)\|_2^2 + \int_0^t \mathcal{H}_\lambda(\dot{p}_\varepsilon(s), \dot{z}_\varepsilon(s)) ds + \frac{1}{2}\|\dot{u}_\varepsilon(t)\|_2^2 + \varepsilon \int_0^t \int_\Omega |E\dot{u}_\varepsilon|^2 dx ds \\ = \mathcal{Q}(e_0) + \frac{1}{2}\|z_0\|_2^2 + \frac{1}{2}\|v_{0,\varepsilon}\|_2^2 + \int_0^t \int_\Omega (\sigma_\varepsilon + \varepsilon E\dot{u}_\varepsilon) : E\dot{w} dx ds \\ + \int_0^t \int_\Omega \ddot{u}_\varepsilon \cdot \dot{w} dx ds + \int_0^t \int_\Omega f \cdot (\dot{u}_\varepsilon - \dot{w}) dx ds. \end{aligned} \quad (4.6)$$

Proof. Multiplying the equation of motion (3.3) by $(\dot{u}_\varepsilon - \dot{w})\chi_{[0,t]} \in L^2(0, T; H_0^1(\Omega; \mathbb{R}^n))$, and integrating by parts yield

$$\int_0^t \int_\Omega \ddot{u}_\varepsilon \cdot (\dot{u}_\varepsilon - \dot{w}) dx ds + \int_0^t \int_\Omega (\sigma_\varepsilon + \varepsilon E\dot{u}_\varepsilon) : (E\dot{u}_\varepsilon - E\dot{w}) dx ds = \int_0^t \int_\Omega f \cdot (\dot{u}_\varepsilon - \dot{w}) dx ds.$$

On the other hand, by the kinematic compatibility for the rates (3.15) $E\dot{u}_\varepsilon = \dot{e}_\varepsilon + \dot{p}_\varepsilon$ a.e. in $\Omega \times [0, T]$ and the flow rule in (3.4) we have

$$\begin{aligned} \int_0^t \int_\Omega \sigma_\varepsilon : E\dot{u}_\varepsilon dx ds &= \int_0^t \int_\Omega \sigma_\varepsilon : \dot{e}_\varepsilon dx ds + \int_0^t \mathcal{H}_\lambda(\dot{p}_\varepsilon(s), \dot{z}_\varepsilon(s)) ds - \int_0^t \int_\Omega \xi_\varepsilon \dot{z}_\varepsilon dx ds \\ &= \mathcal{Q}(e_\varepsilon(t)) - \mathcal{Q}(e_0) + \int_0^t \mathcal{H}_\lambda(\dot{p}_\varepsilon(s), \dot{z}_\varepsilon(s)) ds + \frac{1}{2}\|z_\varepsilon(t)\|_2^2 - \frac{1}{2}\|z_0\|_2^2, \end{aligned}$$

where we used that $\xi_\varepsilon = -z_\varepsilon$. Combining the two previous equalities and applying [15, Section 5.9, Theorem 3], we obtain that (4.6) holds. \square

Remark 4.4. Since $\operatorname{div}(\sigma_\varepsilon + \varepsilon E\dot{u}_\varepsilon) = \ddot{u}_\varepsilon - f \in L^2(0, T; L^2(\Omega; \mathbb{R}^n))$, we deduce that the normal trace $(\sigma_\varepsilon + \varepsilon E\dot{u}_\varepsilon)\nu \in L^2(0, T; H^{-1/2}(\partial\Omega; \mathbb{R}^n))$. Using an integration by parts, the term

$$\int_0^t \int_\Omega (\sigma_\varepsilon + \varepsilon E\dot{u}_\varepsilon) : E\dot{w} \, dx \, ds + \int_0^t \int_\Omega \ddot{u}_\varepsilon \cdot \dot{w} \, dx \, ds + \int_0^t \int_\Omega f \cdot (\dot{u}_\varepsilon - \dot{w}) \, dx \, ds$$

can be equivalently rewritten as

$$\int_0^t \langle (\sigma_\varepsilon(s) + \varepsilon E\dot{u}_\varepsilon(s))\nu, \dot{w}(s) \rangle \, ds + \int_0^t \int_\Omega f \cdot \dot{u}_\varepsilon \, dx \, ds,$$

where the bracket $\langle \cdot, \cdot \rangle$ stands for the duality pairing between $H^{-1/2}(\partial\Omega; \mathbb{R}^n)$ and $H^{1/2}(\partial\Omega; \mathbb{R}^n)$, which is precisely a weak formulation of the work of internal and external forces.

According to Proposition 3.11, there exists a constant $C_1 > 0$ (independent of ε and λ) such that

$$\begin{aligned} & \sup_{t \in [0, T]} \|\ddot{u}_\varepsilon(t)\|_2 + \sqrt{\varepsilon} \|E\dot{u}_\varepsilon\|_{L^2(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))} + \sup_{t \in [0, T]} \|\dot{e}_\varepsilon(t)\|_2 + \sup_{t \in [0, T]} \|\dot{z}_\varepsilon(t)\|_2 \\ & \leq C_1 \left(\varepsilon \|\Delta v_{0, \varepsilon}\|_2 + \|Ev_{0, \varepsilon}\|_2 + \|\dot{f}\|_{L^1(0, T; L^2(\Omega; \mathbb{R}^n))} \right. \\ & \quad \left. + \|\ddot{w}\|_{L^1(0, T; L^2(\Omega; \mathbb{R}^n))} + \|E\dot{w}\|_{L^2(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))} \right). \end{aligned} \quad (4.7)$$

Moreover, as a consequence of Proposition 4.3, and by applying the coercivity properties (2.1), (2.3), and Hölder inequality, we deduce that there is a constant $C > 0$ (independent of ε and λ) such that

$$\begin{aligned} & \frac{\alpha_C}{2} \|e_\varepsilon(t)\|_2^2 + \frac{1}{2} \|z_\varepsilon(t)\|_2^2 + \alpha_H \int_0^t \|\dot{p}_\varepsilon(s)\|_1 \, ds + \frac{1}{2} \|\dot{u}_\varepsilon(t)\|_2^2 + \varepsilon \int_0^t \int_\Omega |E\dot{u}_\varepsilon|^2 \, dx \, ds \\ & \leq C \left(1 + \sup_{s \in [0, T]} \|e_\varepsilon(s)\|_2 + \sup_{s \in [0, T]} \|\dot{u}_\varepsilon(s)\|_2 + \varepsilon \|E\dot{u}_\varepsilon\|_{L^2(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))} \right) \\ & \quad + \frac{\alpha_H}{\sqrt{n}} \int_0^T \|\dot{z}_\varepsilon(s)\|_1 \, ds, \end{aligned}$$

where we used the fact that (\dot{u}_ε) is uniformly bounded in $L^\infty(0, T; L^2(\Omega; \mathbb{R}^n))$ by (4.7), and that $(v_{0, \varepsilon})$ is uniformly bounded in $L^2(\Omega; \mathbb{R}^n)$. By (4.7) again, we also have that \dot{z}_ε is uniformly bounded in $L^\infty(0, T; L^2(\Omega))$; thus, by Young inequality we conclude that

$$\sup_{t \in [0, T]} \|\dot{u}_\varepsilon(t)\|_2 + \sqrt{\varepsilon} \|E\dot{u}_\varepsilon\|_{L^2(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))} + \sup_{t \in [0, T]} \|e_\varepsilon(t)\|_2 + \sup_{t \in [0, T]} \|z_\varepsilon(t)\|_2 \leq C_2, \quad (4.8)$$

$$\int_0^T \|\dot{p}_\varepsilon(s)\|_1 \, ds \leq C_3, \quad (4.9)$$

for some constants $C_2 > 0$ and $C_3 > 0$ both independent of ε and λ .

Let Ω' be a smooth and bounded open set such that $\bar{\Omega} \subset \Omega'$. For every $t \in [0, T]$ we extend $u_\varepsilon(t)$, $e_\varepsilon(t)$, $p_\varepsilon(t)$, and $z_\varepsilon(t)$ to Ω' by setting $u_\varepsilon(t) = w(t)$, $e_\varepsilon(t) = Ew(t)$, $p_\varepsilon(t) = 0$, $z_\varepsilon(t) = 0$ in $\Omega' \setminus \Omega$. Clearly the bounds (4.7), (4.8), and (4.9) remain satisfied if we replace Ω by Ω' .

By (4.7) and (4.8) we deduce the existence of three functions $v \in W^{1,\infty}([0, T]; L^2(\Omega'; \mathbb{R}^n))$, $e \in W^{1,\infty}([0, T]; L^2(\Omega'; \mathbb{M}_{sym}^{n \times n}))$, and $z \in W^{1,\infty}([0, T]; L^2(\Omega'))$ such that, up to a subsequence,

$$\begin{aligned} \dot{u}_\varepsilon &\rightharpoonup v \text{ weakly}^* \text{ in } L^\infty(0, T; L^2(\Omega'; \mathbb{R}^n)), \\ \dot{u}_\varepsilon &\rightharpoonup \dot{v} \text{ weakly}^* \text{ in } L^\infty(0, T; L^2(\Omega'; \mathbb{R}^n)), \end{aligned} \quad (4.10)$$

$$e_\varepsilon \rightharpoonup e \text{ weakly}^* \text{ in } L^\infty(0, T; L^2(\Omega'; \mathbb{M}_{sym}^{n \times n})), \quad (4.11)$$

$$\begin{aligned} \dot{e}_\varepsilon &\rightharpoonup \dot{e} \text{ weakly}^* \text{ in } L^\infty(0, T; L^2(\Omega'; \mathbb{M}_{sym}^{n \times n})), \\ z_\varepsilon &\rightharpoonup z \text{ weakly}^* \text{ in } L^\infty(0, T; L^2(\Omega')), \end{aligned} \quad (4.12)$$

$$\dot{z}_\varepsilon \rightharpoonup \dot{z} \text{ weakly}^* \text{ in } L^\infty(0, T; L^2(\Omega')).$$

By Ascoli-Arzelà Theorem and the bounds (4.7) and (4.8), we also have that for every $t \in [0, T]$,

$$\dot{u}_\varepsilon(t) \rightharpoonup v(t) \text{ weakly in } L^2(\Omega'; \mathbb{R}^n), \quad (4.13)$$

$$e_\varepsilon(t) \rightharpoonup e(t) \text{ weakly in } L^2(\Omega'; \mathbb{M}_{sym}^{n \times n}), \quad (4.14)$$

$$z_\varepsilon(t) \rightharpoonup z(t) \text{ weakly in } L^2(\Omega'). \quad (4.15)$$

Moreover, by (3.4) we have

$$\dot{z}(t) \geq 0 \quad \text{a.e. in } \Omega, \quad \text{for a.e. } t \in [0, T]. \quad (4.16)$$

We next establish some compactness on the sequence (p_ε) of plastic strains. By (4.9) and Helly Theorem (see [23, Theorem 3.2] or [7, Lemma 7.2]) we deduce the existence of a subsequence (not relabeled) and of a function $p \in BV([0, T]; \mathcal{M}(\Omega'; \mathbb{M}_{sym}^{n \times n}))$ such that for every $t \in [0, T]$,

$$p_\varepsilon(t) \rightharpoonup p(t) \text{ weakly}^* \text{ in } \mathcal{M}(\Omega'; \mathbb{M}_{sym}^{n \times n}). \quad (4.17)$$

Finally we state some compactness properties for the sequence (u_ε) of displacements. By (4.14) and (4.17) we have that $Eu_\varepsilon(t) \rightharpoonup e(t) + p(t)$ weakly* in $\mathcal{M}(\Omega'; \mathbb{M}_{sym}^{n \times n})$. Since $u_\varepsilon(t) = w(t)$ a.e. on $\Omega' \setminus \Omega$, we deduce by the Poincaré-Korn inequality (see [34, Chapter II, Remark 2.5 (ii)]) that the sequence $(u_\varepsilon(t))$ is uniformly bounded in $BD(\Omega')$. Therefore, for every $t \in [0, T]$ there exist a subsequence (ε_j) , possibly depending on t , and a function $u(t) \in BD(\Omega')$ such that $u_{\varepsilon_j}(t) \rightharpoonup u(t)$ weakly* in $BD(\Omega')$. Since $u(t) = w(t)$ a.e. in $\Omega' \setminus \Omega$ and $Eu(t) = e(t) + p(t)$ in Ω' , we conclude again by the Poincaré-Korn inequality that the limit $u(t)$ is uniquely determined. Therefore, the convergence result holds for the whole sequence, that is,

$$u_\varepsilon(t) \rightharpoonup u(t) \text{ weakly}^* \text{ in } BD(\Omega') \quad \text{for every } t \in [0, T]. \quad (4.18)$$

In particular, we have shown that for every $t \in [0, T]$,

$$Eu(t) = e(t) + p(t) \text{ in } \Omega, \quad p(t) = (w(t) - u(t)) \odot \nu \mathcal{H}^{n-1} \text{ on } \partial\Omega. \quad (4.19)$$

Moreover, by (4.13) and (4.18) we infer that $v(t) = \dot{u}(t)$ for every $t \in [0, T]$, hence

$$u \in W^{2,\infty}([0, T]; L^2(\Omega; \mathbb{R}^n)).$$

Clearly, from the convergences of the initial data $(u_{0,\varepsilon})$ and $(v_{0,\varepsilon})$, the initial conditions $u(0) = u_0$, $e(0) = e_0$, $p(0) = p_0$, $z(0) = z_0$, and $\dot{u}(0) = v_0$ are satisfied. Inequality (4.5) is an immediate consequence of (4.7).

We define $\sigma(t) := \mathbb{C}e(t)$ and $\xi(t) := -z(t)$. Since $(\sigma_\varepsilon(t), \xi_\varepsilon(t)) \in K_\lambda$ a.e. in Ω and K_λ is a closed and convex set, by (4.14) and (4.15) we immediately deduce that $(\sigma(t), \xi(t)) \in K_\lambda$ a.e. in Ω .

Let $\varphi \in C_c^\infty(\Omega \times (0, T); \mathbb{R}^n)$. By the equation of motion in (3.3) we have

$$\int_0^T \int_\Omega \ddot{u}_\varepsilon \cdot \varphi \, dx \, dt + \int_0^T \int_\Omega (\sigma_\varepsilon + \varepsilon E \dot{u}_\varepsilon) : E \varphi \, dx \, dt = \int_0^T \int_\Omega f \cdot \varphi \, dx \, dt.$$

Since $\varepsilon E\dot{u}_\varepsilon \rightarrow 0$ strongly in $L^2(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))$ by (4.8), we can pass to the limit in the above equality and obtain

$$\int_0^T \int_\Omega \ddot{u} \cdot \varphi \, dx \, dt + \int_0^T \int_\Omega \sigma : E\varphi \, dx \, dt = \int_0^T \int_\Omega f \cdot \varphi \, dx \, dt,$$

which implies

$$\ddot{u} - \operatorname{div} \sigma = f \quad \text{a.e. in } \Omega \times (0, T).$$

4.2. Strong convergences and flow rule. We first improve the weak convergences established at the previous section into strong convergences.

Lemma 4.5. *The following strong convergences hold:*

$$\dot{u}_\varepsilon \rightarrow \dot{u} \text{ strongly in } L^\infty(0, T; L^2(\Omega; \mathbb{R}^n)), \quad (4.20)$$

$$e_\varepsilon \rightarrow e \text{ strongly in } L^\infty(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n})), \quad (4.21)$$

$$z_\varepsilon \rightarrow z \text{ strongly in } L^\infty(0, T; L^2(\Omega)), \quad (4.22)$$

$$\sqrt{\varepsilon} E\dot{u}_\varepsilon \rightarrow 0 \text{ strongly in } L^2(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n})). \quad (4.23)$$

Proof. We first observe that the flow rule in (3.4), together with the fact that $(\sigma(t), \xi(t)) \in K_\lambda$ a.e. in Ω , implies that for a.e. $t \in [0, T]$

$$\int_\Omega \dot{p}_\varepsilon(t) : (\sigma_\varepsilon(t) - \sigma(t)) \, dx + \int_\Omega \dot{z}_\varepsilon(t)(\xi_\varepsilon(t) - \xi(t)) \, dx \geq 0.$$

Using the kinematic compatibility for the rates (3.15) $E\dot{u}_\varepsilon(t) = \dot{e}_\varepsilon(t) + \dot{p}_\varepsilon(t)$ a.e. in Ω , and the definition of $\xi_\varepsilon(t)$ and $\xi(t)$, we obtain

$$\begin{aligned} \int_\Omega \dot{e}_\varepsilon(t) : (\sigma_\varepsilon(t) - \sigma(t)) \, dx + \int_\Omega \dot{z}_\varepsilon(t)(z_\varepsilon(t) - z(t)) \, dx \\ - \int_\Omega E\dot{u}_\varepsilon(t) : (\sigma_\varepsilon(t) + \varepsilon E\dot{u}_\varepsilon(t) - \sigma(t)) \, dx + \varepsilon \int_\Omega |E\dot{u}_\varepsilon(t)|^2 \, dx \leq 0. \end{aligned}$$

Integration by parts yields

$$\begin{aligned} \int_\Omega (\dot{e}_\varepsilon(t) - \dot{e}(t)) : (\sigma_\varepsilon(t) - \sigma(t)) \, dx + \int_\Omega (\dot{z}_\varepsilon(t) - \dot{z}(t))(z_\varepsilon(t) - z(t)) \, dx \\ + \int_\Omega (\ddot{u}_\varepsilon(t) - \ddot{u}(t)) \cdot (\dot{u}_\varepsilon(t) - \dot{u}(t)) \, dx + \varepsilon \int_\Omega |E\dot{u}_\varepsilon(t)|^2 \, dx \\ \leq - \int_\Omega \dot{e}(t) : (\sigma_\varepsilon(t) - \sigma(t)) \, dx - \int_\Omega \dot{z}(t)(z_\varepsilon(t) - z(t)) \, dx - \int_\Omega (\ddot{u}_\varepsilon(t) - \ddot{u}(t)) \cdot (\dot{u}(t) - \dot{u}_\varepsilon(t)) \, dx \\ + \int_\Omega E\dot{u}(t) : (\sigma_\varepsilon(t) + \varepsilon E\dot{u}_\varepsilon(t) - \sigma(t)) \, dx. \end{aligned}$$

By integrating with respect to time between 0 and t , we obtain

$$\begin{aligned} \mathcal{Q}(e_\varepsilon(t) - e(t)) + \frac{1}{2} \|z_\varepsilon(t) - z(t)\|_2^2 + \frac{1}{2} \|\dot{u}_\varepsilon(t) - \dot{u}(t)\|_2^2 + \varepsilon \int_0^t \int_\Omega |E\dot{u}_\varepsilon(s)|^2 \, dx \, ds \\ \leq \frac{1}{2} \|v_{0,\varepsilon} - v_0\|_2^2 - \int_0^t \int_\Omega \dot{e}(s) : (\sigma_\varepsilon(s) - \sigma(s)) \, dx \, ds - \int_0^t \int_\Omega \dot{z}(s)(z_\varepsilon(s) - z(s)) \, dx \, ds \\ - \int_0^t \int_\Omega (\ddot{u}_\varepsilon(s) - \ddot{u}(s)) \cdot (\dot{u}(s) - \dot{u}_\varepsilon(s)) \, dx \, ds + \int_0^t \int_\Omega E\dot{u}(s) : (\sigma_\varepsilon(s) + \varepsilon E\dot{u}_\varepsilon(s) - \sigma(s)) \, dx \, ds, \end{aligned}$$

where we used the fact that $e_\varepsilon(0) - e(0) = 0$, $z_\varepsilon(0) - z(0) = 0$, and $\dot{u}_\varepsilon(0) - \dot{u}(0) = v_{0,\varepsilon} - v_0$. Since the right-hand-side converges to 0 by (4.8) and (4.10)–(4.12), by (2.1) we deduce (4.20)–(4.23). \square

The energy balance (4.6) can be rewritten between two times $0 \leq t_1 \leq t_2 \leq T$ as

$$\begin{aligned} \mathcal{Q}(e_\varepsilon(t_2)) + \frac{1}{2} \|z_\varepsilon(t_2)\|_2^2 + \int_{t_1}^{t_2} \mathcal{H}_\lambda(\dot{p}_\varepsilon(s), \dot{z}_\varepsilon(s)) ds + \frac{1}{2} \|\dot{u}_\varepsilon(t_2)\|_2^2 + \varepsilon \int_{t_1}^{t_2} \int_\Omega |E\dot{u}_\varepsilon|^2 dx ds \\ = \mathcal{Q}(e_\varepsilon(t_1)) + \frac{1}{2} \|z_\varepsilon(t_1)\|_2^2 + \frac{1}{2} \|\dot{u}_\varepsilon(t_1)\|_2^2 + \int_{t_1}^{t_2} \int_\Omega (\sigma_\varepsilon + \varepsilon E\dot{u}_\varepsilon) : E\dot{w} dx ds \\ + \int_{t_1}^{t_2} \int_\Omega \ddot{u}_\varepsilon \cdot \dot{w} dx ds + \int_{t_1}^{t_2} \int_\Omega f \cdot (\dot{u}_\varepsilon - \dot{w}) dx ds. \end{aligned}$$

Owing to Lemma 4.5, Proposition 2.2, and the lower semicontinuity of the dissipation \mathcal{D}_λ , we deduce

$$\begin{aligned} \mathcal{Q}(e(t_2)) + \frac{1}{2} \|z(t_2)\|_2^2 + \mathcal{D}_\lambda(p, z; [t_1, t_2]) + \frac{1}{2} \|\dot{u}(t_2)\|_2^2 \\ \leq \mathcal{Q}(e(t_1)) + \frac{1}{2} \|z(t_1)\|_2^2 + \frac{1}{2} \|\dot{u}(t_1)\|_2^2 \\ + \int_{t_1}^{t_2} \int_\Omega (\sigma : E\dot{w} + \ddot{u} \cdot \dot{w}) dx ds + \int_{t_1}^{t_2} \int_\Omega f \cdot (\dot{u} - \dot{w}) dx ds. \quad (4.24) \end{aligned}$$

So far we only know that the mapping $t \mapsto p(t)$ has bounded variation into $\mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n})$. The previous energy inequality will enable us to prove the absolute continuity in time of the plastic strain.

Lemma 4.6. *We have $p \in AC([0, T]; \mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n}))$ and $u \in AC([0, T]; BD(\Omega))$.*

Proof. We recall that by (2.3) and (4.16)

$$\mathcal{D}_\lambda(p, z; [t_1, t_2]) \geq \mathcal{H}_\lambda(p(t_2) - p(t_1), z(t_2) - z(t_1)) \geq \alpha_H \|p(t_2) - p(t_1)\|_1 - \frac{\alpha_H}{\sqrt{n}} \|z(t_2) - z(t_1)\|_1.$$

Combining this inequality with (4.24) and the Cauchy-Schwarz inequality yields

$$\begin{aligned} \alpha_H \|p(t_2) - p(t_1)\|_1 &\leq \sup_{t \in [0, T]} \|\sigma(t)\|_2 \|e(t_2) - e(t_1)\|_2 + \sup_{t \in [0, T]} \|\dot{u}(t)\|_2 \|\dot{u}(t_2) - \dot{u}(t_1)\|_2 \\ &+ \sup_{t \in [0, T]} \|\sigma(t)\|_2 \int_{t_1}^{t_2} \|E\dot{w}(s)\|_2 ds + \sup_{t \in [0, T]} \|\ddot{u}(t)\|_2 \int_{t_1}^{t_2} \|\dot{w}(s)\|_2 ds \\ &+ \sup_{t \in [0, T]} (\|\dot{u}(t)\|_2 + \|\dot{w}(t)\|_2) \int_{t_1}^{t_2} \|\dot{f}(s)\|_2 ds \\ &+ \sup_{t \in [0, T]} \|z(t)\|_2 \|z(t_2) - z(t_1)\|_2 + \frac{\alpha_H}{\sqrt{n}} \|z(t_2) - z(t_1)\|_1. \quad (4.25) \end{aligned}$$

This implies that $p \in AC([0, T]; \mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n}))$ and, by the kinematic compatibility and the Poincaré-Korn inequality, that $u \in AC([0, T]; BD(\Omega))$. \square

As a consequence of (4.19) and [7, Lemma 5.5], we infer that for a.e. $t \in [0, T]$,

$$(\dot{u}(t), \dot{e}(t), \dot{p}(t)) \in \mathcal{A}_{\text{dyn}}(\dot{w}(t)). \quad (4.26)$$

We are now in position to derive the flow rule.

Proposition 4.7. *For a.e. $t \in [0, T]$, the distribution $[\sigma(t) : \dot{p}(t)]$ is a measure in $\mathcal{M}(\bar{\Omega})$ and satisfies*

$$H_\lambda(\dot{p}(t), \dot{z}(t)) = [\sigma(t) : \dot{p}(t)] + \xi(t)\dot{z}(t) \quad \text{in } \mathcal{M}(\bar{\Omega}).$$

Proof. According to Lemma 4.6 and Proposition 2.2, we have

$$\mathcal{D}_\lambda(p, z; [t_1, t_2]) = \int_{t_1}^{t_2} \mathcal{H}_\lambda(\dot{p}(s), \dot{z}(s)) ds.$$

Dividing the energy inequality (4.24) by $t_2 - t_1$ (with $t_2 > t_1$) and sending t_1 to $t_2 = t$, we obtain that

$$\begin{aligned} \int_{\Omega} \sigma(t) : \dot{e}(t) dx + \int_{\Omega} z(t) \dot{z}(t) dx + \mathcal{H}_\lambda(\dot{p}(t), \dot{z}(t)) + \int_{\Omega} \dot{u}(t) \cdot \ddot{u}(t) dx \\ \leq \int_{\Omega} (\sigma(t) : E\dot{w}(t) + \ddot{u}(t) \cdot \dot{w}(t)) dx + \int_{\Omega} f(t) \cdot (\dot{u}(t) - \dot{w}(t)) dx \end{aligned}$$

for a.e. $t \in [0, T]$. Using the equation of motion, the previous inequality can be equivalently written as

$$\mathcal{H}_\lambda(\dot{p}(t), \dot{z}(t)) \leq \int_{\Omega} \sigma(t) : (E\dot{w}(t) - \dot{e}(t)) dx + \int_{\Omega} \operatorname{div} \sigma(t) \cdot (\dot{w}(t) - \dot{u}(t)) dx + \int_{\Omega} \xi(t) \dot{z}(t) dx.$$

We notice that by (4.26), for a.e. $t \in [0, T]$ we have $\dot{u}(t) \in BD(\Omega) \cap L^2(\Omega; \mathbb{R}^n)$, $\dot{e}(t) \in L^2(\Omega; \mathbb{M}_{sym}^{n \times n})$, $\dot{p}(t) \in \mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n})$, $E\dot{u}(t) = \dot{e}(t) + \dot{p}(t)$ in Ω , $\dot{p}(t) = (\dot{w}(t) - \dot{u}(t)) \odot \nu \mathcal{H}^{n-1}$ on $\partial\Omega$, while $\sigma(t) \in L^2(\Omega; \mathbb{M}_{sym}^{n \times n})$ and $\operatorname{div} \sigma(t) \in L^2(\Omega; \mathbb{R}^n)$. Thus, we can use the duality introduced in Definition 2.3 and (2.7) to get that

$$\mathcal{H}_\lambda(\dot{p}(t), \dot{z}(t)) \leq \langle \sigma(t), \dot{p}(t) \rangle + \int_{\Omega} \xi(t) \dot{z}(t) dx \quad (4.27)$$

for a.e. $t \in [0, T]$.

Inequality (4.27) implies that $H_\lambda(\dot{p}(t), \dot{z}(t)) \in \mathcal{M}(\Omega')$ for a.e. $t \in [0, T]$. Therefore, by (2.12), we have that

$$\begin{aligned} \int_{\Omega'} \varphi d[H_\lambda(\dot{p}(t), \dot{z}(t))] = \sup \left\{ \langle [\tau : \dot{p}(t)], \varphi \rangle + \int_{\Omega} \eta \dot{z}(t) \varphi dx : (\tau, \eta) \in L^2(\Omega; \mathbb{M}_{sym}^{n \times n}) \times L^2(\Omega) \right. \\ \left. \text{with } \operatorname{div} \tau \in L^2(\Omega; \mathbb{R}^n) \text{ and } (\tau(x), \eta(x)) \in K_\lambda \text{ for a.e. } x \in \Omega \right\} \end{aligned}$$

for every $\varphi \in \mathcal{C}_c^\infty(\Omega')$, $\varphi \geq 0$ and a.e. $t \in [0, T]$. In particular,

$$\int_{\Omega'} \varphi d[H_\lambda(\dot{p}(t), \dot{z}(t))] \geq \langle [\sigma(t) : \dot{p}(t)], \varphi \rangle + \int_{\Omega} \xi(t) \dot{z}(t) \varphi dx$$

for every $\varphi \in \mathcal{C}_c^\infty(\Omega')$, $\varphi \geq 0$ and a.e. $t \in [0, T]$, which implies that

$$H_\lambda(\dot{p}(t), \dot{z}(t)) \geq [\sigma(t) : \dot{p}(t)] + \xi(t) \dot{z}(t)$$

for a.e. $t \in [0, T]$, where the inequality above is intended in the sense of distributions on Ω' . In other words, $H_\lambda(\dot{p}(t), \dot{z}(t)) - [\sigma(t) : \dot{p}(t)] - \xi(t) \dot{z}(t)$ is a non negative distribution, hence a Radon measure with zero total variation by (4.27). Therefore, for a.e. $t \in [0, T]$

$$[\sigma(t) : \dot{p}(t)] \in \mathcal{M}(\Omega')$$

and

$$H_\lambda(\dot{p}(t), \dot{z}(t)) = [\sigma(t) : \dot{p}(t)] + \xi(t) \dot{z}(t) \quad \text{in } \mathcal{M}(\Omega').$$

Thanks to our choice of the extensions this last relation implies (4.4) and completes the proof of the proposition. \square

4.3. Uniqueness of the solution. The proof of Theorem 4.1 will be complete once the uniqueness is proved. Let us consider two solutions $(u_1, e_1, p_1, z_1, \sigma_1, \xi_1)$ and $(u_2, e_2, p_2, z_2, \sigma_2, \xi_2)$. Subtracting the equations of motions leads to

$$\ddot{u}_1 - \ddot{u}_2 - \operatorname{div}(\sigma_1 - \sigma_2) = 0 \quad \text{in } L^2(0, T; L^2(\Omega; \mathbb{R}^n)),$$

and since $\dot{u}_1 - \dot{u}_2 \in L^2(0, T; L^2(\Omega; \mathbb{R}^n))$, we infer that

$$\int_0^t \int_{\Omega} (\ddot{u}_1 - \ddot{u}_2) \cdot (\dot{u}_1 - \dot{u}_2) \, dx \, ds - \int_0^t \int_{\Omega} \operatorname{div}(\sigma_1 - \sigma_2) \cdot (\dot{u}_1 - \dot{u}_2) \, dx \, ds = 0. \quad (4.28)$$

Since $\ddot{u}_1 - \ddot{u}_2 \in L^2(0, T; L^2(\Omega; \mathbb{R}^n))$, we get that

$$\int_0^t \int_{\Omega} (\ddot{u}_1(s) - \ddot{u}_2(s)) \cdot (\dot{u}_1(s) - \dot{u}_2(s)) \, dx \, ds = \frac{\|\dot{u}_1(t) - \dot{u}_2(t)\|_2^2}{2} \quad (4.29)$$

where we used that $\dot{u}_1(0) = \dot{u}_2(0) = v_0$. On the other hand, by the stress-strain duality (see Definition 2.3 and (2.7)), we get that

$$\begin{aligned} & - \int_0^t \int_{\Omega} \operatorname{div}(\sigma_1(s) - \sigma_2(s)) \cdot (\dot{u}_1(s) - \dot{u}_2(s)) \, dx \, ds \\ &= \int_0^t \int_{\Omega} (\sigma_1(s) - \sigma_2(s)) : (\dot{e}_1(s) - \dot{e}_2(s)) \, dx \, dt + \int_0^t \langle \sigma_1(s) - \sigma_2(s), \dot{p}_1(s) - \dot{p}_2(s) \rangle \, ds \\ &= \mathcal{Q}(e_1(t) - e_2(t)) + \int_0^t \langle \sigma_1(s) - \sigma_2(s), \dot{p}_1(s) - \dot{p}_2(s) \rangle \, ds, \end{aligned} \quad (4.30)$$

since $e_1(0) = e_2(0) = e_0$. We now estimate the last integral. We first note that, for $i = 1, 2$, $(\sigma_i(s), \xi_i(s)) \in L^2(\Omega; \mathbb{M}_{sym}^{n \times n}) \times L^2(\Omega)$, $\operatorname{div} \sigma_i(s) \in L^2(\Omega; \mathbb{R}^n)$, and $(\sigma_i(s), \xi_i(s)) \in K_{\lambda}$ a.e. in Ω for all $s \in [0, T]$. Therefore, by the flow rule (4.4), the duality formula (2.13) and the kinematic compatibility for the rates (4.26), we get that for a.e. $s \in [0, T]$,

$$\begin{aligned} \langle \sigma_1(s), \dot{p}_1(s) \rangle + \int_{\Omega} \xi_1(s) \dot{z}_1(s) \, dx &= \mathcal{H}_{\lambda}(\dot{p}_1(s), \dot{z}_1(s)) \geq \langle \sigma_2(s), \dot{p}_1(s) \rangle + \int_{\Omega} \xi_2(s) \dot{z}_1(s) \, dx, \\ \langle \sigma_2(s), \dot{p}_2(s) \rangle + \int_{\Omega} \xi_2(s) \dot{z}_2(s) \, dx &= \mathcal{H}_{\lambda}(\dot{p}_2(s), \dot{z}_2(s)) \geq \langle \sigma_1(s), \dot{p}_2(s) \rangle + \int_{\Omega} \xi_1(s) \dot{z}_2(s) \, dx. \end{aligned}$$

Summing up both previous inequalities and integrating in time yields

$$\int_0^t \langle \sigma_1(s) - \sigma_2(s), \dot{p}_1(s) - \dot{p}_2(s) \rangle \, ds \geq \int_0^t \int_{\Omega} (z_1 - z_2)(\dot{z}_1 - \dot{z}_2) \, dx \, ds = \frac{\|z_1(t) - z_2(t)\|_2^2}{2}, \quad (4.31)$$

since $z_1(0) = z_2(0) = z_0$. Gathering (4.28)–(4.31) yields $e_1 = e_2$ (hence $\sigma_1 = \sigma_2$), $z_1 = z_2$ (hence $\xi_1 = \xi_2$), and $\dot{u}_1 = \dot{u}_2$. But since $u_1(0) = u_2(0) = u_0$, we deduce that $u_1 = u_2$ and finally that $p_1 = p_2$.

Remark 4.8. Thanks to the uniqueness of the solution, there is actually no need to extract subsequences in all weak and strong convergences obtained so far.

5. CONVERGENCE OF THE DYNAMIC CAP MODEL

In this section we characterize the asymptotic behaviour of the elasto-plastic dynamic evolutions studied in the previous section when the cap is sent to infinity. In that way we recover a solution of (1.2), namely of a model of perfect elasto-plasticity with a pressure-sensitive yield criterion.

As before, we consider a body load $f \in AC([0, T]; L^2(\Omega; \mathbb{R}^n))$ and a boundary displacement which is the trace on $\partial\Omega \times (0, T)$ of a function $w \in H^2([0, T]; H^1(\Omega; \mathbb{R}^n)) \cap H^3([0, T]; L^2(\Omega; \mathbb{R}^n))$. We also consider initial data $(u_0, e_0, p_0) \in \mathcal{A}_{\text{dyn}}(w(0))$ and $v_0 \in H^1(\Omega; \mathbb{R}^n)$ satisfying

$$v_0 = \dot{w}(0) \quad \mathcal{H}^{n-1}\text{-a.e. on } \partial\Omega,$$

and

$$\sigma_0 := \mathbb{C}e_0 \in K, \quad -\operatorname{div}\sigma_0 = f(0) \quad \text{a.e. in } \Omega.$$

In order to apply Theorem 4.1, we need to fix also an initial datum for the hardening cap variable. To this aim, let us define

$$\xi_0 := (\sigma_0)_m - |(\sigma_0)_m| \quad \text{and} \quad z_0 := -\xi_0.$$

With this definition, and according to (2.2), it is immediate to check that $\xi_0, z_0 \in L^2(\Omega)$ and $(\sigma_0, \xi_0) \in K_1 \subset K_\lambda$ a.e. in Ω , for all $\lambda \geq 1$.

For every $\lambda \geq 1$, let $u_\lambda, e_\lambda, \sigma_\lambda, p_\lambda, z_\lambda, \xi_\lambda$ be the solution to (4.1)–(4.4) constructed in Theorem 4.1 with initial data (u_0, e_0, p_0, z_0) and v_0 .

Theorem 5.1. *There exist*

$$\begin{cases} u \in AC([0, T]; BD(\Omega)) \cap W^{2, \infty}([0, T]; L^2(\Omega; \mathbb{R}^n)), \\ \sigma, e \in W^{1, \infty}([0, T]; L^2(\Omega; \mathbb{M}_{sym}^{n \times n})), \\ p \in AC([0, T]; \mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n})), \end{cases}$$

such that for all $t \in [0, T]$,

$$u_\lambda(t) \rightharpoonup u(t) \text{ weakly* in } BD(\Omega), \quad (5.1)$$

$$e_\lambda(t) \rightharpoonup e(t) \text{ weakly in } L^2(\Omega; \mathbb{M}_{sym}^{n \times n}), \quad (5.2)$$

$$p_\lambda(t) \rightharpoonup p(t) \text{ weakly* in } \mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n}), \quad (5.3)$$

and

$$u_\lambda \rightarrow u \text{ strongly in } W^{1, \infty}([0, T]; L^2(\Omega; \mathbb{R}^n)), \quad (5.4)$$

$$e_\lambda \rightarrow e \text{ strongly in } L^\infty(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n})), \quad (5.5)$$

as $\lambda \rightarrow +\infty$. For every $t \in [0, T]$ we have

$$\begin{cases} Eu(t) = e(t) + p(t) \text{ in } \Omega, & p(t) = (w(t) - u(t)) \odot \nu \mathcal{H}^{n-1} \text{ on } \partial\Omega, \\ \sigma(t) = \mathbb{C}e(t), \\ \sigma(t) \in K \text{ a.e. in } \Omega, \end{cases} \quad (5.6)$$

Moreover,

$$\begin{cases} \ddot{u} - \operatorname{div}\sigma = f \quad \text{a.e. in } \Omega \times (0, T), \\ (u(0), e(0), p(0)) = (u_0, e_0, p_0), \quad \dot{u}(0) = v_0, \end{cases} \quad (5.7)$$

and for a.e. $t \in [0, T]$ the distribution $[\sigma(t) : \dot{p}(t)]$ is a measure in $\mathcal{M}(\bar{\Omega})$ satisfying

$$H(\dot{p}(t)) = [\sigma(t) : \dot{p}(t)] \text{ in } \mathcal{M}(\bar{\Omega}). \quad (5.8)$$

Furthermore, (u, e, σ, p) is the unique solution of (5.6)–(5.8).

Proof. The proof of the theorem is split into several steps.

Step 1: A priori estimates and compactness. From (4.5), it follows that

$$\|\ddot{u}_\lambda\|_{L^\infty(0, T; L^2(\Omega; \mathbb{R}^n))} + \|\dot{e}_\lambda\|_{L^\infty(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))} + \|\dot{z}_\lambda\|_{L^\infty(0, T; L^2(\Omega))} \leq C_1, \quad (5.9)$$

for some positive constant $C_1 > 0$ independent of λ . On the other hand, by the energy inequality (4.24) we have

$$\begin{aligned} & \frac{\alpha_C}{2} \|e_\lambda(t)\|_2^2 + \frac{1}{2} \|z_\lambda(t)\|_2^2 + \frac{1}{2} \|\dot{u}_\lambda(t)\|_2^2 \\ & \leq \frac{\beta_C}{2} \|e_0\|_2^2 + \frac{1}{2} \|z_0\|_2^2 + \frac{1}{2} \|v_0\|_2^2 + \|\sigma_\lambda\|_{L^\infty(0,T;L^2(\Omega;\mathbb{M}_{sym}^{n \times n}))} \|E\dot{w}\|_{L^1(0,T;L^2(\Omega;\mathbb{M}_{sym}^{n \times n}))} \\ & \quad + \|\ddot{u}_\lambda\|_{L^\infty(0,T;L^2(\Omega;\mathbb{R}^n))} \|\dot{w}\|_{L^1(0,T;L^2(\Omega;\mathbb{R}^n))} \\ & \quad + \|f\|_{L^1(0,T;L^2(\Omega;\mathbb{R}^n))} (\|\dot{u}_\lambda\|_{L^\infty(0,T;L^2(\Omega;\mathbb{R}^n))} + \|\dot{w}\|_{L^\infty(0,T;L^2(\Omega;\mathbb{R}^n))}) \end{aligned}$$

for every $t \in [0, T]$. This, together with (5.9), implies that

$$\|e_\lambda\|_{L^\infty(0,T;L^2(\Omega;\mathbb{M}_{sym}^{n \times n}))} + \|z_\lambda\|_{L^\infty(0,T;L^2(\Omega))} + \|\dot{u}_\lambda\|_{L^\infty(0,T;L^2(\Omega;\mathbb{R}^n))} \leq C_2,$$

for some constant $C_2 > 0$ independent of λ . Moreover, combining (4.25) with the previous estimates yields

$$\|p_\lambda(t_2) - p_\lambda(t_1)\|_1 \leq C_3 \int_{t_1}^{t_2} \left(1 + \|\dot{w}(s)\|_{H^1(\Omega;\mathbb{R}^n)} + \|\dot{f}(s)\|_2\right) ds$$

for every $0 \leq t_1 \leq t_2 \leq T$, and some constant $C_3 > 0$ independent of λ .

From the previous bounds, we deduce the existence of functions $v \in W^{1,\infty}([0, T]; L^2(\Omega; \mathbb{R}^n))$, $e \in W^{1,\infty}([0, T]; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))$, and $z \in W^{1,\infty}([0, T]; L^2(\Omega))$ such that, up to subsequences,

$$\begin{aligned} \dot{u}_\lambda & \rightharpoonup v \text{ weakly* in } L^\infty(0, T; L^2(\Omega; \mathbb{R}^n)), \\ \ddot{u}_\lambda & \rightharpoonup \dot{v} \text{ weakly* in } L^\infty(0, T; L^2(\Omega; \mathbb{R}^n)), \end{aligned} \tag{5.10}$$

$$e_\lambda \rightharpoonup e \text{ weakly* in } L^\infty(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n})), \tag{5.11}$$

$$\begin{aligned} \dot{e}_\lambda & \rightharpoonup \dot{e} \text{ weakly* in } L^\infty(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n})), \\ z_\lambda & \rightharpoonup z \text{ weakly* in } L^\infty(0, T; L^2(\Omega)), \end{aligned} \tag{5.12}$$

$$\dot{z}_\lambda \rightharpoonup \dot{z} \text{ weakly* in } L^\infty(0, T; L^2(\Omega)).$$

By Ascoli-Arzelà Theorem we also have that

$$\begin{aligned} \dot{u}_\lambda(t) & \rightharpoonup v(t) \text{ weakly in } L^2(\Omega; \mathbb{R}^n), \\ e_\lambda(t) & \rightharpoonup e(t) \text{ weakly in } L^2(\Omega; \mathbb{M}_{sym}^{n \times n}), \\ z_\lambda(t) & \rightharpoonup z(t) \text{ weakly in } L^2(\Omega) \end{aligned} \tag{5.13}$$

for every $t \in [0, T]$. Moreover, (4.3) implies that

$$\dot{z}(t) \geq 0 \text{ a.e. in } \Omega \tag{5.14}$$

for a.e. $t \in [0, T]$. Finally, again by Ascoli-Arzelà Theorem we deduce the existence of a function $p \in AC([0, T]; \mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n}))$ such that (5.3) is satisfied for every $t \in [0, T]$.

Step 2: Kinematic compatibility, equation of motion, and initial condition. Arguing as in the proof of Theorem 4.1, one can show the existence of $u \in AC([0, T]; BD(\Omega)) \cap W^{2,\infty}([0, T]; L^2(\Omega; \mathbb{R}^n))$ such that (5.1) is satisfied, $v(t) = \dot{u}(t)$ and

$$Eu(t) = e(t) + p(t) \text{ in } \Omega, \quad p(t) = (w(t) - u(t)) \odot \nu \mathcal{H}^{n-1} \text{ on } \partial\Omega$$

for every $t \in [0, T]$, so that the kinematic compatibility holds. Moreover, according to [7, Lemma 5.5], we get, for a.e. $t \in [0, T]$,

$$E\dot{u}(t) = \dot{e}(t) + \dot{p}(t) \text{ in } \Omega, \quad \dot{p}(t) = (\dot{w}(t) - \dot{u}(t)) \odot \nu \mathcal{H}^{n-1} \text{ on } \partial\Omega. \tag{5.15}$$

Clearly the equation of motion and the initial conditions $u(0) = u_0$, $e(0) = e_0$, $p(0) = p_0$, $z(0) = z_0$, and $\dot{u}(0) = v_0$ are satisfied.

Step 3: Stress constraint. For what concerns the stress constraint, we set $\sigma(t) := \mathbb{C}e(t)$ and $\xi(t) := -z(t)$. From the inclusion $(\sigma_\lambda(t), \xi_\lambda(t)) \in K_\lambda$ a.e. in Ω for every $\lambda \geq 1$, it follows that $(\sigma_\lambda(t), \xi_\lambda(t)) \in K \times \mathbb{R}^-$ a.e. in Ω for every $\lambda \geq 1$. By (5.2) and (5.13) we deduce that $(\sigma(t), \xi(t)) \in K \times \mathbb{R}^-$ a.e. in Ω , which implies that $\sigma(t) \in K$ a.e. in Ω for every $t \in [0, T]$.

Step 4: Strong convergences. We now prove the strong convergences (5.4), (5.5), together with

$$z_\lambda \rightarrow z \text{ strongly in } L^\infty(0, T; L^2(\Omega)). \quad (5.16)$$

For every $\lambda \geq 1$ we define

$$\zeta_\lambda := z - \frac{1}{\lambda}(\sigma_m - |\sigma_m|).$$

Using the fact that $z \in W^{1,\infty}([0, T]; L^2(\Omega))$ and $\sigma \in W^{1,\infty}([0, T]; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))$, we get that $\zeta_\lambda \in W^{1,\infty}([0, T]; L^2(\Omega))$ and

$$\zeta_\lambda \rightarrow z \text{ strongly in } W^{1,\infty}([0, T]; L^2(\Omega)), \quad (5.17)$$

as $\lambda \rightarrow \infty$. Moreover, according to (2.2), $(\sigma(t), -\zeta_\lambda(t)) \in K_\lambda$ a.e. in Ω . Since $\text{div} \sigma(t) \in L^2(\Omega; \mathbb{R}^n)$, by integration of (4.4) and the duality formula (2.13), we have

$$\langle \sigma(t), \dot{p}_\lambda(t) \rangle - \int_\Omega \zeta_\lambda(t) \dot{z}_\lambda(t) dx \leq \langle \sigma_\lambda(t), \dot{p}_\lambda(t) \rangle + \int_\Omega \xi_\lambda(t) \dot{z}_\lambda(t) dx$$

for a.e. $t \in [0, T]$. Using the definition of the stress-strain duality (2.7), the kinematic compatibility for the rate (4.26), and the equation of motion this can be rewritten as

$$\begin{aligned} \int_\Omega \dot{e}_\lambda(t) : (\sigma_\lambda(t) - \sigma(t)) dx + \int_\Omega \dot{z}_\lambda(t)(z_\lambda(t) - \zeta_\lambda(t)) dx + \int_\Omega \dot{u}_\lambda(t) \cdot (\ddot{u}_\lambda(t) - \ddot{u}(t)) dx \\ \leq \int_\Omega \dot{w}(t) \cdot (\ddot{u}_\lambda(t) - \ddot{u}(t)) dx + \int_\Omega E \dot{w}(t) : (\sigma_\lambda(t) - \sigma(t)) dx. \end{aligned}$$

By integrating with respect to time we obtain

$$\begin{aligned} \mathcal{Q}(e_\lambda(t) - e(t)) + \frac{1}{2} \|z_\lambda(t) - \zeta_\lambda(t)\|_2^2 + \frac{1}{2} \|\dot{u}_\lambda(t) - \dot{u}(t)\|_2^2 \\ \leq \frac{1}{2} \|z(0) - \zeta_\lambda(0)\|_2^2 - \int_0^t \int_\Omega \dot{e}(s) : (\sigma_\lambda(s) - \sigma(s)) dx ds - \int_0^t \int_\Omega \dot{\zeta}_\lambda(s)(z_\lambda(s) - \zeta_\lambda(s)) dx ds \\ + \int_0^t \int_\Omega (\dot{w}(s) - \dot{u}(s)) \cdot (\ddot{u}_\lambda(s) - \ddot{u}(s)) dx ds + \int_0^t \int_\Omega E \dot{w}(s) : (\sigma_\lambda(s) - \sigma(s)) dx ds, \end{aligned}$$

where we used that $e_\lambda(0) = e(0)$, $z_\lambda(0) = z(0)$, and $\dot{u}_\lambda(0) = \dot{u}(0)$. As $\lambda \rightarrow \infty$, the right-hand side converges to 0 by (5.10)–(5.12) and (5.17). Owing to (2.1), we deduce (5.4), (5.5), and

$$z_\lambda - \zeta_\lambda \rightarrow 0 \text{ strongly in } L^\infty(0, T; L^2(\Omega)),$$

which, together with (5.17), implies (5.16).

Step 5: Flow rule. Owing to the strong convergences proved in the previous step, we are now in position to pass to the limit into the energy inequality (4.24). We first observe that for every $0 \leq t_1 \leq t_2 \leq T$ and every $\lambda_1 \geq 1$ we have by lower semicontinuity of the total dissipation and

Proposition 2.2

$$\begin{aligned} \int_{t_1}^{t_2} \mathcal{H}_{\lambda_1}(\dot{p}(s), \dot{z}(s)) ds &= \mathcal{D}_{\lambda_1}(p, z, [t_1, t_2]) \\ &\leq \liminf_{\lambda \rightarrow \infty} \mathcal{D}_{\lambda_1}(p_\lambda, z_\lambda, [t_1, t_2]) = \liminf_{\lambda \rightarrow \infty} \int_{t_1}^{t_2} \mathcal{H}_{\lambda_1}(\dot{p}_\lambda(s), \dot{z}_\lambda(s)) ds \\ &\leq \liminf_{\lambda \rightarrow \infty} \int_{t_1}^{t_2} \mathcal{H}_\lambda(\dot{p}_\lambda(s), \dot{z}_\lambda(s)) ds. \end{aligned}$$

By monotone convergence, letting $\lambda_1 \rightarrow \infty$ and applying Lemma 2.1 yield

$$\int_{t_1}^{t_2} \mathcal{H}(\dot{p}(s)) ds \leq \liminf_{\lambda \rightarrow \infty} \int_{t_1}^{t_2} \mathcal{H}_\lambda(\dot{p}_\lambda(s), \dot{z}_\lambda(s)) ds,$$

where we used (5.14). Thus passing to the limit in (4.24) yields

$$\begin{aligned} \mathcal{Q}(e(t_2)) + \frac{1}{2} \|z(t_2)\|_2^2 + \int_{t_1}^{t_2} \mathcal{H}(\dot{p}(s)) ds + \frac{1}{2} \|\dot{u}(t_2)\|_2^2 \\ \leq \mathcal{Q}(e(t_1)) + \frac{1}{2} \|z(t_1)\|_2^2 + \frac{1}{2} \|\dot{u}(t_1)\|_2^2 \\ + \int_{t_1}^{t_2} \int_{\Omega} (\sigma : E\dot{w} + \ddot{u} \cdot \dot{w}) dx ds + \int_{t_1}^{t_2} f \cdot (\dot{u} - \dot{w}) dx ds. \end{aligned} \quad (5.18)$$

Since $\dot{z}(t) \geq 0$ a.e. in Ω , we have $z(t_2) \geq z(t_1) \geq z_0 \geq 0$ a.e. in Ω ; thus,

$$\begin{aligned} \mathcal{Q}(e(t_2)) + \int_{t_1}^{t_2} \mathcal{H}(\dot{p}(s)) ds + \frac{1}{2} \|\dot{u}(t_2)\|_2^2 \\ \leq \mathcal{Q}(e(t_1)) + \frac{1}{2} \|\dot{u}(t_1)\|_2^2 + \int_{t_1}^{t_2} \int_{\Omega} (\sigma : E\dot{w} + \ddot{u} \cdot \dot{w}) dx ds + \int_{t_1}^{t_2} f \cdot (\dot{u} - \dot{w}) dx ds. \end{aligned}$$

By differentiation with respect to time, the duality formula (2.7) and the kinematic compatibility for the rates (5.15), this is equivalent to

$$\mathcal{H}(\dot{p}(t)) \leq \langle \sigma(t), \dot{p}(t) \rangle$$

for a.e. $t \in [0, T]$. Using next (2.10), and arguing as in the proof of Theorem 4.1, one can prove that this last inequality yields, in turn, (5.8).

Step 5: Uniqueness. The proof of the uniqueness of the solution is identical to that of Theorem 4.1, and rest on the stress-strain duality (2.7) as well as on the duality formula (2.11). In particular, the uniqueness ensures that there is no need to extract subsequences to get the previous convergences. \square

Remark 5.2. Integrating the flow rule (5.8), and using the duality formula (2.7), together with the kinematic compatibility for the rates (5.15), one can show that the energy inequality is actually an equality:

$$\begin{aligned} \mathcal{Q}(e(t)) + \int_0^t \mathcal{H}(\dot{p}(s)) ds + \frac{1}{2} \|\dot{u}(t)\|_2^2 &= \mathcal{Q}(e(0)) + \frac{1}{2} \|v_0\|_2^2 \\ &+ \int_0^t \int_{\Omega} (\sigma : E\dot{w} + \ddot{u} \cdot \dot{w}) dx ds + \int_0^t \int_{\Omega} f \cdot (\dot{u} - \dot{w}) dx ds. \end{aligned}$$

As a consequence, since $t \mapsto z(t)$ is non-decreasing, comparing with (5.18), we deduce that $z(t) = z_0$ (and thus $\xi(t) = \xi_0$) for all $t \in [0, T]$.

6. THE QUASI-STATIC CASE

In the dynamical elasto-plastic model the kinetic energy gives a natural $L^2(\Omega; \mathbb{R}^n)$ bound on the velocity field, which is crucial in order to define the duality between the stress and the plastic strain rate. Unfortunately, in the quasi-static case, the velocity only belongs, in general, to $BD(\Omega)$ and thus, it is at most in $L^{n/(n-1)}(\Omega; \mathbb{R}^n)$. Consequently, without any further regularity result at our disposal, the stress-strain duality might not be well defined, except of course in the two dimensional setting. This is clearly an obstacle in order to express the flow rule in a measure theoretic sense as we did in (4.4) for the cap model, and in (5.8) for the perfect elasto-plastic model. However, in higher dimension it is possible to give a weak sense to the flow rule by means of an energy equality.

6.1. Quasi-static elasto-plastic cap model. Using a variational evolution approach similar to that of [7], we can show an existence result for solutions of a quasi-static elasto-plastic cap model. In this context, since the kinetic energy is no longer controlled, given a boundary displacement $\hat{w} \in H^1(\Omega; \mathbb{R}^n)$, the space of kinematically admissible fields is defined by

$$\mathcal{A}_{\text{qs}}(\hat{w}) := \left\{ (v, \eta, q) \in BD(\Omega) \times L^2(\Omega; \mathbb{M}_{\text{sym}}^{n \times n}) \times \mathcal{M}(\bar{\Omega}; \mathbb{M}_{\text{sym}}^{n \times n}) : \right. \\ \left. Ev = \eta + q \text{ in } \Omega, \quad q = (\hat{w} - v) \odot \nu \mathcal{H}^{n-1} \text{ on } \partial\Omega \right\}.$$

Let us fix $\lambda \geq 1$ and consider a body load $f \in AC([0, T]; L^n(\Omega; \mathbb{R}^n))$ and a boundary displacement which is the trace on $\partial\Omega \times (0, T)$ of a function $w \in AC([0, T]; H^1(\Omega; \mathbb{R}^n))$. We also consider an initial datum $(u_0, e_0, p_0, z_0) \in \mathcal{A}_{\text{qs}}(w(0)) \times L^2(\Omega)$ satisfying the stability condition

$$\mathcal{Q}(e_0) + \frac{1}{2} \|z_0\|_2^2 - \int_{\Omega} f(0) \cdot u_0 \, dx \leq \mathcal{Q}(\eta) + \mathcal{H}_{\lambda}(q - p_0, \zeta - z_0) + \frac{1}{2} \|\zeta\|_2^2 - \int_{\Omega} f(0) \cdot v \, dx,$$

for any $(v, \eta, q, \zeta) \in \mathcal{A}_{\text{qs}}(w(0)) \times L^2(\Omega)$, and we define $(\sigma_0, \xi_0) := (\mathbb{C}e_0, -z_0)$.

Contrary to the dynamical case, we need also to assume the following safe-load condition: there exist $\chi \in AC([0, T]; L^n(\Omega; \mathbb{M}_{\text{sym}}^{n \times n}))$, $\vartheta \in AC([0, T]; L^2(\Omega))$, and a constant $\alpha_0 > 0$ such that for every $t \in [0, T]$

$$-\text{div} \chi(t) = f(t) \quad \text{a.e. in } \Omega \tag{6.1}$$

and

$$(\chi(t) + \tau, \vartheta(t)) \in K_{\lambda} \quad \text{a.e. in } \Omega \tag{6.2}$$

for every $\tau \in \mathbb{M}_{\text{sym}}^{n \times n}$ with $|\tau| \leq \alpha_0$.

As explained in the introduction, the validity of the safe-load condition ensures a control on the plastic strain (rate) from a control on the dissipation functional. Indeed, the following result establishes a coercivity property of the functional $p \mapsto \mathcal{H}_{\lambda}(p, z) - \langle \chi(t), p \rangle$.

Proposition 6.1. *Let $\hat{w} \in H^1(\Omega; \mathbb{R}^n)$ and let $(u, e, p, z) \in \mathcal{A}_{\text{qs}}(\hat{w}) \times L^2(\Omega)$. Then for every $t \in [0, T]$ the following coercivity estimate holds:*

$$\mathcal{H}_{\lambda}(p, z) - \langle \chi(t), p \rangle \geq \alpha_0 \|p\|_1 - \alpha_1 \|z\|_2,$$

where $\alpha_1 := \|\vartheta\|_{L^\infty(0, T; L^2(\Omega))}$.

Proof. We notice that the duality $\langle \chi(t), p \rangle$ is well defined owing to Remark 2.6. Moreover, we can assume $\mathcal{H}_{\lambda}(p, z) < \infty$, otherwise the result is trivial. By the duality formula (2.13) and Remark 2.6 we have

$$\mathcal{H}_{\lambda}(p, z) - \langle \chi(t), p \rangle = \sup \left\{ \langle \sigma - \chi(t), p \rangle + \int_{\Omega} \xi z \, dx : (\sigma, \xi) \in L^n(\Omega; \mathbb{M}_{\text{sym}}^{n \times n}) \times L^2(\Omega) \text{ with} \right. \\ \left. \text{div} \sigma \in L^n(\Omega; \mathbb{R}^n) \text{ and } (\sigma(x), \xi(x)) \in K_{\lambda} \text{ a.e. in } \Omega \right\}.$$

Using (2.9) and (6.2), this implies that

$$\begin{aligned} \mathcal{H}_\lambda(p, z) - \langle \chi(t), p \rangle &\geq \sup \left\{ \int_{\bar{\Omega}} \tau : dp + \int_{\Omega} \vartheta(t) z \, dx : \tau \in C^\infty(\bar{\Omega}; \mathbb{R}^n) \text{ and } |\tau| \leq \alpha_0 \text{ in } \bar{\Omega} \right\} \\ &= \alpha_0 \|p\|_1 + \int_{\Omega} \vartheta(t) z \, dx, \end{aligned}$$

from which the thesis immediately follows. \square

The following result concerns the optimality conditions of a suitable minimum problem arising in the definition of the incremental evolution.

Lemma 6.2. *Let $\hat{w} \in H^1(\Omega; \mathbb{R}^n)$, $f \in L^n(\Omega; \mathbb{R}^n)$, $(u, e, p, z) \in \mathcal{A}_{\text{qs}}(\hat{w}) \times L^2(\Omega)$, and $(\sigma, \xi) := (\mathbb{C}e, -z)$. Then the following conditions are equivalent:*

(a) (u, e, p, z) is a solution of

$$\min_{(v, \eta, q, \zeta) \in \mathcal{A}_{\text{qs}}(\hat{w}) \times L^2(\Omega)} \left\{ \mathcal{Q}(\eta) + \frac{1}{2} \|\zeta\|_2^2 + \mathcal{H}_\lambda(q - p, \zeta - z) - \int_{\Omega} f \cdot v \, dx \right\};$$

(b) (σ, ξ) satisfies

$$- \int_{\Omega} \sigma : \eta \, dx + \int_{\Omega} \xi \zeta \, dx + \int_{\Omega} f \cdot v \, dx \leq \mathcal{H}_\lambda(q, \zeta)$$

for every $(v, \eta, q, \zeta) \in \mathcal{A}_{\text{qs}}(0) \times L^2(\Omega)$.

If (a) or (b) holds, then the following conditions are satisfied:

(c) $(\sigma, \xi) \in K_\lambda$ and $-\text{div} \sigma = f$ a.e. in Ω .

If, in addition, $\sigma \in L^n(\Omega; \mathbb{M}_{\text{sym}}^{n \times n})$ or $u \in L^2(\Omega; \mathbb{R}^n)$, then (c) is equivalent to (a) and (b).

Proof. The proof is an immediate adaptation of [7, Theorem 3.6], once we notice that if $\sigma \in L^n(\Omega; \mathbb{M}_{\text{sym}}^{n \times n})$ or $u \in L^2(\Omega; \mathbb{R}^n)$ the stress-strain duality is well defined by Remark 2.6. \square

We are now in a position to prove the first main result of this section.

Theorem 6.3. *There exist*

$$\begin{cases} u \in AC([0, T]; BD(\Omega)), \\ \sigma, e \in AC([0, T]; L^2(\Omega; \mathbb{M}_{\text{sym}}^{n \times n})), \\ p \in AC([0, T]; \mathcal{M}(\bar{\Omega}; \mathbb{M}_{\text{sym}}^{n \times n})), \\ \xi, z \in AC([0, T]; L^2(\Omega)), \end{cases}$$

with $(u(0), e(0), p(0), z(0)) = (u_0, e_0, p_0, z_0)$, satisfying the following properties: for all $t \in [0, T]$

$$\begin{cases} Eu(t) = e(t) + p(t) \text{ in } \Omega, & p(t) = (w(t) - u(t)) \odot \nu \mathcal{H}^{n-1} \text{ on } \partial\Omega, \\ \sigma(t) = \mathbb{C}e(t), & \xi(t) = -z(t), \end{cases} \quad (6.3)$$

and

$$\begin{cases} -\text{div} \sigma(t) = f(t) \text{ a.e. in } \Omega, \\ (\sigma(t), \xi(t)) \in K_\lambda. \end{cases}$$

Moreover, for a.e. $t \in [0, T]$

$$\dot{z}(t) \geq 0 \quad \text{a.e. in } \Omega, \quad (6.4)$$

and the following energy equality holds for all $t \in [0, T]$,

$$\begin{aligned} \mathcal{Q}(e(t)) + \frac{1}{2} \|z(t)\|_2^2 + \int_0^t \mathcal{H}_\lambda(\dot{p}(s), \dot{z}(s)) ds &= \mathcal{Q}(e_0) + \frac{1}{2} \|z_0\|_2^2 \\ &+ \int_0^t \int_\Omega \sigma(s) : E\dot{w}(s) dx ds - \int_0^t \int_\Omega f(s) \cdot (\dot{w}(s) - \dot{u}(s)) dx ds. \end{aligned} \quad (6.5)$$

Proof of Theorem 6.3. Let us sketch the proof of Theorem 6.3. As in [7], we introduce a time discretization

$$0 = t_k^0 < t_k^1 < \dots < t_k^{N_k} = T, \quad \text{with} \quad \delta_k := \sup_{1 \leq i \leq N_k} (t_k^i - t_k^{i-1}) \rightarrow 0$$

of the time interval $[0, T]$. For each $i = 0$, we set $(u_k^0, e_k^0, p_k^0, z_k^0) := (u_0, e_0, p_0, z_0)$ and for all $i \in \{1, \dots, N_k\}$ we define $(u_k^i, e_k^i, p_k^i, z_k^i) \in \mathcal{A}_{\text{qs}}(w(t_k^i)) \times L^2(\Omega)$ as a solution of the minimum problem

$$\min_{(v, \eta, q, \zeta) \in \mathcal{A}_{\text{qs}}(w(t_k^i)) \times L^2(\Omega)} \left\{ \mathcal{Q}(\eta) + \frac{1}{2} \|\zeta\|_2^2 + \mathcal{H}_\lambda(q - p_k^{i-1}, \zeta - z_k^{i-1}) - \int_\Omega f(t_k^i) \cdot v dx \right\}.$$

Such a solution exists: indeed, by (6.1) and (2.7) the minimum problem above is equivalent to

$$\begin{aligned} \min_{(v, \eta, q, \zeta) \in \mathcal{A}_{\text{qs}}(w(t_k^i)) \times L^2(\Omega)} \left\{ \mathcal{Q}(\eta) - \int_\Omega \chi(t_k^i) : \eta dx + \frac{1}{2} \|\zeta\|_2^2 \right. \\ \left. + \mathcal{H}_\lambda(q - p_k^{i-1}, \zeta - z_k^{i-1}) - \langle \chi(t_k^i), q - p_k^{i-1} \rangle \right\}, \end{aligned} \quad (6.6)$$

for which existence of solutions can be proved by the direct method, owing to Proposition 6.1 and the Poincaré-Korn inequality.

By minimality and (2.4), we have that $z_k^i \geq z_k^{i-1}$ a.e. in Ω for all $i \in \{1, \dots, N_k\}$. Moreover, since H_λ satisfies the triangle inequality, the quadruplet $(u_k^i, e_k^i, p_k^i, z_k^i)$ is also a solution of

$$\min_{(v, \eta, q, \zeta) \in \mathcal{A}_{\text{qs}}(w(t_k^i)) \times L^2(\Omega)} \left\{ \mathcal{Q}(\eta) + \frac{1}{2} \|\zeta\|_2^2 + \mathcal{H}_\lambda(q - p_k^i, \zeta - z_k^i) - \int_\Omega f(t_k^i) \cdot v dx \right\}.$$

By Lemma 6.2, setting $(\sigma_k^i, \xi_k^i) := (\mathbb{C}e_k^i, -z_k^i)$, we have

$$- \int_\Omega \sigma_k^i : \eta dx - \int_\Omega z_k^i \zeta dx + \int_\Omega f(t_k^i) \cdot v dx \leq \mathcal{H}_\lambda(q, \zeta) \quad (6.7)$$

for every $(v, \eta, q, \zeta) \in \mathcal{A}_{\text{qs}}(0) \times L^2(\Omega)$. Moreover, using the minimality property in (6.6), the following discrete energy inequality can be proved: for all $j \in \{1, \dots, N_k\}$,

$$\begin{aligned} \mathcal{Q}(e_k^j) - \int_\Omega \chi(t_k^j) : (e_k^j - Ew(t_k^j)) dx + \frac{1}{2} \|z_k^j\|_2^2 \\ + \sum_{i=1}^j \left(\mathcal{H}_\lambda(p_k^i - p_k^{i-1}, z_k^i - z_k^{i-1}) - \langle \chi(t_k^i), p_k^i - p_k^{i-1} \rangle \right) \\ \leq \mathcal{Q}(e_0) - \int_\Omega \chi(0) : (e_0 - Ew(0)) dx + \frac{1}{2} \|z_0\|_2^2 + \sum_{i=1}^j \int_{t_k^{i-1}}^{t_k^i} \int_\Omega \sigma_k^{i-1} : E\dot{w}(s) dx ds \\ - \sum_{i=1}^j \int_{t_k^{i-1}}^{t_k^i} \int_\Omega \dot{\chi}(s) : (e_k^{i-1} - Ew(t_k^{i-1})) dx ds + o(1) \quad \text{as } k \rightarrow \infty. \end{aligned}$$

Let $u_k(t)$, $e_k(t)$, $p_k(t)$, $z_k(t)$, $\sigma_k(t)$, and $\xi_k(t)$ be the piecewise constant right-continuous interpolations of $\{u_k^i\}_{0 \leq i \leq N_k}$, $\{e_k^i\}_{0 \leq i \leq N_k}$, $\{p_k^i\}_{0 \leq i \leq N_k}$, $\{z_k^i\}_{0 \leq i \leq N_k}$, $\{\sigma_k^i\}_{0 \leq i \leq N_k}$, and $\{\xi_k^i\}_{0 \leq i \leq N_k}$. By applying Proposition 6.1 and Helly's Theorem, we have that, up to a subsequence, $p_k(t) \rightharpoonup$

$p(t)$ weakly* in $\mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n})$ and $z_k(t) \rightharpoonup z(t)$ weakly in $L^2(\Omega)$ for all $t \in [0, T]$, where $p \in BV([0, T]; \mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n}))$ and $z \in BV([0, T]; L^2(\Omega))$ with $z(t) \geq z(s)$ for all $0 \leq s \leq t \leq T$. Thanks to a uniqueness argument, we get that, for the same subsequence, $e_k(t) \rightharpoonup e(t)$ weakly in $L^2(\Omega; \mathbb{M}_{sym}^{n \times n})$ and $u_k(t) \rightharpoonup u(t)$ weakly* in $BD(\Omega)$ for all $t \in [0, T]$, for some weakly measurable map $e : [0, T] \rightarrow L^2(\Omega; \mathbb{M}_{sym}^{n \times n})$ and some weakly* measurable map $u : [0, T] \rightarrow BD(\Omega)$, satisfying $(u(t), e(t), p(t)) \in \mathcal{A}_{qs}(w(t))$. Passing to the lower limit in the energy inequality yields

$$\begin{aligned}
 \mathcal{Q}(e(t)) - \int_{\Omega} \chi(t) : (e(t) - Ew(t)) dx + \frac{1}{2} \|z(t)\|_2^2 + \mathcal{D}_{\lambda}(p, z; [0, t]) - \langle \chi(t), p(t) \rangle \\
 \leq \mathcal{Q}(e_0) - \int_{\Omega} \chi(0) : (e_0 - Ew(0)) dx + \frac{1}{2} \|z_0\|_2^2 - \langle \chi(0), p_0 \rangle \\
 + \int_0^t \int_{\Omega} \sigma(s) : E\dot{w}(s) dx ds - \int_0^t \int_{\Omega} \dot{\chi}(s) : (e(s) - Ew(s)) dx ds - \int_0^t \langle \dot{\chi}(s), p(s) \rangle ds,
 \end{aligned}$$

where we used that

$$\sum_{i=1}^j \langle \chi(t_k^i), p_k^i - p_k^{i-1} \rangle = - \int_0^{t_k^j} \langle \dot{\chi}(s), p_k(s) \rangle ds + \langle \chi(t_k^j), p_k^j \rangle - \langle \chi(0), p_0 \rangle.$$

For all $t \in [0, T]$, we define $(\sigma(t), \xi(t)) := (\mathbb{C}e(t), -z(t))$. Passing to the limit in the Euler-Lagrange equation (6.7) leads to

$$- \int_{\Omega} \sigma(t) : \eta dx - \int_{\Omega} z(t) \zeta dx + \int_{\Omega} f(t) \cdot v dx \leq \mathcal{H}_{\lambda}(q, \zeta) \quad (6.8)$$

for every $(v, \eta, q, \zeta) \in \mathcal{A}_{qs}(0) \times L^2(\Omega)$. By Lemma 6.2 this implies that $-\operatorname{div} \sigma(t) = f(t)$ and $(\sigma(t), \xi(t)) \in K_{\lambda}$ a.e. in Ω . This is true also at time $t = 0$ thanks to the assumptions on the initial datum. In order to prove the converse energy inequality, we may proceed as in the proof of [7, Theorem 4.7]. The argument is based on a use of the minimality property together with an approximation of $t \mapsto p(t)$ and $t \mapsto e(t)$ by means of piecewise constant mappings. For p , this is possible since the map $t \mapsto p(t)$ has (at most) countably many discontinuity points for the strong $\mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n})$ -topology. For what concerns $t \mapsto e(t)$, this can be done by approximating its $L^2(\Omega; \mathbb{M}_{sym}^{n \times n})$ -Bochner integral by suitable Riemann sums (see, e.g., [6, Lemma 4.12]). Therefore, the previous energy inequality is actually an equality, namely

$$\begin{aligned}
 \mathcal{Q}(e(t)) - \int_{\Omega} \chi(t) : (e(t) - Ew(t)) dx + \frac{1}{2} \|z(t)\|_2^2 + \mathcal{D}_{\lambda}(p, z; [0, t]) - \langle \chi(t), p(t) \rangle \\
 = \mathcal{Q}(e_0) - \int_{\Omega} \chi(0) : (e_0 - Ew(0)) dx + \frac{1}{2} \|z_0\|_2^2 - \langle \chi(0), p_0 \rangle \\
 + \int_0^t \int_{\Omega} \sigma(s) : E\dot{w}(s) dx ds - \int_0^t \int_{\Omega} \dot{\chi}(s) : (e(s) - Ew(s)) dx ds - \int_0^t \langle \dot{\chi}(s), p(s) \rangle ds.
 \end{aligned}$$

Arguing as in [7, Theorem 5.2], we obtain that $e \in AC([0, T]; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))$, $z \in AC([0, T]; L^2(\Omega))$, $p \in AC([0, T]; \mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n}))$, and by the Poincaré-Korn inequality, $u \in AC([0, T]; BD(\Omega))$. Moreover, the following estimate holds true:

$$\begin{aligned}
 \|\dot{e}\|_{L^1(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))} + \|\dot{z}\|_{L^1(0, T; L^2(\Omega))} + \|\dot{p}\|_{L^1(0, T; \mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n}))} + \|\dot{u}\|_{L^1(0, T; BD(\Omega))} \\
 \leq C \left(\|\dot{w}\|_{L^1(0, T; H^1(\Omega; \mathbb{R}^n))} + \|\dot{\chi}\|_{L^1(0, T; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))} + \|\dot{f}\|_{L^1(0, T; L^n(\Omega; \mathbb{R}^n))} \right), \quad (6.9)
 \end{aligned}$$

where $C > 0$ is a constant depending on the norms $\|\chi\|_{L^\infty(0,T;L^2(\Omega;\mathbb{M}_{sym}^{n \times n}))}$, $\|\vartheta\|_{L^\infty(0,T;L^2(\Omega))}$, $\|e\|_{L^\infty(0,T;L^2(\Omega;\mathbb{M}_{sym}^{n \times n}))}$, $\|p\|_{L^\infty(0,T;\mathcal{M}(\bar{\Omega};\mathbb{M}_{sym}^{n \times n}))}$, $\|z\|_{L^\infty(0,T;L^2(\Omega))}$, and $\|w\|_{L^\infty(0,T;H^1(\Omega;\mathbb{R}^n))}$, but independent of λ . Note that in order to derive estimate (6.9), we use at some point the coercivity property of Proposition 6.1 and the following estimate for the stress/strain duality: for all $\tau \in L^n(\Omega;\mathbb{M}_{sym}^{n \times n})$ with $\operatorname{div}\tau \in L^n(\Omega;\mathbb{R}^n)$, and for all $s \in (0, T)$,

$$|\langle \tau, p(s) \rangle| \leq C(\|\operatorname{div}\tau\|_n + \|\tau\|_2)(\|e(s)\|_2 + \|p(s)\|_1 + \|w(s)\|_{H^1(\Omega;\mathbb{R}^n)}),$$

for some constant $C > 0$ depending only on Ω . In particular, $\dot{z}(t) \geq 0$ for a.e. $t \in [0, T]$ and a.e. $x \in \Omega$, and according to Proposition 2.2, we infer that for all $t \in [0, T]$

$$\begin{aligned} \mathcal{Q}(e(t)) - \int_{\Omega} \chi(t) : (e(t) - Ew(t)) \, dx + \frac{1}{2}\|z(t)\|_2^2 + \int_0^t \mathcal{H}_\lambda(\dot{p}(s), \dot{z}(s)) \, ds - \langle \chi(t), p(t) \rangle \\ = \mathcal{Q}(e_0) - \int_{\Omega} \chi(0) : (e_0 - Ew(0)) \, dx + \frac{1}{2}\|z_0\|_2^2 - \langle \chi(0), p_0 \rangle \\ + \int_0^t \int_{\Omega} \sigma(s) : E\dot{w}(s) \, dx \, ds - \int_0^t \int_{\Omega} \dot{\chi}(s) : (e(s) - Ew(s)) \, dx \, ds - \int_0^t \langle \dot{\chi}(s), p(s) \rangle \, ds. \end{aligned} \quad (6.10)$$

By applying again the duality formula (2.7) and (6.1) we get

$$\begin{aligned} \mathcal{Q}(e(t)) + \frac{1}{2}\|z(t)\|_2^2 + \int_0^t \mathcal{H}_\lambda(\dot{p}(s), \dot{z}(s)) \, ds + \int_{\Omega} f(t) \cdot (w(t) - u(t)) \, dx \\ = \mathcal{Q}(e_0) + \frac{1}{2}\|z_0\|_2^2 + \int_{\Omega} f(0) \cdot (w(0) - u(0)) \, dx \\ + \int_0^t \int_{\Omega} \sigma(s) : E\dot{w}(s) \, dx \, ds + \int_0^t \int_{\Omega} \dot{f}(s) \cdot (w(s) - u(s)) \, dx \, ds, \end{aligned}$$

which is equivalent to (6.5) by integration by parts in time of the last integral. This completes the proof of the theorem. \square

The following result states a more precise formulation of the energy equality as a flow rule, whenever additional integrability properties are satisfied for the stress and/or the velocity.

Theorem 6.4. *Assume that either $\sigma(t) \in L^n(\Omega;\mathbb{M}_{sym}^{n \times n})$ or $\dot{u}(t) \in L^2(\Omega;\mathbb{R}^n)$ for a.e. $t \in [0, T]$. Then the distribution $[\sigma(t) : \dot{p}(t)]$ is well defined for a.e. $t \in [0, T]$, and it is a measure in $\mathcal{M}(\bar{\Omega})$ satisfying*

$$H_\lambda(\dot{p}(t), \dot{z}(t)) = [\sigma(t) : \dot{p}(t)] + \xi(t)z(t) \quad \text{in } \mathcal{M}(\bar{\Omega}), \quad \text{for a.e. } t \in [0, T]. \quad (6.11)$$

Moreover, the stress σ and the cap variable ξ are unique.

Proof. By Definition 2.3, if $\dot{u}(t) \in L^2(\Omega;\mathbb{R}^n)$, then the distribution $[\tau : \dot{p}(t)]$ is well defined for every $\tau \in L^2(\Omega;\mathbb{M}_{sym}^{n \times n})$ with $\operatorname{div}\tau \in L^2(\Omega;\mathbb{R}^n)$. Equality (6.11) is obtained from (6.5) exactly as in the proof of Proposition 4.7. By Remark 2.6, since $\operatorname{div}\sigma(t) = -f(t) \in L^n(\Omega;\mathbb{R}^n)$, the same conclusion holds if $\sigma(t) \in L^n(\Omega;\mathbb{M}_{sym}^{n \times n})$.

Once the stress-strain duality pairing is defined, it is possible to argue as in [7, Theorem 5.9] to establish the uniqueness of σ and ξ . \square

Note that the assumptions of Theorem 6.4 are clearly satisfied for $n = 2$. However, it is not clear to us if such integrability properties for the stress and/or the velocity are true in higher dimension.

6.2. Convergence of the quasi-static cap model. In this section we characterize the asymptotic behaviour of the quasi-static evolutions studied in Theorems 6.3 and 6.4 when the cap is sent to infinity.

We consider a body load $f \in AC([0, T]; L^n(\Omega; \mathbb{R}^n))$ satisfying the following safe-load condition: there exist $\chi \in AC([0, T]; L^n(\Omega; \mathbb{M}_{sym}^{n \times n}))$ and a constant $\alpha_0 > 0$ such that for every $t \in [0, T]$

$$-\operatorname{div}\chi(t) = f(t) \quad \text{a.e. in } \Omega \quad (6.12)$$

and

$$\chi(t) + \tau \in K \quad \text{a.e. in } \Omega$$

for every $\tau \in \mathbb{M}_{sym}^{n \times n}$ with $|\tau| \leq \alpha_0$. We also consider a boundary displacement which is the trace on $\partial\Omega \times (0, T)$ of a function $w \in AC([0, T]; H^1(\Omega; \mathbb{R}^n))$ and an initial datum $(u_0, e_0, p_0) \in \mathcal{A}_{qs}(w(0))$ satisfying the stability condition

$$\mathcal{Q}(e_0) - \int_{\Omega} f(0) \cdot u_0 \, dx \leq \mathcal{Q}(\eta) + \mathcal{H}(q - p_0) - \int_{\Omega} f(0) \cdot v \, dx,$$

for any $(v, \eta, q) \in \mathcal{A}_{qs}(w(0))$. We next define $(\sigma_0, \xi_0) := (\mathbb{C}e_0, -z_0)$ and we further assume that $\sigma_0 \in L^n(\Omega; \mathbb{M}_{sym}^{n \times n})$.

In order to apply Theorems 6.3 and 6.4, we set $\vartheta := \chi_m - |\chi_m| - \alpha_0/\sqrt{n} \in AC([0, T]; L^2(\Omega))$ and we observe that

$$(\chi(t) + \tau, \vartheta(t)) \in K_{\lambda} \quad \text{in } \Omega$$

for every $\tau \in \mathbb{M}_{sym}^{n \times n}$ with $|\tau| \leq \alpha_0$ and every $\lambda \geq 1$, so that f satisfies (6.1) and (6.2). We define an initial datum for the hardening cap variable as in the dynamical case by

$$\xi_0 := (\sigma_0)_m - |(\sigma_0)_m|, \quad \text{and } z_0 := -\xi_0,$$

so that $\xi_0, z_0 \in L^2(\Omega)$ and $(\sigma_0, \xi_0) \in K_{\lambda}$ a.e. in Ω , for all $\lambda \geq 1$. Since $\sigma_0 \in L^n(\Omega; \mathbb{M}_{sym}^{n \times n})$ by assumption, by Lemma 6.2 we infer that (u_0, e_0, p_0, z_0) is a solution of

$$\min_{(v, \eta, q, \zeta) \in \mathcal{A}_{qs}(w(0)) \times L^2(\Omega)} \left\{ \mathcal{Q}(\eta) + \frac{1}{2} \|\zeta\|_2^2 + \mathcal{H}_{\lambda}(q - p_0, \zeta - z_0) - \int_{\Omega} f(0) \cdot v \, dx \right\}.$$

For every $\lambda \geq 1$ let $u_{\lambda}, e_{\lambda}, \sigma_{\lambda}, p_{\lambda}, z_{\lambda}, \xi_{\lambda}$ be the solution to (6.3)–(6.5) constructed in Theorem 6.3 with initial datum (u_0, e_0, p_0, z_0) .

Theorem 6.5. *There exist*

$$\begin{cases} u \in AC([0, T]; BD(\Omega)), \\ \sigma, e \in AC([0, T]; L^2(\Omega; \mathbb{M}_{sym}^{n \times n})), \\ p \in AC([0, T]; \mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n})), \end{cases}$$

with $(u(0), e(0), p(0)) = (u_0, e_0, p_0)$, such that for all $t \in [0, T]$

$$\begin{aligned} u_{\lambda}(t) &\rightharpoonup u(t) \text{ weakly* in } BD(\Omega), \\ e_{\lambda}(t) &\rightarrow e(t) \text{ strongly in } L^2(\Omega; \mathbb{M}_{sym}^{n \times n}), \\ p_{\lambda}(t) &\rightharpoonup p(t) \text{ weakly* in } \mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n}), \end{aligned} \quad (6.13)$$

as $\lambda \rightarrow \infty$. For every $t \in [0, T]$, we have

$$\begin{cases} Eu(t) = e(t) + p(t) \text{ in } \Omega, & p(t) = (w(t) - u(t)) \odot \nu \mathcal{H}^{n-1} \text{ on } \partial\Omega, \\ \sigma(t) = \mathbb{C}e(t), \end{cases}$$

and

$$\begin{cases} -\operatorname{div}\sigma(t) = f(t) \text{ a.e. in } \Omega, \\ \sigma(t) \in K \text{ a.e. in } \Omega. \end{cases}$$

Moreover, the following energy equality holds

$$\begin{aligned} \mathcal{Q}(e(t)) + \int_0^t \mathcal{H}(\dot{p}(s)) ds &= \mathcal{Q}(e_0) \\ &+ \int_0^t \int_{\Omega} \sigma(s) : E\dot{w}(s) dx ds - \int_0^t \int_{\Omega} f(s) \cdot (\dot{w}(s) - \dot{u}(s)) dx ds. \end{aligned} \quad (6.14)$$

Proof. Some *a priori* estimates for the sequences (u_λ) , (e_λ) , (p_λ) , and (z_λ) can be obtained from the uniform estimate (6.9) and the energy equality (6.10). They imply the existence of $u \in AC([0, T]; BD(\Omega))$, $e \in AC([0, T]; L^2(\Omega; \mathbb{M}_{sym}^{n \times n}))$, $p \in AC([0, T]; \mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n}))$, and $z \in AC([0, T]; L^2(\Omega))$ such that $u_\lambda(t) \rightharpoonup u(t)$ weakly* in $BD(\Omega)$, $e_\lambda(t) \rightharpoonup e(t)$ weakly in $L^2(\Omega; \mathbb{M}_{sym}^{n \times n})$, $p_\lambda(t) \rightharpoonup p(t)$ weakly* in $\mathcal{M}(\bar{\Omega}; \mathbb{M}_{sym}^{n \times n})$, and $z_\lambda(t) \rightharpoonup z(t)$ weakly in $L^2(\Omega)$. Moreover, we have that $(u(0), e(0), p(0), z(0)) = (u_0, e_0, p_0, z_0)$, $Eu(t) = e(t) + p(t)$ in Ω , $p(t) = (w(t) - u(t)) \odot \nu \mathcal{H}^{n-1}$ on $\partial\Omega$ for every $t \in [0, T]$, and $\dot{z}(t) \geq 0$ a.e. in Ω , for a.e. $t \in [0, T]$. In particular, we infer that $z(t) \geq z_0$ a.e. in Ω for every $t \in [0, T]$.

Setting $\sigma(t) := \mathbb{C}e(t)$, passing to the limit in (6.8) as $\lambda \rightarrow +\infty$, and applying Lemma 2.1, we obtain

$$- \int_{\Omega} \sigma(t) : \eta dx - \int_{\Omega} z(t) \zeta dx + \int_{\Omega} f(t) \cdot v dx \leq \mathcal{H}(q)$$

for every $(v, \eta, q, \zeta) \in \mathcal{A}_{qs}(0) \times L^2(\Omega)$ with $\zeta \geq 0$ in Ω . For $\zeta \equiv 0$ this implies that

$$- \int_{\Omega} \sigma(t) : \eta dx + \int_{\Omega} f(t) \cdot v dx \leq \mathcal{H}(q)$$

for every $(v, \eta, q) \in \mathcal{A}_{qs}(0)$. By [7, Theorem 3.6] this condition is equivalent to saying that $(u(t), e(t), p(t))$ minimizes the functional

$$\mathcal{Q}(e) + \mathcal{H}(p - p(t)) - \int_{\Omega} f(t) \cdot u dx$$

over all $(u, e, p) \in \mathcal{A}_{qs}(w(t))$. This, in turn, implies that $\sigma(t) \in K$ and $-\text{div}\sigma(t) = f(t)$ a.e. in Ω .

Furthermore, arguing as in the proof of Theorem 5.1, we deduce the following energy inequality:

$$\begin{aligned} \mathcal{Q}(e(t)) - \int_{\Omega} \chi(t) : (e(t) - Ew(t)) dx + \int_0^t \mathcal{H}(\dot{p}(s)) ds - \langle \chi(t), p(t) \rangle \\ \leq \mathcal{Q}(e_0) - \int_{\Omega} \chi(0) : (e_0 - Ew(0)) dx - \langle \chi(0), p_0 \rangle \\ + \int_0^t \int_{\Omega} \sigma(s) : E\dot{w}(s) dx ds - \int_0^t \int_{\Omega} \dot{\chi}(s) : (e(s) - Ew(s)) dx ds - \int_0^t \langle \dot{\chi}(s), p(s) \rangle ds. \end{aligned}$$

Finally, we argue as in the proof of Theorem 6.3 to show that this inequality is actually an equality leading to (6.14) by (2.7), (6.12), and integration by parts with respect to t . The strong convergence (6.13) can be proved as in [7, Theorem 4.8]. \square

Once again, provided the stress and/or the velocity have enough integrability in such a way that the stress-strain duality is well defined, one can write the flow rule in a measure theoretic sense.

Theorem 6.6. *Assume that either $\sigma(t) \in L^n(\Omega; \mathbb{M}_{sym}^{n \times n})$ or $\dot{u}(t) \in L^2(\Omega; \mathbb{R}^n)$ for a.e. $t \in [0, T]$. Then the distribution $[\sigma(t) : \dot{p}(t)]$ is well defined for a.e. $t \in [0, T]$, and it is a measure in $\mathcal{M}(\bar{\Omega})$ satisfying*

$$H(\dot{p}(t)) = [\sigma(t) : \dot{p}(t)] \quad \text{in } \mathcal{M}(\bar{\Omega}), \quad \text{for a.e. } t \in [0, T].$$

Moreover, the stress σ is unique.

We again observe that the assumptions of Theorem 6.6 are clearly satisfied in the two-dimensional setting.

ACKNOWLEDGEMENTS

The authors wish to thank the hospitality of the Laboratoire Jacques-Louis Lions at Pierre et Marie Curie University and of the Mathematics Department of the University of Pavia, where a large part of this work was carried out. The research of JFB has been supported by the Agence Nationale de la Recherche under Grant No. ANR 10-JCJC 0106. MGM acknowledges support from the European Research Council under Grant No. 290888 “Quasistatic and Dynamic Evolution Problems in Plasticity and Fracture”.

REFERENCES

- [1] L. AMBROSIO, N. FUSCO & D. PALLARA: *Functions of bounded variation and free discontinuity problems*, Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York (2000).
- [2] G. ANZELLOTTI: On the existence of the rates of stress and displacements for Prandtl-Reuss plasticity, *Quart. Appl. Math.* **41** (1984) 181–208.
- [3] G. ANZELLOTTI & S. LUCKHAUS: Dynamical evolution in elasto-perfectly plastic bodies, *Appl. Math. Opt.* **15** (1987) 121–140.
- [4] A. BENSOUSSAN & J. FREHSE: Asymptotic behaviour of the time dependent Norton-Hoff law in plasticity theory and H^1 regularity, *Comment. Math. Univ. Carolin.* **37** (1996) 285–304.
- [5] H. BRÉZIS: *Opérateurs maximaux monotones et semi groupes de contractions dans les espaces de Hilbert*, North-Holland, Amsterdam-London; American Elsevier, New York (1973).
- [6] G. DAL MASO, G. A. FRANCFORT & R. TOADER: Quasistatic crack growth in nonlinear elasticity, *Arch. Rational Mech. Anal.* **176** (2005) 165–225.
- [7] G. DAL MASO, A. DE SIMONE & M. G. MORA: Quasistatic evolution problems for linearly elastic–perfectly plastic materials, *Arch. Rational Mech. Anal.* **180** (2006) 237–291.
- [8] G. DAL MASO, A. DE SIMONE & F. SOLOMBRINO: Quasistatic evolution for Cam-Clay plasticity: a weak formulation via viscoplastic regularization and time rescaling, *Calc. Var. PDEs* **40** (2011) 125–181.
- [9] F. DEMENGEL & R. TEMAM: Convex function of a measure, *Indiana Univ. Math. J.* **33** (1984) 673–709.
- [10] F. DEMENGEL & R. TEMAM: Convex function of a measure: the unbounded case, FERMAT days 1985: mathematics for optimization, 103–134, North-Holland (1986).
- [11] A. DEMYANOV: Regularity of stresses in Prandtl-Reuss perfect plasticity, *Calc. Var. PDEs* **34** (2009) 23–72.
- [12] F. L. DIMAGGIO & I. S. SANDLER: Material model for granular soils, *J. Engrg. Mech.* **97** (1971) 935–950.
- [13] D. C. DRUCKER & W. PRAGER: Soil mechanics and plastic analysis of rock and concrete, *Quart. Appl. Math.* **10** (1952) 157–175.
- [14] G. DUVAUT & J.-L. LIONS: *Les inéquations en mécanique et en physique*, Dunod, Paris (1972).
- [15] L. C. EVANS: *Partial Differential equations*, Second edition, Graduate Studies in Mathematics, 19. American Mathematical Society, Providence, RI (2010).
- [16] G. A. FRANCFORT & A. GIACOMINI: Small strain heterogeneous elasto-plasticity revisited, *Comm. Pure Appl. Math.* **65** (2012) 1185–1241.
- [17] C. GOFFMAN & J. SERRIN: Sublinear functions of measures and variational integrals, *Duke Math. J.* **31** (1964) 159–178.
- [18] R. HILL: *The mathematical theory of plasticity*, Clarendon press, Oxford (1950).
- [19] C. JOHNSON: Existence theorems for plasticity problems, *J. Math. Pures Appl.* **55** (1976) 431–444.
- [20] R.V. KOHN & R. TEMAM: Dual spaces of stresses and strains, with applications to Hencky plasticity, *Appl. Math. Optim.* **10** (1983) 1–35.
- [21] J. LUBLINER: *Plasticity Theory*, Macmillan Publishing Company, New York (1990).
- [22] S. LUCKHAUS: Elastisch-plastische Materialien mit Viskosität, *Preprint no. 65 of Heidelberg University, SFB 123*.
- [23] A. MAINIK & A. MIELKE: Existence results for energetic models for rate-independent systems, *Calc. Var. PDEs* **22** (2005) 73–99.
- [24] H. MATTHIES, G. STRANG & E. CHRISTIANSEN: The saddle point of a differential program, *Energy methods in finite element analysis*, Ed. Rodin, Zienkiewicz, Glowinski, Wiley, New York (1979).
- [25] J.-J. MOREAU: Application of convex analysis to the treatment of elasto-plastic systems, Springer Lecture Notes in Math., no. 503, Ed. P. Germain and B. Nayroles (1975).
- [26] S. REPIN & G. SEREGIN: Existence of weak solution of the minimax problem arising in Coulomb-Mohr plasticity, *Amer. Math. Soc. Transl.* **164** (1995) 189–220.
- [27] L. RESENDE & J.B. MARTIN: Formulation of Drucker-Prager Cap Model, *J. Engrg. Mech.* **111** (1985) 855–881.
- [28] L. RESENDE & J.B. MARTIN: Closure of “Formulation of Drucker-Prager Cap Model”, *J. Engrg. Mech.* **113** (1987) 1257–1259.

- [29] J. SALENÇON: *Application of the Theory of Plasticity to Soil Mechanics*, Wiley Ser. Geotechnical Engrg., John Wiley, New York (1977).
- [30] G. SEREGIN: On the differentiability of the stress tensor in the Coulomb-Mohr theory of plasticity, *St. Petersburg Math. J.* **4** (1993) 1257–1271.
- [31] P. SUQUET: Sur les équations de la plasticité: existence et régularité des solutions, *J. Mécanique* **20** (1981) 3–39.
- [32] P. SUQUET: Evolution problems for a class of dissipative materials, *Quart. Appl. Math.* **38** (1981) 391–414.
- [33] P. SUQUET: Un espace fonctionnel pour les équations de la plasticité, *Ann. Fac. Sci. Toulouse Math.* **1** (1979) 77–87.
- [34] R. TEMAM: *Mathematical problems in plasticity*, Gauthier-Villars, Paris (1985). Translation of *Problèmes mathématiques en plasticité*, Gauthier-Villars, Paris (1983).
- [35] R. TEMAM: A generalized Norton-Hoff model and the Prandtl-Reuss law of plasticity, *Arch. Rational Mech. Anal.* **95** (1986) 137–183.

(J.-F. Babadjian) UNIVERSITÉ PIERRE ET MARIE CURIE – PARIS 6, CNRS, UMR 7598 LABORATOIRE JACQUES-LOUIS LIONS, PARIS, F-75005, FRANCE
E-mail address: babadjian@ann.jussieu.fr

(M.G. Mora) DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PAVIA, VIA FERRATA 1, 27100 PAVIA, ITALY
E-mail address: mariagiovanna.mora@unipv.it