



Addendum to Pontryagin's maximum principle for dynamic systems on time scales

Loïc Bourdin^a, Oleksandr Stanzhytskyi^b and Emmanuel Trélat^c

^aXLIM Research Institute, University of Limoges, CNRS UMR 7252, Limoges, France; ^bTaras Shevchenko National University of Kyiv, Kyiv, Ukraine; ^cLaboratoire Jacques-Louis Lions, Sorbonne Universités, UPMC Univ Paris 06, CNRS UMR 7598, Paris, France

ABSTRACT

This note is an addendum to [L. Bourdin and E. Trélat, *SIAM J. Cont. Optim.*, 2013] and [M. Bohner, K. Kenzhebaev, O. Lavrova and O. Stanzhytskyi, *J. Differ. Equ. Appl.*, 2017], pointing out the differences between these papers and raising open questions.

ARTICLE HISTORY

Received 9 May 2017
Accepted 16 July 2017

KEYWORDS

Time scale; optimal control; Pontryagin maximum principle; Ekeland variational principle; packages of needle-like variations

AMS SUBJECT

CLASSIFICATIONS

34K35; 34N99; 39A12;
39A13; 49K15; 93C15; 93C55

The main differences. In view of establishing a time scale version of the Pontryagin Maximum Principle (PMP), the authors of [1, Theorem 1] have developed in 2013 a strategy of proof based on the *Ekeland variational principle*. This strategy was originally considered for the classical continuous case by Ivar Ekeland in his seminal paper [3].

The authors of [2, Theorem 2.11] developed in 2017 a different approach, with *packages of needle-like variations* and *necessary conditions for an extreme in a cone*. Note that the authors of [2] prove moreover in [2, Theorem 2.13] that the necessary conditions derived in the PMP are also sufficient in the linear-convex case.

In the sequel of this paragraph, we focus on the major pros and cons of each approach:

- (1) In [1]:
 - (a) The set Ω of control constraints is assumed to be closed. This is in order to apply the Ekeland variational principle on a complete metric space.
 - (b) There is no assumption on the time scale \mathbb{T} .
- (2) In [2]:
 - (a) The set Ω of control constraints is assumed to be convex, but need not to be closed.

(b) The time scale \mathbb{T} is assumed to satisfy *density conditions* (see [2, Definition 2.4]) of the kind

$$\lim_{\substack{\beta \rightarrow 0^+ \\ s+\beta \in \mathbb{T}}} \frac{\mu(s + \beta)}{\beta} = 0,$$

for every right-dense points s , in order to guarantee that

$$\lim_{\beta \rightarrow 0^+} \frac{1}{\beta} \int_{[s, s+\beta)_{\mathbb{T}}} x(\tau) \Delta \tau = x(s),$$

for Δ -integrable function x and for right-dense Δ -Lebesgue points s , even for $\beta > 0$ such that $s + \beta \notin \mathbb{T}$. Note that a discussion about this issue was provided in [1, Section 3.1].

Hence, the method developed in [2] allows to remove the closedness assumption done on Ω in [1], but this is at the price of an additional assumption on the time scale \mathbb{T} .

In [1, Section 3.1], the authors explained why other approaches (other than the Ekeland variational principle), based for instance on implicit function arguments, or on Brouwer fixed point arguments, or on separation (Hahn-Banach) arguments, fail for general time scales.

As a conclusion, a time scale version of the PMP without closedness assumption on Ω and without any assumption on the time scale \mathbb{T} still remains an open challenge.

Additional comments on the terminal constraints. In [1] the authors considered constraints on the initial/final state of the kind $g(x(t_0), x(t_1)) \in S$, where S is a nonempty closed convex set and g is a general smooth function.

In [2] the authors considered terminal state constraints of the kind $\Phi_i(x(t_0), x(t_1)) = 0$ for $i = 1, \dots, k$, and $\Phi_i(x(t_0), x(t_1)) \leq 0$ for $i = k + 1, \dots, n$, where Φ_i are general smooth functions.

Contrarily to what is claimed in [2], the terminal constraints considered in [2] are only a particular case of the ones considered in [1]. Indeed, it suffices to take

$$g = (\Phi_1, \dots, \Phi_k, \Phi_{k+1}, \dots, \Phi_n)$$

and

$$S = \{0\} \times \dots \times \{0\} \times \mathbb{R}^- \times \dots \times \mathbb{R}^-.$$

Moreover, note that the necessary condition $-\Psi \in \mathcal{O}_S(g(x(t_0), x(t_1)))$ obtained in [1, Theorem 1] encompasses both the *sign condition* (1) and the *complementary slackness* (2) obtained in [2, Theorem 2.11]. For the sign condition, it is sufficient to recall that the orthogonal of \mathbb{R}^- at a point $x \in \mathbb{R}^-$ is included in \mathbb{R}^+ . For the complementary slackness, it is sufficient to recall that the orthogonal of S at $g(x(t_0), x(t_1))$ is reduced to $\{0\}$ when $g(x(t_0), x(t_1))$ belongs to the interior of S .

Additional comments on the convexity of Ω . The set Ω is assumed to be convex in [2], while it is not in [1]. As explained in [1, Section 3.1], in order to apply necessary conditions of an extreme in a cone, the authors of [2] require that the parameters of perturbations live in intervals. As a consequence, in order to remove the convexity assumption on Ω , one would need (local-directional) convexity of the set Ω for perturbations at right-scattered

points, which is a concept that differs from the stable Ω -dense directions used in [1]. Hence, in spite of the claim done in [2], the convexity assumption on Ω does not seem to be easily removable.

On the universal Lagrange multipliers. This paragraph is devoted to providing more details on the existence of universal Lagrange multipliers claimed in [2, p.25]. In the sequel, we use the notations of [2], and we denote by \mathcal{S} the unit sphere of \mathbb{R}^{n+1} .

A package P consists of:

- $N \in \mathbb{N}$ and $\nu \in \mathbb{N}$;
- $\bar{\tau} = (\tau_1, \dots, \tau_N)$ where τ_i are right-dense points of \mathbb{T} ;
- $\bar{\nu} = (\nu_1, \dots, \nu_N)$ where $\nu_i \in U$;
- $\bar{r} = (r_1, \dots, r_\nu)$ where r_i are right-scattered points of \mathbb{T} .
- $\bar{z} = (z_1, \dots, z_\nu)$ where $z_i \in U$.

Let $(P_i)_{i \in I}$ denotes the set of all possible packages.

Following the proof of [2, Theorem 2.11], for every $i \in I$, there exists a nonzero vector $\lambda = (\lambda_0, \dots, \lambda_n)$ (that we renormalize in \mathcal{S}) of Lagrange multipliers such that:

- (i) (1) and (2) in [2, Theorem 2.11] are satisfied;
- (ii) the adjoint vector Ψ solution of (2.9), with the final condition (3.65) which depends on λ , satisfies the initial condition $\Psi(t_0) = L_{x_0}$;
- (iii) (4a) and (4b) in [2, Theorem 2.11] are satisfied, but only at the points contained in $\bar{\tau}$ and \bar{r} respectively.

For every $i \in I$, the above vector λ is not necessarily unique. Then, for every $i \in I$, we denote by K_i the set of all nonzero and renormalized Lagrange multiplier vectors associated with P_i satisfying the above properties.

By continuity of the adjoint vector Ψ with respect to the Lagrange multipliers (dependence from its final condition), we infer that K_i is a nonempty closed subset contained in the compact \mathcal{S} . This is true for every $i \in I$.

Now, let us prove that the family $(K_i)_{i \in I}$ satisfies the finite intersection property. Let $J \subset I$ be a finite subset and let us prove that $\bigcap_{i \in J} K_i \neq \emptyset$. Note that we can construct a package P corresponding to the union of all packages P_i with $i \in J$. It follows that $P \in (P_i)_{i \in I}$, and thus there exists a nonzero and renormalized Lagrange multiplier vector λ associated with P satisfying the above properties. Since $\lambda \in K_i$ for every $i \in J$, we conclude that $\bigcap_{i \in J} K_i \neq \emptyset$.

It follows from the lemma of a centered system in a compact set that $\bigcap_{i \in I} K_i \neq \emptyset$, and we deduce the existence of a universal Lagrange multiplier vector.

On the density conditions and the Cantor set. Contrarily to what is claimed in [2, Example 2.5], the classical Cantor set does not satisfy the density conditions. However, generalized versions of the Cantor set (see, e.g., [4]) that satisfy density conditions can be constructed as follows.

Let $(\alpha_k)_{k \in \mathbb{N}}$ be a real sequence such that $0 < \alpha_k < \frac{1}{2}$ for all $k \in \mathbb{N}$, and such that $\lim_{k \rightarrow +\infty} \alpha_k = \frac{1}{2}$. Let $(A_k)_{k \in \mathbb{N}}$ be the sequence of compact subsets defined by the induction

$$A_0 = [0, 1], \quad A_{k+1} = \mathcal{T}_k(A_k) \quad \forall k \in \mathbb{N},$$

where \mathcal{T}_k denotes the operator removing the open $(\alpha_k, 1 - \alpha_k)$ -central part of all intervals. Note that the classical Cantor set corresponds to the case where $\alpha_k = \frac{1}{3}$ for every $k \in \mathbb{N}$.

In our situation, we obtain

$$A_1 = [0, \alpha_0] \cup [1 - \alpha_0, 1],$$

$$A_2 = \left([0, \alpha_1 \alpha_0] \cup [(1 - \alpha_1) \alpha_0, \alpha_0] \right) \cup \left([1 - \alpha_0, 1 - (1 - \alpha_1) \alpha_0] \cup [1 - \alpha_1 \alpha_0, 1] \right),$$

etc. We define the generalized Cantor set $\mathbb{T} = \bigcap_{k \in \mathbb{N}} A_k$. In order to prove that the time scale \mathbb{T} satisfies the density conditions, from the fractal properties of \mathbb{T} , it suffices to prove that the density condition is satisfied at the right-dense point $0 \in \mathbb{T}$. More precisely, it is sufficient to prove that

$$\lim_{\substack{\beta \rightarrow 0^+ \\ \beta \in \mathbb{T}}} \frac{\mu(\beta)}{\beta} = 0.$$

Since $\mu(\beta) = 0$ for every right-dense point β , we only have to consider the case where β is a right-scattered point of \mathbb{T} . In that case, one can easily see that $\frac{\mu(\beta)}{\beta} \leq \frac{1 - 2\alpha_k}{\alpha_k}$ for some $k \in \mathbb{N}$ and that k tends to $+\infty$ when β tends to 0. The conclusion follows from the fact that $\lim_{k \rightarrow +\infty} \alpha_k = \frac{1}{2}$.

References

- [1] L. Bourdin and E. Trélat, *Pontryagin maximum principle for finite dimensional nonlinear optimal control problems on time scales*, SIAM J. Control Optim. 51(5) (2013), pp. 3781–3813.
- [2] M. Bohner, K. Kenzhebaev, O. Lavrova, and O. Stanzhytskyi, *Pontryagin’s maximum principle for dynamic systems on time scales*, J. Differ. Equ. Appl. (2017). DOI:10.1080/10236198.2017.1284829
- [3] I. Ekeland, *On the variational principle*, J. Math. Anal. Appl. 47 (1974), pp. 324–353.
- [4] A.F. Turbin and N.V. Pratsevity, *Fractal Sets, Functions, Distribution*, Naukova Dumka, Kyiv, 1992.