




Contents lists available at ScienceDirect

Systems & Control Letters

journal homepage: www.elsevier.com/locate/sysconle

Asymptotic approach on conjugate points for minimal time bang–bang controls

C. Silva^a, E. Trélat^{b,*}^a Department of Mathematics, University of Aveiro, Portugal and Math., Labo. MAPMO, Université d'Orléans, France^b Université d'Orléans, UFR Sciences, Math., Labo. MAPMO, UMR 6628, Route de Chartres, BP 6759, 45067 Orléans cedex 2, France

ARTICLE INFO

Article history:

Received 27 April 2010

Received in revised form

23 August 2010

Accepted 24 August 2010

Available online 22 September 2010

Keywords:

Optimal control

Minimal time problem

Bang–bang control

Conjugate time

ABSTRACT

We focus on the minimal time control problem for single-input control-affine systems $\dot{x} = X(x) + u_1 Y_1(x)$ in \mathbb{R}^n with fixed initial and final time conditions $x(0) = \hat{x}_0, x(t_f) = \hat{x}_1$, and where the scalar control u_1 satisfies the constraint $|u_1(\cdot)| \leq 1$. For these systems a concept of conjugate time t_c has been defined in e.g. Agrachev et al. (2002) [23], Maurer and Osmolovskii (2004) [21], and Noble and Schättler (2002) [28] in the bang–bang case. Besides, theoretical and practical issues for conjugate time theory are well known in the smooth case (see e.g. Agrachev and Sachkov (2004) [43] and Milyutin and Osmolovskii (1998) [15]), and efficient implementation tools are available (see Bonnard et al. (2007) [35]). The first conjugate time along an extremal is the time at which the extremal loses its local optimality. In this work, we use the asymptotic approach developed in Silva and Trélat (in press) [36] and investigate the convergence properties of conjugate times. More precisely, for $\varepsilon > 0$ small and arbitrary vector fields Y_1, \dots, Y_m , we consider the minimal time problem for the control system $\dot{x}^\varepsilon = X(x^\varepsilon) + u_1^\varepsilon Y_1(x^\varepsilon) + \varepsilon \sum_{i=2}^m u_i^\varepsilon Y_i(x^\varepsilon)$, under the constraint $\sum_{i=1}^m (u_i^\varepsilon)^2 \leq 1$, with the fixed boundary conditions $x^\varepsilon(0) = \hat{x}_0, x^\varepsilon(t_f) = \hat{x}_1$ of the initial problem. Under appropriate assumptions, the optimal controls of the latter regularized optimal control problem are smooth, and the computation of associated conjugate times t_c^ε falls into the standard theory; our main result asserts the convergence, as ε tends to 0, of t_c^ε towards the conjugate time t_c of the initial bang–bang optimal control problem, as well as the convergence of the associated extremals. As a byproduct, we obtain an efficient algorithmic way to compute conjugate times in the bang–bang case.

© 2010 Elsevier B.V. All rights reserved.

1. Introduction

1.1. The optimal control problem

Consider the single-input control-affine system in \mathbb{R}^n

$$\dot{x} = X(x) + u_1 Y_1(x), \quad (1)$$

where X and Y_1 are smooth vector fields, and the control u_1 is a measurable scalar function satisfying the constraint

$$|u_1(\cdot)| \leq 1. \quad (2)$$

Let \hat{x}_0 and \hat{x}_1 be two points of \mathbb{R}^n . Assume that \hat{x}_1 is reachable from \hat{x}_0 , that is, there exists a time $T > 0$ and a control function $u_1(\cdot) \in L^\infty(0, T)$ satisfying the constraint (2), such that the trajectory $x(\cdot)$, solution of (1) with $x(0) = \hat{x}_0$, satisfies $x(T) = \hat{x}_1$.

We consider the optimal control problem (OCP) of determining a solution $\hat{x}(\cdot)$ associated to a control $\hat{u}_1(\cdot)$, on $[0, t_f]$, satisfying

(1)–(2) and steering \hat{x}_0 to \hat{x}_1 in minimal time t_f . We assume that such a solution $\hat{x}(\cdot)$ for (OCP) exists.¹

According to the Pontryagin maximum principle (see [2]), there exists a absolutely continuous mapping $\hat{p}(\cdot) : [0, t_f] \rightarrow \mathbb{R}^n$, called *adjoint vector*, and a real number $p^0 \leq 0$, with $(\hat{p}(\cdot), p^0) \neq (0, 0)$, such that

$$\begin{aligned} \dot{\hat{p}}(t) &= -\frac{\partial H}{\partial x}(\hat{x}(t), \hat{p}(t), p^0, \hat{u}_1(t)) \\ &= -\left\langle \hat{p}(t), \frac{\partial X}{\partial x}(\hat{x}(t)) \right\rangle - \hat{u}_1(t) \left\langle \hat{p}(t), \frac{\partial Y_1}{\partial x}(\hat{x}(t)) \right\rangle \end{aligned} \quad (3)$$

where the function

$$H(x, p, p^0, u_1) = \langle p, X(x) + u_1 Y_1(x) \rangle + p^0$$

is called the *Hamiltonian*, and the maximization condition

$$H(\hat{x}(t), \hat{p}(t), p^0, \hat{u}_1(t)) = \max_{|w| \leq 1} H(\hat{x}(t), \hat{p}(t), p^0, w) \quad (4)$$

holds almost everywhere on $[0, t_f]$. Moreover, $\max_{|w| \leq 1} H(\hat{x}(t), \hat{p}(t), p^0, w) = 0$ for every $t \in [0, t_f]$. The quadruple $(\hat{x}(\cdot),$

* Corresponding author. Tel.: +33 02 38 49 25 48; fax: +33 02 38 41 72 05.

E-mail addresses: cjoaosilva@ua.pt (C. Silva), emmanuel.trelat@univ-orleans.fr (E. Trélat).

¹ See e.g. [1] for existence results of optimal solutions.

$\hat{p}(\cdot), p^0, \hat{u}_1(\cdot)$ is called an *extremal*. The extremal is said *normal* whenever $p^0 \neq 0$, and in that case it is usual to normalize the adjoint vector so that $p^0 = -1$; otherwise it is said *abnormal*. It follows from (4) that

$$\hat{u}_1(t) = \text{sign}(\hat{p}(t), Y_1(\hat{x}(t))) \quad (5)$$

for almost every t , provided that the (continuous) switching function

$$\varphi_1(t) = \langle \hat{p}(t), Y_1(\hat{x}(t)) \rangle$$

does not vanish on any subinterval of $[0, t_f]$.² Such an extremal $(\hat{x}(\cdot), \hat{p}(\cdot), p^0, \hat{u}_1(\cdot))$ is then completely determined by the initial adjoint vector $\hat{p}(0)$. This extremal is a priori defined on the time interval $[0, t_f]$, but since it is completely determined by the differential system (1)–(3) and its initial condition, it may be extended forward on a maximal time interval I of $[0, +\infty)$, containing $[0, t_f]$. In this way, we consider the trajectory $\hat{x}(\cdot)$ on this maximal interval I .

Note that, since $\hat{x}(\cdot)$ is optimal on $[0, t_f]$, and since the control system under study is autonomous, it follows that $\hat{x}(\cdot)$ is as well optimal for the problem of steering the system (1) from $\hat{x}(0) = \hat{x}_0$ to $\hat{x}(t)$, for every $t \in (0, t_f]$.

We assume that the extremal $(\hat{x}(\cdot), \hat{p}(\cdot), p^0, \hat{u}_1(\cdot))$ is *bang–bang* on the interval I , that is, the switching function φ_1 does not vanish on any subinterval of I . Denote by $\hat{\tau}_1, \dots, \hat{\tau}_s, \dots$ the zeros of φ_1 on I (possibly in infinite number), called switching times, such that $0 < \hat{\tau}_1 < \dots < \hat{\tau}_s < \dots$. As explained above, there holds

$$\hat{u}_1(t) = \begin{cases} 1 & \text{if } \varphi_1(t) > 0, \\ -1 & \text{if } \varphi_1(t) < 0, \end{cases}$$

for every $t \in I$.

We assume moreover that the extremal $(\hat{x}(\cdot), \hat{p}(\cdot), p^0, \hat{u}_1(\cdot))$ satisfies the *strict bang–bang Legendre condition*, that is,

$$\dot{\varphi}_1(\hat{\tau}_j) = \left. \frac{d}{dt} \langle \hat{p}(t), Y_1(\hat{x}(t)) \rangle \right|_{t=\hat{\tau}_j} \neq 0,$$

for every switching time. This condition implies that the switching times are isolated, and moreover are in finite number on every compact subinterval of I . In particular, we assume that there are exactly s switching times on the interval $[0, t_f]$, such that $0 < \hat{\tau}_1 < \dots < \hat{\tau}_s < t_f$.

Definition 1.1. Let $T > 0, T \in I$. The trajectory $\hat{x}(\cdot)$ is said to be *locally minimal time on $[0, T]$ in C^0 topology* if there exist a neighborhood W of the trajectory $\hat{x}(\cdot)$ in \mathbb{R}^n and a real number $\eta > 0$ such that, for every trajectory $y(\cdot)$ that is solution of (1), contained in W , associated with a control v on $[0, T + \eta]$ satisfying the constraint (2), satisfying $y(0) = \hat{x}_0$ and $y(t_1) = \hat{x}(T)$ with $t_1 \in [0, T + \eta]$, there holds $t_1 \geq T$.

The C^0 local optimality is also called *strong local optimality*. The notion of global optimality is defined similarly, with $W = \mathbb{R}^n$ and $\eta = +\infty$.

The Pontryagin maximum principle mentioned formerly is a necessary first order condition for optimality; conversely, extremals are not necessarily locally optimal, and there have been many works on high order necessary optimality conditions (see e.g. [3]) and on sufficient (first and second order) optimality conditions detailed in the next section.

1.2. Second order optimality conditions and conjugate times for bang–bang controls

Consider the extremal $(\hat{x}(\cdot), \hat{p}(\cdot), p^0, \hat{u}_1(\cdot))$ of the problem (OCP) introduced previously.

Definition 1.2. The *cut time* $t_{\text{cut}}(\hat{x}_0)$ is defined as the first positive time of I beyond which the trajectory $\hat{x}(\cdot)$ loses its global optimality status for the problem of steering the system (1)–(2) from \hat{x}_0 to \hat{x}_1 in minimal time, with the agreement that $t_{\text{cut}}(\hat{x}_0) = +\infty$ whenever $\hat{x}(\cdot)$ is globally optimal on every interval $[0, T], T > 0, T \in I$. The point $\hat{x}(t_{\text{cut}}(\hat{x}_0))$ is called a *cut point*.

Whereas such a global optimality status is difficult to characterize, the local optimality status of a trajectory may be characterized using the concept of conjugate time, that is, the time at which the optimal trajectory $\hat{x}(\cdot)$ loses its local optimality. We next recall well known facts on first conjugate times of solutions $\hat{x}(\cdot)$ of the optimal control problem (OCP) associated to bang–bang controls $\hat{u}_1(\cdot)$.

The definition and computation of conjugate points are an important topic in the theory of calculus of variations (see e.g. [4]). In [5] the investigation of the definition and computation of conjugate points for minimal time control problems is based on the study of second order conditions. In fact, second order necessary and/or sufficient conditions are crucial for the study of the first conjugate time of the problem (OCP). In [6], the theory of envelopes and conjugate points is used for the study of the structure of locally optimal bang–bang trajectories for the problem (OCP) in \mathbb{R}^2 and \mathbb{R}^3 ; these results were generalized in [7].

Second order optimality conditions. When the optimal control problem has a nonlinear control system and the extremal controls are continuous, the literature on first and/or second order sufficient conditions is vast; see e.g. [8–16] and references therein. In this case numerical procedures are available to test second order sufficient conditions; see e.g. [17,18,12]. For second order necessary and/or sufficient conditions of optimal control problems with nonlinear control systems and discontinuous controls; see e.g. [19] and references therein.

We will next focus on second order necessary and/or sufficient optimality conditions for optimal control problems with affine-control systems and bang–bang optimal controls.

In [20] a minimal time control problem for affine-control systems is considered and first and second order sufficient optimality conditions are derived, for bang–bang Pontryagin extremal controls which are L^1 -locally optimal. In [21] the same optimal control problem is studied and the authors provide sufficient conditions for strong local optimality and develop numerical methods to test the positive definiteness of a specific quadratic form. In both papers [20] and [21], the sufficient optimality conditions are expressed in terms of quadratic forms, however although the same critical subspace is used, the quadratic form in [20] is a lower bound for the one in [21]. In fact, the second order sufficient optimality condition in [21] is always fulfilled whenever the corresponding condition in [20] is.

In [22,21] optimization methods are given to test second order sufficient optimality conditions for optimal control problems with bounded scalar controls [21], and vector-valued controls [22].

In [23] the authors derive second order sufficient conditions, under the same regularity assumptions as [21], for an optimal control problem in the Mayer form with fixed final time, with affine-control systems and bang–bang optimal controls. In [24] the authors showed that, in certain cases, the second order sufficient conditions given in [21] are equivalent to the ones in [23]. In the cases where the equivalence holds, the results obtained in [24] extend those in [23] to the problem of free final time, with mixed initial and terminal conditions of equality and inequality type. The

² The case where the switching function may vanish on a subinterval is related to singular trajectories, and is outside the scope of this article where we focus on the bang–bang case.

detailed proofs of the main results in [24] are given in [25]. In [23] a finite-dimensional subproblem is considered which consists in moving the switching times and a second variation is defined as a certain quadratic form associated to this subproblem; then, finding a conjugate time consists in testing the positivity of that quadratic form. The authors prove that this can only happen at a switching time.

In [26] the minimal time problem for control-affine systems is studied. An analogous quadratic form to the one in [23] is defined, but the kind of optimality studied is a stronger one (state local optimality).

Quadratic forms. As mentioned above the quadratic forms defined in [23,21] are equivalent (see [24,25]), although the way they are defined is different. We only give a brief sketch of a possible procedure to define the quadratic form.

Let $F(t; \tau_1, \dots, \tau_s) = x(t; \tau_1, \dots, \tau_s)$ be the mapping associated with the finite-dimensional problem associated to (OCP) that consists in moving the switching times τ_1, \dots, τ_s in a neighborhood of the reference switching times $\hat{\tau}_1, \dots, \hat{\tau}_s$ (see [23,22,24–26]), where $x(t; \tau_1, \dots, \tau_s)$ is the trajectory solution of (1), on $[0, t]$, with $x(0) = \hat{x}_0$, associated to the bang–bang control $u_1(\cdot)$ with switching times τ_1, \dots, τ_s and such that it coincides with the reference trajectory $\hat{x}(\cdot)$ whenever $\tau_i = \hat{\tau}_i$ for every i . Note that the trajectory $x(\cdot; \tau_1, \dots, \tau_s)$ is not the projection of an extremal whenever $\tau_i \neq \hat{\tau}_i$. The mapping F is well defined for t in a neighborhood of t_f and τ_i in a neighborhood of $\hat{\tau}_i$ for every i , and is the composition of smooth mappings, therefore is differentiable. Denoting $\tau = (\tau_1, \dots, \tau_s)$, one has

$$\frac{\partial F}{\partial \tau}(t; \tau_1, \dots, \tau_s) = \begin{pmatrix} \frac{\partial x_1}{\partial \tau_1}(\cdot) & \dots & \frac{\partial x_1}{\partial \tau_s}(\cdot) \\ \vdots & \vdots & \vdots \\ \frac{\partial x_n}{\partial \tau_1}(\cdot) & \dots & \frac{\partial x_n}{\partial \tau_s}(\cdot) \end{pmatrix},$$

and

$$\frac{\partial F}{\partial t}(t; \tau_1, \dots, \tau_s) = \dot{x}(t; \tau_1, \dots, \tau_s).$$

Since $\hat{x}(\cdot)$ is optimal, it follows that

$$\text{rank} \left(\frac{\partial F}{\partial \tau}(t; \hat{\tau}_1, \dots, \hat{\tau}_s) \right) \leq n - 1.$$

Indeed, otherwise, if $\text{rank} \left(\frac{\partial F}{\partial \tau}(t; \hat{\tau}_1, \dots, \hat{\tau}_s) \right) = n$ then F would be a local submersion, which contradicts the optimality of $\hat{x}(\cdot)$. Therefore, there exists a multiplier $\psi_t \in \mathbb{R}^n \setminus \{0\}$ such that $\psi_t \cdot \frac{\partial F}{\partial \tau}(t; \hat{\tau}_1, \dots, \hat{\tau}_s) = 0$. Denote by Q_t the intrinsic second derivative of the mapping F , defined by

$$Q_t = \psi_t \cdot \frac{\partial^2 F}{\partial \tau^2}(t; \hat{\tau}_1, \dots, \hat{\tau}_s) \Big|_{\ker \frac{\partial F}{\partial \tau}(t; \hat{\tau}_1, \dots, \hat{\tau}_s)}. \quad (6)$$

Explicit formulas of Q_t are given in [23,27,21,26]; in particular formulas in terms of Lie brackets of the vector fields can be derived.

The next theorem, combination of several known results, provides a necessary and/or sufficient condition for strong local optimality.

Theorem 1 ([23,27,21,28,26]). *Let $(\hat{x}(\cdot), \hat{p}(\cdot), p^0, \hat{u}_1(\cdot))$ be a bang–bang extremal for (OCP) defined on a maximal time interval I of $[0, +\infty)$ containing $[0, t_f]$. If this extremal satisfies the strict bang–bang Legendre condition on I , then for every $t \in I$, the following holds:*

- If the quadratic form Q_t is positive definite then $\hat{x}(\cdot)$ is a local minimizer in the C^0 topology on $[0, t]$.

- Assume moreover that $\hat{x}(\cdot)$ has a unique extremal lift (up to a multiplicative scalar) $(\hat{x}(\cdot), \hat{p}(\cdot), p^0, \hat{u}_1(\cdot))$, which is moreover normal ($p^0 = -1$). If $\hat{x}(\cdot)$ is locally optimal in the C^0 topology on $[0, t]$ then Q_t is nonnegative.

Remark 1.1. Under the assumptions of Theorem 1, the set

$$\{t > 0 | Q_t \text{ has a nontrivial kernel}\}$$

is discrete and can only consist of some switching times (see [23]). This remark permits to define the notion of first conjugate time.

Definition 1.3. The first conjugate time t_c of $\hat{x}(\cdot)$ is defined by

$$t_c = \sup\{t | Q_t \text{ is positive definite}\} = \inf\{t | Q_t \text{ is indefinite}\}.$$

The point $\hat{x}(t_c)$ is called the first conjugate point of the trajectory $\hat{x}(\cdot)$.

Remark 1.2. A conjugate time can only occur at a switching time.

Extremal field approach. Sufficient optimality conditions for a general optimal control problem are provided in [28] (see also [23,26]) with a different point of view than the one recalled in the previous paragraph. In [28] the authors study local optimality conditions for both continuous and piecewise continuous (including bang–bang) controls. The sufficient conditions developed in that article are based on the method of characteristics and the theory of extremal fields. Sufficient optimality conditions are given for embedding a reference trajectory into a local field of broken extremals.³ The occurrence of a conjugate point is related with a so-called overlap of the flow near the switching surface. Second order sufficient optimality conditions stated in [28] have been tested numerically for bang–bang control problems; see e.g. [29]. See also [30] where sufficient optimality conditions for bang–bang controls based on the extremal field approach are studied.

In [31,32,23], using Hamiltonian methods and the extremal field theory, the authors construct, under certain conditions, a non-intersecting field of state extremals⁴ that covers a given extremal trajectory $\hat{x}(\cdot)$. In [23] the authors associate the occurrence of a conjugate point with a fold point of the flow of the extremal field. We next recall the Hamiltonian approach presented in [23,26].

For every $z_0 = (x_0, p_0) \in \mathbb{R}^n \times \mathbb{R}^n$, let $z(\cdot, z_0) = (x(\cdot, z_0), p(\cdot, z_0))$ denote the solution of the system of Eqs. (1) and (3), with the control (5), such that $z(0, z_0) = z_0$. The exponential mapping is then defined by

$$\exp(t, z_0) = x(t, z_0).$$

In (OCP) as in the problems considered in [23] and [26] the initial point is not free (\hat{x}_0 is a fixed point of \mathbb{R}^n). To apply the Hamiltonian approach presented in [23,26], we consider a C^2 function $\alpha : \mathbb{R}^n \rightarrow \mathbb{R}$ such that $\alpha'(\hat{x}_0) = \hat{p}_0$, where $\alpha'(x_0)$ denotes $\frac{\partial \alpha}{\partial x}(x_0)$ and $\hat{p}_0 = \hat{p}(0)$. The function α represents a penalization on the initial point \hat{x}_0 and a new finite-dimensional subproblem is considered, with free initial point $\alpha(\hat{x}_0)$, that consists in moving the switching times and minimizing $\alpha(\hat{x}_0) + t_f$.

The existence of a function α in the previous conditions was proved in [33]. Moreover, in [26] the authors proved that if the quadratic form (6) is positive definite, then the quadratic form associated to the finite-dimensional subproblem of moving the switching times with free initial point is also positive definite.

³ Broken extremals are associated to piecewise continuous controls.

⁴ By non-intersecting extremals we mean that for any fixed $t \in (0, t_c)$ and any extremal trajectories $x(\cdot), y(\cdot)$ with initial points x_0, y_0 , respectively, with x_0, y_0 close to \hat{x}_0 , we have $x(t) \neq y(t)$.

Let \mathcal{O} be a neighborhood of the initial point \hat{x}_0 . Let $x_0 \in \mathcal{O}$; define the switching time functions $\tau_j : \mathcal{O} \rightarrow \mathbb{R}$ with

$$\tau_0(x_0) = 0 \quad \text{and} \quad \tau_j(\hat{x}_0) = \hat{\tau}_j, \quad j = 1, \dots, s,$$

such that, for $j = 1, \dots, s$,

$$\varphi_1(\tau_j(x_0)) = \langle p(\tau_j(x_0), x_0, \alpha'(x_0)), Y_1(x(\tau_j(x_0), x_0, \alpha'(x_0))) \rangle = 0.$$

In other words, $\tau_j(x_0)$ is the j th-switching time of the extremal $x(\cdot, x_0, \alpha'(x_0))$, $p(\cdot, x_0, \alpha'(x_0))$ starting from $(x_0, \alpha'(x_0))$, with x_0 close to \hat{x}_0 .

Since $\hat{x}(\cdot)$ is a minimal time trajectory, there holds $\max_{|w| \leq 1} H(\hat{x}_0, \hat{p}_0, p^0, w) = 0$. Consider the set

$$X = \{x_0 \in \mathcal{O} \mid \max_{|w| \leq 1} H(x_0, \alpha'(x_0), p^0, w) = 0\}.$$

We claim that X is a $(n - 1)$ -dimensional manifold.⁵ Indeed, consider the map

$$G : \mathcal{O} \rightarrow \mathbb{R}$$

$$x_0 \mapsto G(x_0) = \max_{|w| \leq 1} H(x_0, \alpha'(x_0), p^0, w)$$

and the vector field $h_1(x_0) = X(x_0) + u_1 Y_1(x_0)$ that defines the extremal trajectory $x(\cdot)$ on the interval $[0, \tau_1(x_0))$, associated to the value u_1 that satisfies the maximization condition (4) on the referred interval. Proving that X is a $(n - 1)$ -dimensional manifold amounts to proving that, for every function $\alpha \in C^2$ such that $\alpha'(x_0) = p_0$, there holds $dG(x_0) \neq 0$ before the first conjugate time t_c . The second variation formula given in [26, p. 275, Eq. (12)] taken at $(\delta x, \varepsilon) = (h_1(x_0), -1, 0, \dots, 0)$ is equal to, after some simplifications, $dG(x_0) \cdot h_1(x_0)$. Since the second variation is positive definite on $(0, t_c)$ then $dG(x_0) \cdot h_1(x_0) \neq 0$ before t_c . The claim is proved.

Define the j th switching surface Σ_j , for $j = 1, \dots, s$, as the image of the mapping

$$x_0 \mapsto \exp(\tau_j(x_0), x_0, \alpha'(x_0)),$$

where $x_0 \in X$.

Remark 1.3. If the strict bang–bang Legendre condition holds, then the flow associated to the maximized Hamiltonian crosses the switching surface Σ_j at the instant $\hat{\tau}_j$ transversally, for $j = 1, \dots, s$ (see [23]).

Theorem 2 ([23,34,21,28]). Let $(\hat{x}(\cdot), \hat{p}(\cdot), p^0, \hat{u}_1(\cdot))$ be a bang–bang extremal for (OCP) that satisfies the strict bang–bang Legendre condition on $[0, t_c)$, with $t_c < +\infty$. The trajectory $\hat{x}(\cdot)$ is strong locally optimal if and only if there exists a function $\alpha \in C^2$ with $\alpha'(\hat{x}_0) = \hat{p}_0$ such that:

- the trajectory $\hat{x}(\cdot)$ can be embedded into the field of non-intersecting (broken) extremals $(t, x_0) \mapsto \exp(t, x_0, \alpha'(x_0))$ where $x_0 \in \mathcal{O}$;
- this field of extremals crosses the switching surfaces Σ_j transversally, for $j = 1, \dots, s$, and for $j = 1, \dots, s + 1$, with $\tau_{s+1}(\hat{x}_0) = t_c$, the mapping

$$\begin{aligned} (\tau_{j-1}(x_0), \tau_j(x_0)) \times X &\longrightarrow \mathbb{R}^n \\ (t, x_0) &\longmapsto \exp(t, x_0, \alpha'(x_0)) \end{aligned}$$

is of rank n .

Remark 1.4. In the conditions of Theorem 2, at the first conjugate point $\hat{x}(t_c)$, the flow of the extremal field reflects off the switching surface, causing an overlap of the flow near this surface (see Fig. 1 – switching surface Σ_{s+1} , and see [29,28]).




Fig. 1. Field of extremals.

Remark 1.5. Let $f_j(x_0) = X(x_0) + u_j Y_1(x_0)$, for $j = 1, \dots, s + 2$ and $x_0 \in \mathcal{O}$, be the vector fields that define the extremal trajectory $x(\cdot)$ on $(\tau_{j-1}(x_0), \tau_j(x_0))$, with $\tau_{s+1}(\hat{x}_0) = t_c$ and where u_j is the value (1 or -1) of the control that satisfies the maximization condition (4) in each respective interval. If we take $x_0 \in X$ and $j = 1, \dots, s + 1$, then for $(t, x_0) \in (\tau_{j-1}, \tau_j) \times X$

$$\det(\exp(t, x_0, \alpha'(x_0)), f_j(x_0))$$

has constant sign (see [26]).

Moreover, the determinants

$$\det \left(\frac{d}{dx_0} \exp(t, x_0, \alpha'(x_0)) \Big|_{(t, x_0) \in (\tau_s(x_0), \tau_{s+1}(x_0)) \times X}, f_{s+1}(x_0) \right)$$

and

$$\det \left(\frac{d}{dx_0} \exp(t, x_0, \alpha'(x_0)) \Big|_{(t, x_0) \in (\tau_{s+1}(x_0), \tau_{s+2}(x_0)) \times X}, f_{s+2}(x_0) \right)$$

have different signs (see [26]).

The computation of conjugate times in the bang–bang case is difficult in practice. In the last few years several methods have been developed for the numerical implementation of second order sufficient optimality conditions (see, e.g., [22,21] and references cited therein). These numerical procedures allow the computation of the first conjugate time, for bang–bang optimal control problems with affine-control systems, whenever it exists and is attained at a j th switching time. Besides, in the smooth case, efficient tools are available; see e.g. [35]. We next propose a regularization procedure which allows the use of these tools for the computation of the first conjugate time for the problem (OCP). However, in practice, if j is too large then the numerical computation of the first conjugate time may become very difficult either using the methods for bang–bang or smooth controls.

1.3. The regularization procedure

Let ε be a positive real parameter and let Y_2, \dots, Y_m be $m - 1$ arbitrary smooth vector fields on \mathbb{R}^n , where $m \geq 2$ is an integer. Consider the control-affine system

$$\dot{x}^\varepsilon(t) = X(x^\varepsilon(t)) + u_1^\varepsilon(t) Y_1(x^\varepsilon(t)) + \varepsilon \sum_{i=2}^m u_i^\varepsilon(t) Y_i(x^\varepsilon(t)), \quad (7)$$

where the control $u^\varepsilon(t) = (u_1^\varepsilon(t), \dots, u_m^\varepsilon(t))$ satisfies the constraint

$$\sum_{i=1}^m (u_i^\varepsilon(t))^2 \leq 1. \quad (8)$$

Consider the optimal control problem (OCP) $_\varepsilon$ of determining a trajectory $x^\varepsilon(\cdot)$, solution of (7)–(8) on $[0, t_f^\varepsilon]$, such that $x^\varepsilon(0) = \hat{x}_0$ and $x^\varepsilon(t_f^\varepsilon) = \hat{x}_1$, and minimizing the time of transfer t_f^ε . The

⁵ The argument that follows is due to L. Poggiolini.

parameter ε is viewed as a penalization parameter. The existence of at least one solution for $(\mathbf{OCP})_\varepsilon$ is proved in [36], and it is also proved that any solution $x^\varepsilon(\cdot)$ of $(\mathbf{OCP})_\varepsilon$ converges uniformly on $[0, t_f]$ to a solution $x(\cdot)$ of (\mathbf{OCP}) when ε tends to zero (see Theorem 4).

According to the Pontryagin maximum principle, any optimal solution $\hat{x}^\varepsilon(\cdot)$ of $(\mathbf{OCP})_\varepsilon$, associated with controls $(\hat{u}_1^\varepsilon, \dots, \hat{u}_m^\varepsilon)$ satisfying the constraint (8), is the projection of an extremal $(\hat{x}^\varepsilon(\cdot), \hat{p}^\varepsilon(\cdot), p^{0\varepsilon}, \hat{u}^\varepsilon(\cdot))$ such that

$$\begin{aligned} \dot{\hat{p}}^\varepsilon(t) &= -\frac{\partial H^\varepsilon}{\partial x}(\hat{x}^\varepsilon(t), \hat{p}^\varepsilon(t), p^{0\varepsilon}, \hat{u}^\varepsilon(t)) \\ &= -\left\langle \hat{p}^\varepsilon(t), \frac{\partial X}{\partial x}(\hat{x}^\varepsilon(t)) \right\rangle - \hat{u}_1^\varepsilon(t) \left\langle \hat{p}^\varepsilon(t), \frac{\partial Y_1}{\partial x}(\hat{x}^\varepsilon(t)) \right\rangle \\ &\quad - \varepsilon \sum_{i=2}^m \hat{u}_i^\varepsilon(t) \left\langle \hat{p}^\varepsilon(t), \frac{\partial Y_i}{\partial x}(\hat{x}^\varepsilon(t)) \right\rangle \end{aligned} \quad (9)$$

where $H^\varepsilon(x^\varepsilon, p^\varepsilon, p^{0\varepsilon}, u^\varepsilon) = \langle p^\varepsilon, X(x^\varepsilon) + u_1^\varepsilon Y_1(x^\varepsilon) + \varepsilon \sum_{i=2}^m u_i^\varepsilon Y_i(x^\varepsilon) \rangle + p^{0\varepsilon}$ is the Hamiltonian, and

$$\begin{aligned} H^\varepsilon(\hat{x}^\varepsilon(t), \hat{p}^\varepsilon(t), p^{0\varepsilon}, \hat{u}^\varepsilon(t)) \\ = \max_{\sum_{i=1}^m w_i^2 \leq 1} H^\varepsilon(\hat{x}^\varepsilon(t), \hat{p}^\varepsilon(t), p^{0\varepsilon}, w) \end{aligned} \quad (10)$$

almost everywhere on $[0, t_f^\varepsilon]$. Moreover, the maximized Hamiltonian is equal to 0 on $[0, t_f^\varepsilon]$. The maximization condition (10) turns into

$$\begin{aligned} \hat{u}_1^\varepsilon(t) \langle \hat{p}^\varepsilon(t), Y_1(\hat{x}^\varepsilon(t)) \rangle + \varepsilon \sum_{i=2}^m \hat{u}_i^\varepsilon(t) \langle \hat{p}^\varepsilon(t), Y_i(\hat{x}^\varepsilon(t)) \rangle \\ = \max_{\sum_{i=1}^m w_i^2 \leq 1} \left(w_1 \langle \hat{p}^\varepsilon(t), Y_1(\hat{x}^\varepsilon(t)) \rangle + \varepsilon \sum_{i=2}^m w_i \langle \hat{p}^\varepsilon(t), Y_i(\hat{x}^\varepsilon(t)) \rangle \right), \end{aligned} \quad (11)$$

and two cases may occur: either the maximum is attained in the interior of the domain, or it is attained at the boundary. In the first case, there must hold $\langle p^\varepsilon(t), Y_i(x^\varepsilon(t)) \rangle = 0$, for every $i \in \{1, \dots, m\}$; in particular, if the m functions $t \mapsto \langle p^\varepsilon(t), Y_i(x^\varepsilon(t)) \rangle$, $i = 1, \dots, m$, do not vanish simultaneously, then the maximum is attained on the boundary of the domain.

We make the following assumption.

Assumption 3. The integer m and the vector fields Y_2, \dots, Y_m are chosen such that

$$\text{Span}\{Y_i | i = 1, \dots, m\} = \mathbb{R}^n.$$

Under the Assumption 3, the maximization condition (11) yields, for $i = 2, \dots, m$,

$$\begin{aligned} \hat{u}_1^\varepsilon(t) &= \frac{\langle \hat{p}^\varepsilon(t), Y_1(\hat{x}^\varepsilon(t)) \rangle}{\sqrt{\langle \hat{p}^\varepsilon(t), Y_1(\hat{x}^\varepsilon(t)) \rangle^2 + \varepsilon^2 \sum_{i=2}^m \langle \hat{p}^\varepsilon(t), Y_i(\hat{x}^\varepsilon(t)) \rangle^2}}, \\ \hat{u}_i^\varepsilon(t) &= \frac{\varepsilon \langle \hat{p}^\varepsilon(t), Y_i(\hat{x}^\varepsilon(t)) \rangle}{\sqrt{\langle \hat{p}^\varepsilon(t), Y_1(\hat{x}^\varepsilon(t)) \rangle^2 + \varepsilon^2 \sum_{i=2}^m \langle \hat{p}^\varepsilon(t), Y_i(\hat{x}^\varepsilon(t)) \rangle^2}}, \end{aligned} \quad (12)$$

for almost every $t \in [0, t_f^\varepsilon]$, and moreover the control functions $\hat{u}_i^\varepsilon(\cdot)$, $i = 1, \dots, m$ are smooth functions of t (so that the above formula holds actually for every $t \in [0, t_f^\varepsilon]$).

Note that this assumption implies that $m \geq n$. Actually, this assumption can be weakened (see [36] for details).

Theorem 4 ([36]). Assume that the problem (\mathbf{OCP}) has a unique solution $\hat{x}(\cdot)$, defined on $[0, t_f]$, associated with a bang–bang control $\hat{u}_1(\cdot)$ on $[0, t_f]$. Moreover, assume that $\hat{x}(\cdot)$ has a unique extremal lift (up to a multiplicative scalar), which is moreover normal, denoted $(\hat{x}(\cdot), \hat{p}(\cdot), -1, \hat{u}_1(\cdot))$.

Then, under the Assumption 3, there exists $\varepsilon_0 > 0$ such that, for every $\varepsilon \in (0, \varepsilon_0)$, the problem $(\mathbf{OCP})_\varepsilon$ has at least one solution $\hat{x}^\varepsilon(\cdot)$, defined on $[0, t_f^\varepsilon]$ with $t_f^\varepsilon \leq t_f$, associated with a smooth control $u^\varepsilon(\cdot) = (u_1^\varepsilon(\cdot), \dots, u_m^\varepsilon(\cdot))$ satisfying the constraint (8), every extremal lift of which is normal. Let $(\hat{x}^\varepsilon(\cdot), \hat{p}^\varepsilon(\cdot), -1, \hat{u}^\varepsilon(\cdot))$ be such a normal extremal lift. Then, as ε tends to 0,

- t_f^ε converges to t_f ;
- $\hat{x}^\varepsilon(\cdot)$ converges uniformly⁶ to $\hat{x}(\cdot)$, and $\hat{p}^\varepsilon(\cdot)$ converges uniformly to $\hat{p}(\cdot)$ on $[0, t_f]$;
- $\hat{u}_i^\varepsilon(\cdot)$ converges to $\hat{u}_i(\cdot)$ and $u_i^\varepsilon(\cdot)$, $i = 2, \dots, m$, converge to 0 almost everywhere on $[0, t_f]$, and thus in particular for the strong $L^1(0, t_f)$ topology.

Remark 1.6. This result remains true if we extend forward the interval $[0, t_f]$ on an interval $[0, T]$ for $T \in I$, where I is a maximal time interval of $[0, +\infty)$ containing $[0, t_f]$.

Remark 1.7. It is assumed that the problem (\mathbf{OCP}) has a unique solution $\hat{x}(\cdot)$, having a unique extremal lift that is normal. Such an assumption holds true whenever the minimum time function (the value function of the optimal control problem) enjoys differentiability properties (see e.g. [37,38] for a precise relationship, see also [39–42] for results on the size of the set where the value function is differentiable).

1.4. Conjugate times in the smooth case

We recall how to define the concept of first conjugate time for the smooth optimal control problem $(\mathbf{OCP})_\varepsilon$. A first possible definition of conjugate times is in terms of a quadratic form, which is the second order intrinsic derivative of the end-point mapping defined by $E(\varepsilon, t_f^\varepsilon, \hat{x}_0, u^\varepsilon) = x^\varepsilon(t_f^\varepsilon)$ where $t \mapsto x^\varepsilon(\varepsilon, t, \hat{x}_0, u^\varepsilon)$ is the trajectory solution of (7), associated to the control u^ε , such that $x^\varepsilon(\varepsilon, 0, \hat{x}_0, u^\varepsilon) = \hat{x}_0$. Testing a conjugate time amounts to testing the positivity of that quadratic form. However, this definition requires a corank one assumption, and we will rather use a geometric concept of conjugate time, defined below. We refer the reader to [35] for a survey on that theory and to [43] for extensive explanations and for the more general Morse index theory.

Geometric conjugate time.

Definition 1.4. Let $x_0 \in \mathcal{O}$. The point $x^\varepsilon(t_c^\varepsilon)$ is geometrically conjugate to $x^\varepsilon(0)$ if and only if the mapping $x_0 \mapsto \exp^\varepsilon(t_c^\varepsilon, x_0, \alpha'(x_0))$ is not immersive, that is,

$$\det \left(\frac{d}{dx_0} \exp^\varepsilon(t_c^\varepsilon, x_0, \alpha'(x_0)) \right) = 0.$$

The time t_c^ε is called a *geometric conjugate time*.

Remark 1.8. Given an extremal $(\hat{x}^\varepsilon(\cdot), \hat{p}^\varepsilon(\cdot), p^{0\varepsilon}, u^\varepsilon(\cdot))$, the notion of geometric conjugate time coincides with the notion of conjugate time defined in terms of quadratic form, provided the following assumptions hold:

⁶ We consider any continuous extension of the extremal $(\hat{x}^\varepsilon(\cdot), \hat{p}^\varepsilon(\cdot), -1, \hat{u}_1^\varepsilon(\cdot))$ on $[0, t_f]$.

- the strong Legendre condition holds along the extremal, that is, there exists $\gamma > 0$ such that

$$\frac{\partial^2 H}{\partial u^2}(\hat{x}^\varepsilon(\cdot), \hat{p}^\varepsilon(\cdot), p^{0\varepsilon}, u_1^\varepsilon(\cdot)) \cdot (v, v) \leq -\gamma \|v\|^2,$$

for every $v \in \mathbb{R}^m$;

- the control u^ε is of corank one on every subinterval (assumption of strong regularity, see [5]).

Moreover, in that case the first conjugate time t_c^ε characterizes the optimality status of the extremal: the trajectory $\hat{x}^\varepsilon(\cdot)$ is strongly locally optimal on $[0, t]$, for every $t < t_c^\varepsilon$; for $t > t_c^\varepsilon$, the trajectory $\hat{x}^\varepsilon(\cdot)$ is not locally optimal on $[0, t]$ (see, e.g., [43,35,5]).

Remark 1.9. None of the two assumptions of the previous remark will be made for the extremal $(\hat{x}^\varepsilon(\cdot), \hat{p}^\varepsilon(\cdot), p^{0\varepsilon}, \hat{u}^\varepsilon(\cdot))$. In fact, our aim is to prove that the first geometric conjugate time t_c^ε converges to the first conjugate time t_c of the bang–bang case, when ε tends to 0. This result, derived in Theorem 6 (Section 1.5), will permit to use as well in the bang–bang case the available efficient implementation procedures that exist in the smooth case, like for instance the free package *COTCOT*⁷ (see [35]).

For normal extremals $(x^\varepsilon(\cdot), p^\varepsilon(\cdot), -1, u^\varepsilon(\cdot))$ that satisfy the strong Legendre condition, the absence of conjugate points is a sufficient condition for local optimality (see e.g. [43]). This sufficient optimality condition will be expressed using the extremal field approach.

Extremal field approach. From Theorem 4 every extremal lift of the problem $(\mathbf{OCP})_\varepsilon$ is normal ($p^{0\varepsilon} = -1$). Analogously to the bang–bang case, the aim is to construct a family of extremals containing the reference normal extremal $(\hat{x}^\varepsilon(\cdot), \hat{p}^\varepsilon(\cdot), -1, \hat{u}^\varepsilon(\cdot))$, sharing nice non-intersection properties before the first conjugate time.

For every $z_0 = (x_0, p_0) \in \mathbb{R}^n \times \mathbb{R}^n$, let $z^\varepsilon(\cdot, z_0) = (x^\varepsilon(\cdot, z_0), p^\varepsilon(\cdot, z_0))$ be the solution of the system of Eqs. (7) and (9), with the controls (12), such that $z^\varepsilon(0, z_0) = z_0$. The exponential mapping associated to $(\mathbf{OCP})_\varepsilon$ is defined by

$$\exp^\varepsilon(t, z_0) = x^\varepsilon(t, z_0).$$

Let $x_0 \in \mathcal{O}$ and $\alpha^\varepsilon : \mathbb{R}^n \rightarrow \mathbb{R}$ be a C^2 function such that $\alpha^{\varepsilon'}(x_0) = p^\varepsilon(0)$, and such that the family of functions (α^ε) converges to the function α associated with the problem (\mathbf{OCP}) in C^2 topology, as ε tends to 0. As in the bang–bang case, define

$$X^\varepsilon = \{x_0 \in \mathcal{O} \mid \max_{\sum_{i=1}^m w_i^2 \leq 1} H^\varepsilon(x_0, \alpha^{\varepsilon'}(x_0), -1, w^\varepsilon) = 0\}.$$

For $\varepsilon > 0$ small enough, X^ε is a $(n - 1)$ -dimensional manifold. Indeed, let G^ε be defined on \mathcal{O} by $G^\varepsilon(x_0) = \max_{\sum_{i=1}^m w_i^2 \leq 1} H^\varepsilon(x_0, \alpha^{\varepsilon'}(x_0), -1, w^\varepsilon)$. It follows from Theorem 4 that G^ε converges to G (defined in Section 1.2) as ε goes to 0, and therefore, for $\alpha \in C^2$ such that $\alpha'(x_0) = p_0$, there holds $dG^\varepsilon(x_0) \neq 0$, since $dG(x_0) \neq 0$.

Theorem 5 ([43]). *If the normal extremal $(\hat{x}^\varepsilon(\cdot), \hat{p}^\varepsilon(\cdot), -1, \hat{u}^\varepsilon(\cdot))$ satisfies the strong Legendre condition and, moreover can be embedded into the family of extremals $\exp^\varepsilon(t, x_0, \alpha^{\varepsilon'}(x_0))$ such that the mapping*

$$(0, t_c^\varepsilon) \times X^\varepsilon \rightarrow \mathbb{R}^n \\ (t, x_0) \mapsto \exp^\varepsilon(t, x_0, \alpha^{\varepsilon'}(x_0))$$

is of rank n , then $(\hat{x}^\varepsilon(\cdot), \hat{p}^\varepsilon(\cdot), -1, \hat{u}^\varepsilon(\cdot))$ is a local minimum in C^0 topology for the problem $(\mathbf{OCP})_\varepsilon$.




Fig. 2. Field of extremals in the smooth case.

Remark 1.10. The typical behavior of the flow of the extremal field at the first conjugate point is a fold point (see Fig. 2, and see [43,44]).

Remark 1.11. If one considers $x_0 \in X^\varepsilon$, then $x^\varepsilon(t_c^\varepsilon)$ is geometrically conjugate to $x^\varepsilon(0)$ if and only if

$$\det \left(\frac{d}{dx_0} \exp^\varepsilon(t_c^\varepsilon, x_0, \alpha^{\varepsilon'}(x_0))|_{X^\varepsilon}, f^\varepsilon(x_0) \right) = 0,$$

where $f^\varepsilon(x_0) = X(x^\varepsilon(x_0)) + \varepsilon \sum_{i=1}^m u_i^\varepsilon(x_0, \alpha^{\varepsilon'}(x_0))$ and $u_i^\varepsilon(x_0, \alpha^{\varepsilon'}(x_0))$ are smooth functions that satisfy the maximization condition (11).

Remark 1.12. Note that, as long as the minimum time function is differentiable at the point $\hat{x}^\varepsilon(t)$, the optimal trajectory $\hat{x}^\varepsilon(\cdot)$ can be embedded into a non-intersecting extremal field.

Remark 1.13. To derive a necessary optimality condition, a corank one assumption is required for the extremal $(\hat{x}^\varepsilon(\cdot), \hat{p}^\varepsilon(\cdot), p^{0\varepsilon}, \hat{u}^\varepsilon(\cdot))$ (see [35]).

1.5. Main result

We first recall the context. Let $\hat{x}(\cdot)$ denote the strong locally optimal trajectory of (\mathbf{OCP}) , corresponding to the control \hat{u}_1 on $[0, t_f]$. In particular, t_f is the minimal time so that $\hat{x}(0) = \hat{x}_0$ and $\hat{x}(t_f) = \hat{x}_1$. We extend $\hat{x}(\cdot)$ on a maximal interval $I \subset [0, +\infty)$ containing $[0, t_f]$, and denote by t_c its first conjugate time. For every $\varepsilon > 0$, let $\hat{x}^\varepsilon(\cdot)$ denote an optimal trajectory solution of $(\mathbf{OCP})_\varepsilon$, corresponding to a control $\hat{u}^\varepsilon = (\hat{u}_1^\varepsilon, \dots, \hat{u}_m^\varepsilon)$ on $[0, t_f^\varepsilon]$. In particular, t_f^ε is the minimal time so that $\hat{x}^\varepsilon(0) = \hat{x}_0$ and $\hat{x}^\varepsilon(t_f^\varepsilon) = \hat{x}_1$. We extend $\hat{x}^\varepsilon(\cdot)$ on a maximal interval of $[0, +\infty)$ containing $[0, t_f^\varepsilon]$, and denote by t_c^ε its first geometrically conjugate time.

Theorem 6. *Assume that the problem (\mathbf{OCP}) has a unique solution $\hat{x}(\cdot)$, associated with a bang–bang control $\hat{u}_1(\cdot)$, on a maximal interval I . Moreover, assume that $\hat{x}(\cdot)$ has a unique extremal lift (up to a multiplicative scalar), which is moreover normal, and denoted by $(\hat{x}(\cdot), \hat{p}(\cdot), -1, \hat{u}_1(\cdot))$. If the extremal $(\hat{x}(\cdot), \hat{p}(\cdot), -1, \hat{u}_1(\cdot))$ satisfies, moreover, the strict bang–bang Legendre condition on $[0, t_c]$, then the first geometric conjugate time t_c^ε converges to the first conjugate time t_c as ε tends to 0.*

Remark 1.14. Let t_{cut} denote the cut time along the extremal $(\hat{x}(\cdot), \hat{p}(\cdot), p^0, \hat{u}(\cdot))$. Analogously to the bang–bang case, we can define the cut time $t_{\text{cut}}^\varepsilon$ of the optimal trajectory $\hat{x}^\varepsilon(\cdot)$ for the problem $(\mathbf{OCP})_\varepsilon$ as the first time at which $\hat{x}^\varepsilon(\cdot)$ loses its optimality. We claim that, under the assumptions of Theorem 4, there holds $\limsup_{\varepsilon \rightarrow 0} t_{\text{cut}}^\varepsilon \leq t_{\text{cut}}$.

⁷ Conditions of Order Two, COnjugate Times, <http://apo.enseiht.fr/cotcot/>.

2. Proof of the main result

The next proposition is the key result to derive [Theorem 6](#).

Proposition 7. *Let \mathcal{O} be a neighborhood of \hat{x}_0 and $x_0 \in \mathcal{O}$. The exponential mapping $(t, x_0) \mapsto \exp^\varepsilon(t, x_0, \alpha^{\varepsilon'}(x_0))$ converges to $(t, x_0) \mapsto \exp(t, x_0, \alpha'(x_0))$ piecewise in C^1 topology on $I \times \mathcal{O}$, with $\tau_{s+1}(\hat{x}_0) = t_c$, as ε tends to 0. More precisely, on every compact subinterval of $(\tau_{j-1}(x_0), \tau_j(x_0)) \times \mathcal{O}$, with $(\tau_{j-1}(x_0), \tau_j(x_0)) \subset I$ and $j \in \mathbb{N}$, the mapping $(t, x_0) \mapsto \exp^\varepsilon(t, x_0, \alpha^{\varepsilon'}(x_0))$ converges to $(t, x_0) \mapsto \exp(t, x_0, \alpha'(x_0))$ uniformly in C^1 topology.*

Proof. In what follows, when it is convenient, we simplify the notation and write $\exp(t, x_0)$ or $x(t, x_0)$ (respectively, $\exp^\varepsilon(t, x_0)$ or $x^\varepsilon(t, x_0)$) for $\exp(t, x_0, \alpha'(x_0))$ (respectively, for $\exp^\varepsilon(t, x_0, \alpha^{\varepsilon'}(x_0))$).

Let $\varepsilon > 0$ small enough. For $x_0 \in \mathcal{O}$, consider the function

$$\varphi_1(\varepsilon, t, x_0) = (p(\varepsilon, t, x_0), Y_1(x(\varepsilon, t, x_0))).$$

For $(\varepsilon, t, x_0) = (0, \hat{\tau}_j, x_0)$, by definition of the switching time, one has $\varphi_1(0, \hat{\tau}_j, x_0) = 0$, and by the strict bang–bang Legendre condition, $\frac{\partial \varphi_1}{\partial t}(0, \hat{\tau}_j, x_0) \neq 0$. By the implicit function theorem there exists a neighborhood $(-\varepsilon_0, \varepsilon_0)$ of $0 \in \mathbb{R}$, such that for $\varepsilon \in (-\varepsilon_0, \varepsilon_0)$, there exists a C^1 function $\tau_j^\varepsilon(x_0) = \tau_j^\varepsilon(\varepsilon, x_0)$, with $j = 1, \dots, s$, satisfying $\varphi_1(\varepsilon, \tau_j^\varepsilon(x_0)) = 0$ and such that, as ε tends to 0, $\tau_j^\varepsilon(x_0)$ converges to $\tau_j(x_0)$, and $\frac{\partial \tau_j^\varepsilon}{\partial x_0}(x_0)$ converges to $\frac{\partial \tau_j}{\partial x_0}(x_0)$.

Analogously to the definition of switching time function of an extremal trajectory $x(\cdot)$, we have thus defined some functions $\tau_j^\varepsilon(\cdot) : \mathcal{O} \rightarrow \mathbb{R}$, that are however not switching functions.

Lemma 2.1. *The mapping $(t, x_0) \mapsto \exp^\varepsilon(t, x_0, \alpha^{\varepsilon'}(x_0))$ converges to $(t, x_0) \mapsto \exp(t, x_0, \alpha'(x_0))$ uniformly in C^1 topology on $J \times \mathcal{O}$, where J is any compact subinterval of $[0, \tau_1(x_0))$, as ε tends to 0.*

Proof. Let J be a compact subinterval of $[0, \tau_1(x_0))$. The uniform C^0 convergence on $J \times \mathcal{O}$ of the mapping $(t, x_0) \mapsto \exp^\varepsilon(t, x_0)$ to $(t, x_0) \mapsto \exp(t, x_0)$, as ε tends to 0, is a direct consequence of [Theorem 4](#). We have

$$\frac{\partial \exp^\varepsilon}{\partial t}(t, x_0) = \dot{x}^\varepsilon(t, x_0)$$

where $\dot{x}^\varepsilon(t, x_0)$ is given by (7); from [Theorem 4](#), $\dot{x}^\varepsilon(t, x_0)$ converges to $\dot{x}(t, x_0) = \frac{d \exp}{dt}(t, x_0)$ as ε tends to 0. On the other hand,

$$\begin{aligned} \frac{d}{dx_0} \exp^\varepsilon(t, x_0, \alpha^{\varepsilon'}(x_0)) &= \frac{\partial \exp^\varepsilon}{\partial x_0}(t, x_0, \alpha^{\varepsilon'}(x_0)) \\ &+ \frac{\partial \exp^\varepsilon}{\partial p_0}(t, x_0, \alpha^{\varepsilon'}(x_0)) \alpha^{\varepsilon''}(x_0) \end{aligned}$$

where $\frac{\partial \exp^\varepsilon}{\partial x_0}(t, x_0, \alpha^{\varepsilon'}(x_0))$ and $\frac{\partial \exp^\varepsilon}{\partial p_0}(t, x_0, \alpha^{\varepsilon'}(x_0))$ are solutions of the linearized system associated with the Hamiltonian system (7)–(9), for the problem $(\mathbf{OCP})_\varepsilon$ on $[0, t]$. From [Theorem 4](#), $(x^\varepsilon(\cdot), p^\varepsilon(\cdot))$ converges uniformly to the solution of the Hamiltonian system associated with the problem (\mathbf{OCP}) as ε tends to 0. This convergence clearly holds as well for the solutions of the linearized system associated with the Hamiltonian system for $(\mathbf{OCP})_\varepsilon$; therefore, as ε tends to 0, $\frac{\partial \exp^\varepsilon}{\partial x_0}(t, x_0, \alpha^{\varepsilon'}(x_0))$ (respectively, $\frac{\partial \exp^\varepsilon}{\partial p_0}(t, x_0, \alpha^{\varepsilon'}(x_0))$) converges to $\frac{\partial \exp}{\partial x_0}(t, x_0, \alpha'(x_0))$ (respectively, $\frac{\partial \exp}{\partial p_0}(t, x_0, \alpha'(x_0))$) uniformly on $[0, t]$. \square

In what follows, the notation $\tau_j^+(x_0)$ (resp. $\tau_j^-(x_0)$) stands for the right limit (resp. the left limit). For $x_0 \in \mathcal{O}$ and $j = 1, \dots, s$, we call the *jump* of $\frac{\partial \exp}{\partial x_0}(t, x_0)$ at $\tau_j(x_0)$ the difference

$$\frac{\partial \exp}{\partial x_0}(\tau_j^+(x_0), x_0) - \frac{\partial \exp}{\partial x_0}(\tau_j^-(x_0), x_0),$$

which is, according to [28, Eq. 3.10, p. 123] given by

$$\begin{aligned} &\frac{\partial \exp}{\partial x_0}(\tau_j^+(x_0), x_0) - \frac{\partial \exp}{\partial x_0}(\tau_j^-(x_0), x_0) \\ &= (u_1(\tau_j^+(x_0), x_0) - u_1(\tau_j^-(x_0), x_0)) Y_1(x(\tau_1(x_0), x_0)) \frac{\partial \tau_j}{\partial x_0}(x_0) \\ &= (\text{sign}(\varphi_1(\tau_j^+)) - \text{sign}(\varphi_1(\tau_j^-))) Y_1(x(\tau_j(x_0), x_0)) \frac{\partial \tau_j}{\partial x_0}(x_0). \end{aligned} \quad (13)$$

Due to this jump condition one cannot expect to get a C^1 convergence result on the whole interval. We will next estimate the difference

$$\frac{\partial \exp^\varepsilon}{\partial x_0}(\tau_j^\varepsilon(x_0) + \eta, x_0) - \frac{\partial \exp^\varepsilon}{\partial x_0}(\tau_j^\varepsilon(x_0) - \eta, x_0),$$

for $\eta > 0$ small, and show that it converges to (13), whenever ε tends to 0, and then η tends to 0.

Lemma 2.2. *There holds*

$$\begin{aligned} &\lim_{\eta \rightarrow 0} \lim_{\varepsilon \rightarrow 0} \left(\frac{\partial \exp^\varepsilon}{\partial x_0}(\tau_1^\varepsilon(x_0) + \eta, x_0) - \frac{\partial \exp^\varepsilon}{\partial x_0}(\tau_1^\varepsilon(x_0) - \eta, x_0) \right) \\ &= \frac{\partial \exp}{\partial x_0}(\tau_1^+(x_0), x_0) - \frac{\partial \exp}{\partial x_0}(\tau_1^-(x_0), x_0). \end{aligned}$$

Proof. One has

$$\begin{aligned} \frac{\partial}{\partial t} \left(\frac{\partial x^\varepsilon}{\partial x_0}(t, x_0) \right) &= \left(\frac{\partial X}{\partial x_0}(x^\varepsilon(t, x_0)) + u_1^\varepsilon(t, x_0) \frac{\partial Y_1}{\partial x_0}(x^\varepsilon(t, x_0)) \right. \\ &+ \varepsilon \sum_{i=2}^m u_i^\varepsilon(t, x_0) \frac{\partial Y_i}{\partial x_0}(x^\varepsilon(t, x_0)) \left. \right) \frac{\partial x^\varepsilon}{\partial x_0}(t, x_0) \\ &+ Y_1(x^\varepsilon(t, x_0)) \frac{\partial u_1^\varepsilon}{\partial x_0}(t, x_0) + \varepsilon \sum_{i=2}^m Y_i(x^\varepsilon(t, x_0)) \frac{\partial u_i^\varepsilon}{\partial x_0}(t, x_0). \end{aligned}$$

It follows that

$$\begin{aligned} &\frac{\partial x^\varepsilon}{\partial x_0}(\tau_1^\varepsilon(x_0) + \eta, x_0) - \frac{\partial x^\varepsilon}{\partial x_0}(\tau_1^\varepsilon(x_0) - \eta, x_0) \\ &= \int_{\tau_1^\varepsilon(x_0) - \eta}^{\tau_1^\varepsilon(x_0) + \eta} \left(\frac{\partial X}{\partial x_0}(x^\varepsilon(t, x_0)) + u_1^\varepsilon(t, x_0) \frac{\partial Y_1}{\partial x_0}(x^\varepsilon(t, x_0)) \right. \\ &+ \varepsilon \sum_{i=2}^m u_i^\varepsilon(t, x_0) \frac{\partial Y_i}{\partial x_0}(x^\varepsilon(t, x_0)) \left. \right) \frac{\partial x^\varepsilon}{\partial x_0}(t, x_0) dt \\ &+ \int_{\tau_1^\varepsilon(x_0) - \eta}^{\tau_1^\varepsilon(x_0) + \eta} Y_1(x^\varepsilon(t, x_0)) \frac{\partial u_1^\varepsilon}{\partial x_0}(t, x_0) dt \\ &+ \int_{\tau_1^\varepsilon(x_0) - \eta}^{\tau_1^\varepsilon(x_0) + \eta} \varepsilon \sum_{i=2}^m Y_i(x^\varepsilon(t, x_0)) \frac{\partial u_i^\varepsilon}{\partial x_0}(t, x_0) dt. \end{aligned}$$

It is easy to see that the limit when η tends to zero of the limit when ε tends to zero of the first and third term of the right side of the last equation is equal to zero. Only the limit term

$$\lim_{\eta \rightarrow 0} \lim_{\varepsilon \rightarrow 0} \int_{\tau_1^\varepsilon(x_0) - \eta}^{\tau_1^\varepsilon(x_0) + \eta} Y_1(x^\varepsilon(t, x_0)) \frac{\partial u_1^\varepsilon}{\partial x_0}(t, x_0) dt$$

deserves a special attention. Let us denote

$$\varphi_i^\varepsilon(t, x_0) = \langle p^\varepsilon(t, x_0), Y_i(x^\varepsilon(t, x_0)) \rangle, \quad i = 1, \dots, m.$$

From (12), we compute easily

$$\frac{\partial u_1^\varepsilon}{\partial x_0}(t, x_0) = \frac{\varepsilon^2 \left(\frac{\partial \varphi_1^\varepsilon}{\partial x_0}(t, x_0) \sum_{i=2}^m \varphi_i^\varepsilon(t, x_0)^2 - \varphi_1^\varepsilon(t, x_0) \sum_{i=2}^m \varphi_i^\varepsilon(t, x_0) \frac{\partial \varphi_i^\varepsilon}{\partial x_0}(t, x_0) \right)}{\left(\varphi_1^\varepsilon(t, x_0)^2 + \varepsilon^2 \sum_{i=2}^m \varphi_i^\varepsilon(t, x_0)^2 \right)^{3/2}}.$$

We will consider asymptotic expansions of these quantities around $\tau_1^\varepsilon(x_0)$. Since $\varphi_1^\varepsilon(\tau_1^\varepsilon(x_0), x_0) = 0$ for every x_0 , it follows that

$$\frac{\partial \varphi_1^\varepsilon}{\partial x_0}(\tau_1^\varepsilon(x_0), x_0) = -\frac{\partial \varphi_1^\varepsilon}{\partial t}(\tau_1^\varepsilon(x_0), x_0) \frac{\partial \tau_1^\varepsilon}{\partial x_0}(x_0).$$

In what follows, denote $\tau_1^\varepsilon = (\tau_1^\varepsilon(x_0), x_0)$. One has

$$\begin{aligned} & \int_{\tau_1^\varepsilon - \eta}^{\tau_1^\varepsilon + \eta} Y_1^\varepsilon(x^\varepsilon(t, x_0)) \frac{\partial u_1^\varepsilon}{\partial x_0}(t, x_0) dt \\ &= \int_{\tau_1^\varepsilon - \eta}^{\tau_1^\varepsilon + \eta} (Y_1(x^\varepsilon(\tau_1^\varepsilon)) + O(t - \tau_1^\varepsilon)) \\ & \times \left[\frac{\varepsilon^2 \left(\frac{\partial \varphi_1^\varepsilon}{\partial x_0}(\tau_1^\varepsilon) + O(t - \tau_1^\varepsilon) \right) \sum_{i=2}^m (\varphi_i^\varepsilon(\tau_1^\varepsilon) + O(t - \tau_1^\varepsilon))^2}{D} \right. \\ & \left. - \frac{\varepsilon^2 (\varphi_1^\varepsilon(\tau_1^\varepsilon) + O(t - \tau_1^\varepsilon)) \sum_{i=2}^m (\varphi_i^\varepsilon(\tau_1^\varepsilon) + O(t - \tau_1^\varepsilon))}{D} \right. \\ & \left. \times \left(\frac{\partial \varphi_1^\varepsilon}{\partial x_0}(\tau_1^\varepsilon) + O(t - \tau_1^\varepsilon) \right) \right] dt \end{aligned} \quad (14)$$

where

$$D = \left(\left(\frac{\partial \varphi_1^\varepsilon}{\partial t}(\tau_1^\varepsilon)(t - \tau_1^\varepsilon) + o(t - \tau_1^\varepsilon) \right)^2 + \varepsilon^2 \sum_{i=2}^m \left(\varphi_i^\varepsilon(\tau_1^\varepsilon) + \frac{\partial \varphi_i^\varepsilon}{\partial t}(\tau_1^\varepsilon)(t - \tau_1^\varepsilon) + o(t - \tau_1^\varepsilon) \right)^2 \right)^{3/2}$$

and simplifying the expression (14) (the terms of order $O((t - \tau_1^\varepsilon)^k)$ and $o((t - \tau_1^\varepsilon)^l)$, with $k = 2, 3$ and $l = 1, 2, 3$, are omitted) we get

$$\begin{aligned} & \int_{\tau_1^\varepsilon - \eta}^{\tau_1^\varepsilon + \eta} Y_1^\varepsilon(x^\varepsilon(t, x_0)) \frac{\partial u_1^\varepsilon}{\partial x_0}(t, x_0) dt \\ &= \int_{\tau_1^\varepsilon - \eta}^{\tau_1^\varepsilon + \eta} (Y_1(x^\varepsilon(\tau_1^\varepsilon))) \\ & \times \frac{-\varepsilon^2 \frac{\partial \varphi_1^\varepsilon}{\partial t}(\tau_1^\varepsilon) N_1}{\left(\left(\left(\frac{\partial \varphi_1^\varepsilon}{\partial t}(\tau_1^\varepsilon) \right)^2 + \varepsilon^2 N_2 \right) (t - \tau_1^\varepsilon)^2 + \varepsilon^2 N_3 (t - \tau_1^\varepsilon) + \varepsilon^2 N_1 \right)^{3/2}} \\ & \times \frac{\partial \tau_1^\varepsilon}{\partial x_0}(x_0) \\ & + \frac{\varepsilon^2 (M_1 - M_2) O(t - \tau_1^\varepsilon) - \varepsilon^2 \frac{\partial \varphi_1^\varepsilon}{\partial t}(\tau_1^\varepsilon) N_1 \frac{\partial \tau_1^\varepsilon}{\partial x_0}(x_0) O(t - \tau_1^\varepsilon)}{\left(\left(\left(\frac{\partial \varphi_1^\varepsilon}{\partial t}(\tau_1^\varepsilon) \right)^2 + \varepsilon^2 N_2 \right) (t - \tau_1^\varepsilon)^2 + \varepsilon^2 N_3 (t - \tau_1^\varepsilon) + \varepsilon^2 N_1 \right)^{3/2}} dt \end{aligned}$$

where

$$N_1 = \sum_{i=2}^m (\varphi_i^\varepsilon(\tau_1^\varepsilon))^2, \quad N_2 = \sum_{i=2}^m \left(\frac{\partial \varphi_i^\varepsilon}{\partial t}(\tau_1^\varepsilon) \right)^2,$$

$$N_3 = 2 \sum_{i=2}^m \varphi_i^\varepsilon(\tau_1^\varepsilon) \frac{\partial \varphi_i^\varepsilon}{\partial t}(\tau_1^\varepsilon),$$

$$M_1 = 2 \frac{\partial \varphi_1^\varepsilon}{\partial x_0}(\tau_1^\varepsilon) \sum_{i=2}^m \varphi_i^\varepsilon(\tau_1^\varepsilon) + \sum_{i=2}^m (\varphi_i^\varepsilon(\tau_1^\varepsilon))^2,$$

$$M_2 = \sum_{i=2}^m \left(\frac{\partial \varphi_i^\varepsilon}{\partial x_0}(\tau_1^\varepsilon) \right)^2.$$

Notice that the denominator never vanishes, since by **Assumption 3** the functions $(t, x_0) \mapsto \varphi_i(t, x_0)$, $i = 1, \dots, m$ do not vanish simultaneously.

The limit when η tends to zero of the limit when ε tends to zero, of the first and second term of the right side of the last equality are respectively equal to $(\text{sign}(\varphi_1(\tau_1^+)) - \text{sign}(\varphi_1(\tau_1^-))) Y_1(x(\tau_1(x_0), x_0)) \frac{\partial \tau_1}{\partial x_0}(x_0)$ and 0. Since

$$\lim_{\varepsilon \rightarrow 0} \frac{\partial \tau_1^\varepsilon}{\partial x_0}(x_0) = \frac{\partial \tau_1}{\partial x_0}(x_0),$$

it follows that

$$\begin{aligned} & \lim_{\eta \rightarrow 0} \lim_{\varepsilon \rightarrow 0} \left(\frac{\partial x^\varepsilon}{\partial x_0}(\tau_1^\varepsilon(x_0) + \eta, x_0) - \frac{\partial x^\varepsilon}{\partial x_0}(\tau_1^\varepsilon(x_0) - \eta, x_0) \right) \\ &= (\text{sign}(\varphi_1(\tau_1^+)) - \text{sign}(\varphi_1(\tau_1^-))) Y_1(x(\tau_1(x_0), x_0)) \frac{\partial \tau_1}{\partial x_0}(x_0), \end{aligned}$$

and the lemma follows. \square

A similar lemma holds for $\frac{\partial \exp}{\partial p_0}$. This result permits to extend the convergence result beyond the first switching time; the extension of **Lemma 2.1** to every further interval (τ_{j-1}, τ_j) is then straightforward. This proves the proposition. \square

We are now in a position to prove **Theorem 6**. From **Theorem 2**, the trajectory $\hat{x}(\cdot)$ can be embedded into the field of extremals $x_0 \mapsto \exp(t, x_0, \alpha'(x_0))$ with $x_0 \in \mathcal{O}$ and the mapping

$$\begin{aligned} & (0, t_c) \times X \rightarrow \mathbb{R}^n \\ & (t, x_0) \mapsto \exp(t, x_0, \alpha'(x_0)) \end{aligned}$$

is of rank n , where $X = \{x_0 \in \mathcal{O} \mid \max_{|w| \leq 1} H(x_0, \alpha'(x_0), p^0, w) = 0\}$, \mathcal{O} is a neighborhood of \hat{x}_0 , and t_c is the first conjugate time of $\hat{x}(\cdot)$.

From **Remark 1.5**, the determinants

$$\det \left(\frac{d}{dx_0} \exp(t, x_0, \alpha'(x_0)) \Big|_{(t, x_0) \in (\tau_5(x_0), \tau_{5+1}(x_0)) \times X}, f_{s+1}(x_0) \right)$$

and

$$\det \left(\frac{d}{dx_0} \exp(t, x_0, \alpha'(x_0)) \Big|_{(t, x_0) \in (\tau_{5+1}(x_0), \tau_{5+2}(x_0)) \times X}, f_{s+2}(x_0) \right)$$

have different signs, with $\tau_{5+1}(\hat{x}_0) = t_c$.

By **Definition 1.4**, the point $x^\varepsilon(\tau_c^\varepsilon(x_0))$ is geometrically conjugate to $x^\varepsilon(0) = x_0$, with $x_0 \in X^\varepsilon$, if and only if

$$\det \left(\frac{d}{dx_0} \exp^\varepsilon(t, x_0, \alpha^{\varepsilon'}(x_0)), f^\varepsilon(x_0) \Big|_{x_0 \in X^\varepsilon} \right) = 0$$

for $t = \tau_c^\varepsilon(x_0)$. Let $x_0 \in X^\varepsilon$, we have

$$\begin{aligned} & \frac{\partial \exp^\varepsilon}{\partial x_0}(\tau^\varepsilon(x_0), x_0, \alpha^{\varepsilon'}(x_0)) \\ &= \frac{\partial \exp^\varepsilon}{\partial t}(\tau^\varepsilon(x_0), x_0, \alpha^{\varepsilon'}(x_0)) \frac{\partial \tau^\varepsilon}{\partial x_0}(x_0) \\ & + \frac{\partial \exp^\varepsilon}{\partial x_0}(\tau^\varepsilon(x_0), x_0, \alpha^{\varepsilon'}(x_0)) \\ & + \frac{\partial \exp^\varepsilon}{\partial p_0}(\tau^\varepsilon(x_0), x_0, \alpha^{\varepsilon'}(x_0)) \alpha^{\varepsilon''}(x_0). \end{aligned}$$

Since $\frac{\partial \exp^\varepsilon}{\partial t}(\tau^\varepsilon(x_0), x_0, \alpha^{\varepsilon'}(x_0)) = \dot{x}^\varepsilon(x_0) = f^\varepsilon(x_0)$, there holds, clearly,

$$\det \left(\frac{\partial \exp^\varepsilon}{\partial t}(\tau^\varepsilon(x_0), x_0, \alpha^{\varepsilon'}(x_0)) \frac{\partial \tau^\varepsilon}{\partial x_0}(x_0), f^\varepsilon(x_0) \right) = 0,$$

and it thus follows that

$$\begin{aligned} & \det \left(\frac{d}{dx_0} \exp^\varepsilon(\tau^\varepsilon(x_0), x_0, \alpha^{\varepsilon'}(x_0)), f^\varepsilon(x_0) \right) \\ &= \det \left(\frac{\partial \exp^\varepsilon}{\partial x_0}(\tau^\varepsilon(x_0), x_0, \alpha^{\varepsilon'}(x_0)) \right. \\ & \quad \left. + \frac{\partial \exp^\varepsilon}{\partial p_0}(\tau^\varepsilon(x_0), x_0, \alpha^{\varepsilon'}(x_0)) \alpha^{\varepsilon''}(x_0), f^\varepsilon(x_0) \right) \\ &= \det \left(\frac{d}{dx_0} \exp^\varepsilon(t, x_0, \alpha^{\varepsilon'}(x_0)), f^\varepsilon(x_0) \right) \end{aligned}$$

for $t = \tau^\varepsilon(x_0)$. By Proposition 7, on every compact subinterval of $(\tau_{j-1}^\varepsilon(x_0), \tau_j^\varepsilon(x_0))$, the mapping $(t, x_0) \mapsto \exp^\varepsilon(t, x_0, \alpha^{\varepsilon'}(x_0))$ converges to $(t, x_0) \mapsto \exp(t, x_0, \alpha'(x_0))$ uniformly in C^1 topology, therefore the determinants

$$\det \left(\frac{d}{dx_0} \exp^\varepsilon(t, x_0, \alpha^{\varepsilon'}(x_0)) \Big|_{(t, x_0) \in (\tau_s^\varepsilon(x_0), \tau_{s+1}^\varepsilon(x_0)) \times X^\varepsilon}, f^\varepsilon(x_0) \right)$$

and

$$\det \left(\frac{d}{dx_0} \exp^\varepsilon(t, x_0, \alpha^{\varepsilon'}(x_0)) \Big|_{(t, x_0) \in (\tau_{s+1}^\varepsilon(x_0), \tau_{s+2}^\varepsilon(x_0)) \times X^\varepsilon}, f^\varepsilon(x_0) \right)$$

have different signs before and after $\tau_{s+1}^\varepsilon(x_0)$. Therefore, by continuity, the function $t \mapsto \det \left(\frac{d}{dx_0} \exp^\varepsilon(t, x_0, \alpha^{\varepsilon'}(x_0)), f^\varepsilon(x_0) \right)$ vanishes for some time, close to $\tau_{s+1}^\varepsilon(x_0)$. By Definition 1.4, this time $t_c^\varepsilon(x_0)$ is a geometrically conjugate time, and when ε tends to 0, $t_c^\varepsilon(x_0)$ converges to the bang–bang conjugate time $t_c = \tau_{s+1}(\hat{x}_0)$. This ends the proof of the theorem.

3. Examples

In this section we illustrate Theorem 6 with two examples of minimal time control problems.

3.1. First example: the Rayleigh problem

In this section we consider the minimal time control problem for the Rayleigh control system (see e.g. [45,21]),

$$\begin{aligned} \dot{x}_1(t) &= x_2(t), \\ \dot{x}_2(t) &= -x_1(t) + x_2(t)(1.4 - 0.14x_2(t)^2) + u_1(t), \end{aligned} \quad (15)$$

with the control constraint

$$|u_1(\cdot)| \leq 4, \quad (16)$$

and with boundary conditions given by

$$x_1(0) = -4, \quad x_2(0) = -3, \quad x_1(t_f) = x_2(t_f) = 0. \quad (17)$$

According to the Pontryagin maximum principle, any optimal solution $\hat{x}(\cdot)$ of (15)–(17) is the projection of an extremal $(\hat{x}(\cdot), \hat{p}(\cdot), p^0, \hat{u}_1(\cdot))$ such that

$$\begin{aligned} \dot{\hat{p}}_1(t) &= \hat{p}_2(t) \\ \dot{\hat{p}}_2(t) &= -\hat{p}_1(t) - \hat{p}_2(t) (1.4 - 0.42\hat{x}_2(t)^2) \end{aligned} \quad (18)$$

and the maximization condition $\hat{p}_2(t)\hat{u}_1(t) = \max_{|w| \leq 4} (\hat{p}_2(t)w)$ holds almost everywhere on $[0, t_f]$. It is easy to see that $\hat{p}_2(\cdot)$ cannot vanish on some subinterval, and it follows that the




Fig. 3. Optimal trajectory.




Fig. 4. Optimal control.

optimal control $\hat{u}_1(\cdot)$ is bang–bang, equal to $\hat{u}_1(t) = 4 \text{ sign}(\hat{p}_2(t))$. Applying a shooting method to problem (15)–(17) (with $p^0 = -1$), we determine the initial adjoint vector $\hat{p}(0) \simeq (0.53095052; 0.34206485)$, and observe that the trajectory has only one switching time $\hat{\tau}_1 \simeq 0.57613128$ on $[0, t_f]$, that is, $\hat{u}_1(\cdot)$ is given by

$$\hat{u}_1(t) = \begin{cases} +4 & \text{for } 0 \leq t \leq \hat{\tau}_1 \\ -4 & \text{for } \hat{\tau}_1 \leq t \leq t_f, \end{cases}$$

with a final time $t_f \simeq 2.97812917$ (see Figs. 3 and 4). Furthermore, $\hat{x}(\cdot)$ is the unique minimal time solution and has a unique extremal lift (up to a multiplicative scalar), which is moreover normal.

Prolongating the trajectory $\hat{x}(\cdot)$ to the interval $[0, 4]$, we observe a second switching time at $\hat{\tau}_2 \simeq 3.14750955$.

Notice that the second order sufficient conditions of [22,13,34, 21] are satisfied before $\hat{\tau}_2$, confirming the local optimality status of the trajectory, but are no longer satisfied beyond this second switching time; we can thus expect the trajectory not to be locally optimal beyond $\hat{\tau}_2$. To investigate this optimality status we use the extremal field approach.

From Theorem 2 and Remark 1.4, the first conjugate point $\hat{x}(t_c)$ is an overlap point of the extremal field emanating from the horizontal one-dimensional manifold $X = \{x_0 \in \mathcal{O} \mid \max_{|w| \leq 1} H(x_0, \alpha'(x_0), -1, w) = 0\}$. In practice, the function α is not known, and we rather use the field of extremals emanating from the vertical manifold $X_p = \{p_0 \in \mathcal{O}_p \mid$




 Fig. 5. Extremal field $t \in [0, 4]$.


Fig. 6. Overlap of the flow.

$\max_{|w| \leq 1} H(\hat{x}_0, p_0, -1, w) = 0$ (see [35,26]), where \mathcal{O}_p is a neighborhood of the initial value of the adjoint vector $\hat{p}(0)$. The characterization in terms of fold point still holds for this vertical manifold (see [26]). We observe on Figs. 5 and 6 that this field of extremals reflects off the switching surface at the second switching time; the point $\hat{x}(\hat{\tau}_2)$ is a fold point and the first conjugate time is equal to the second switching time, $t_c = \hat{\tau}_2 \simeq 3.14750955$.

We next propose a regularization procedure, for which we compute the first geometric conjugate time t_c^ε and check that it indeed converges to the first conjugate time t_c of the bang–bang case as ε tends to 0.

We consider the regularized control system

$$\begin{aligned} \dot{x}_1^\varepsilon(t) &= x_2^\varepsilon(t) + \varepsilon u_2^\varepsilon(t), \\ \dot{x}_2^\varepsilon(t) &= -x_1^\varepsilon(t) + x_2^\varepsilon(t)(1.4 - 0.14x_2^\varepsilon(t)^2) + u_1^\varepsilon(t), \end{aligned} \quad (19)$$

with the boundary conditions (17), and where the control $u^\varepsilon(\cdot) = (u_1^\varepsilon(\cdot), u_2^\varepsilon(\cdot))$ satisfies the constraint

$$(u_1^\varepsilon(\cdot))^2 + (u_2^\varepsilon(\cdot))^2 \leq 16. \quad (20)$$

Any optimal solution $\hat{x}^\varepsilon(\cdot)$ of (17)–(20) is the projection of an extremal $(\hat{x}^\varepsilon(\cdot), \hat{p}^\varepsilon(\cdot), p^{0\varepsilon}, \hat{u}^\varepsilon(\cdot))$ such that

$$\begin{aligned} \dot{\hat{p}}_1^\varepsilon(t) &= \hat{p}_2^\varepsilon(t) \\ \dot{\hat{p}}_2^\varepsilon(t) &= -\hat{p}_1^\varepsilon(t) - \hat{p}_2^\varepsilon(t)(1.4 - 0.42\hat{x}_2^\varepsilon(t)^2). \end{aligned}$$




Fig. 7. Trajectory.




Fig. 8. Adjoint vector.

Assumption 3 is verified, and the controls that satisfy the Pontryagin maximization condition (11) are given by

$$\begin{aligned} \hat{u}_1^\varepsilon(t) &= \frac{4\hat{p}_2^\varepsilon(t)}{\sqrt{(\hat{p}_2^\varepsilon(t))^2 + \varepsilon^2(\hat{p}_1^\varepsilon(t))^2}}, \\ \hat{u}_2^\varepsilon(t) &= \frac{4\varepsilon\hat{p}_1^\varepsilon(t)}{\sqrt{(\hat{p}_2^\varepsilon(t))^2 + \varepsilon^2(\hat{p}_1^\varepsilon(t))^2}}. \end{aligned} \quad (21)$$

Applying a shooting method to this problem, we determine the optimal trajectory of the regularized problem, and we indeed observe the expected convergence of $(\hat{x}^\varepsilon(\cdot), \hat{p}^\varepsilon(\cdot), -1, \hat{u}^\varepsilon)$ towards $(\hat{x}(\cdot), \hat{p}(\cdot), -1, \hat{u})$, as ε tends to 0, in agreement with Theorem 4 (see Figs. 7–9).

The optimal controls (21) are smooth functions of t , therefore the algorithms presented in [35] to compute the first conjugate time along a smooth extremal curve can be applied. Here we will apply the test for conjugate times explained in [35] when the final time is free and the extremal is normal. Let us briefly recall this test. The maximized Hamiltonian writes as

$$\begin{aligned} H_r^\varepsilon(\hat{x}^\varepsilon, \hat{p}^\varepsilon) &= \hat{p}_1^\varepsilon \left(\hat{x}_2^\varepsilon + \frac{4\varepsilon^2\hat{p}_1^\varepsilon}{\sqrt{(\hat{p}_2^\varepsilon)^2 + \varepsilon^2(\hat{p}_1^\varepsilon)^2}} \right) \\ &+ \hat{p}_2^\varepsilon \left(-\hat{x}_1^\varepsilon + \hat{x}_2^\varepsilon(1.4 - 0.14(\hat{x}_2^\varepsilon)^2) + \frac{4\hat{p}_2^\varepsilon}{\sqrt{(\hat{p}_2^\varepsilon)^2 + \varepsilon^2(\hat{p}_1^\varepsilon)^2}} \right) - 1. \end{aligned}$$




Fig. 9. Control.




Fig. 10. Determinant ($\varepsilon = 0.01$).

The aim is to compute the solution $Z^\varepsilon(\cdot) = (\delta x_1^\varepsilon(\cdot), \delta x_2^\varepsilon(\cdot), \delta p_1^\varepsilon(\cdot), \delta p_2^\varepsilon(\cdot))^T$ of the so-called variational system $\dot{Z}^\varepsilon(t) = V(t)Z^\varepsilon(t)$ along the extremal $(\hat{x}^\varepsilon(\cdot), \hat{p}^\varepsilon(\cdot))$, where

$$V(t) = \begin{pmatrix} \frac{\partial^2 H_r^\varepsilon}{\partial x \partial p}(\hat{x}^\varepsilon(t), \hat{p}^\varepsilon(t)) & \frac{\partial^2 H_r^\varepsilon}{\partial p^2}(\hat{x}^\varepsilon(t), \hat{p}^\varepsilon(t)) \\ -\frac{\partial^2 H_r^\varepsilon}{\partial x^2}(\hat{x}^\varepsilon(t), \hat{p}^\varepsilon(t)) & -\frac{\partial^2 H_r^\varepsilon}{\partial x \partial p}(\hat{x}^\varepsilon(t), \hat{p}^\varepsilon(t)) \end{pmatrix}$$

with initial conditions $(\delta x_1^\varepsilon(0), \delta x_2^\varepsilon(0)) = (0, 0)$ and $(\delta p_1^\varepsilon(0), \delta p_2^\varepsilon(0))$ such that the scalar product $\langle (f_1^\varepsilon(0), f_2^\varepsilon(0)), (\delta p_1^\varepsilon(0), \delta p_2^\varepsilon(0)) \rangle$ is equal to 0, where $(f_1^\varepsilon, f_2^\varepsilon)$ is the dynamics, given by

$$\begin{cases} f_1^\varepsilon(t) = x_2^\varepsilon(t) + \frac{4\varepsilon^2 p_1^\varepsilon(t)}{\sqrt{(p_2^\varepsilon(t))^2 + \varepsilon^2 (p_1^\varepsilon(t))^2}}, \\ f_2^\varepsilon(t) = -x_1^\varepsilon(t) + x_2^\varepsilon(t)(1.4 - 0.14x_2^\varepsilon(t)^2) \\ \quad + \frac{4p_2^\varepsilon(t)}{\sqrt{(p_2^\varepsilon(t))^2 + \varepsilon^2 (p_1^\varepsilon(t))^2}}. \end{cases}$$

The first geometric conjugate time is then the first positive zero of the function $t \mapsto \det(\delta x_1^\varepsilon(t), \delta x_2^\varepsilon(t), f_1^\varepsilon(t), f_2^\varepsilon(t))$ (see Fig. 10).

We report on Table 1 the values of the first geometric conjugate time of the optimal trajectory $\hat{x}^\varepsilon(\cdot)$, for different values of ε . We observe that, as expected, t_c^ε converges to $t_c \simeq 3.14750955$ as ε tends to 0.

Another possible test (see [35]) is to compute numerically solutions $Z_i(\cdot) = (\delta x_{1i}^\varepsilon(\cdot), \delta x_{2i}^\varepsilon(\cdot), \delta p_{1i}^\varepsilon(\cdot), \delta p_{2i}^\varepsilon(\cdot))$, $i = 1, 2$, of the variational system considered previously, with initial conditions

Table 1
Values of t_c^ε .

ε	t_c^ε
0.1	3.26735859
0.01	3.1559626
0.001	3.14844987
0.0001	3.14760515




Fig. 11. First singular value of $J^\varepsilon(t)$ ($\varepsilon = 0.01$).

$(\delta p_{11}^\varepsilon(0), \delta p_{21}^\varepsilon(0)) = (1, 0)$ and $(\delta p_{12}^\varepsilon(0), \delta p_{22}^\varepsilon(0)) = (0, 1)$, and then to compute the rank of the matrix

$$J^\varepsilon(t) = \begin{pmatrix} \delta x_{11}^\varepsilon(t) & \delta x_{21}^\varepsilon(t) \\ \delta x_{12}^\varepsilon(t) & \delta x_{22}^\varepsilon(t) \end{pmatrix}.$$

This rank must be equal to 1 outside a conjugate time, and 0 at a conjugate time. In order to compute it, we use a singular value decomposition of $J^\varepsilon(t)$; then, a conjugate time occurs whenever the first singular value of $J^\varepsilon(t)$ vanishes (see Fig. 11).

In this first example, the first conjugate time t_c of the optimal bang–bang trajectory $\hat{x}(\cdot)$ coincides with the second switching time. We next provide an example where the first conjugate time is equal to the third switching time.

3.2. Second example

Consider the minimal time control problem for the control system

$$\begin{aligned} \dot{x}_1(t) &= \sin(x_2(t)), \\ \dot{x}_2(t) &= -\sin(x_1(t)) + u_1(t), \end{aligned} \quad (22)$$

with the control constraint

$$|u_1(\cdot)| \leq 1, \quad (23)$$

and with the boundary conditions

$$x_1(0) = x_2(0) = 0, \quad x_1(t_f) = 2.9, \quad x_2(t_f) = 0.1. \quad (24)$$

From the Pontryagin maximum principle, any optimal solution $\hat{x}(\cdot)$ of (22)–(24) is the projection of an extremal $(\hat{x}(\cdot), \hat{p}(\cdot), p^0, \hat{u}_1(\cdot))$ such that

$$\begin{aligned} \dot{\hat{p}}_1(t) &= \hat{p}_2(t) \cos(\hat{x}_1(t)), \\ \dot{\hat{p}}_2(t) &= -\hat{p}_1(t) \cos(\hat{x}_2(t)), \end{aligned}$$

and the maximization condition $\hat{p}_2(t)\hat{u}_1(t) = \max_{|w| \leq 1} (\hat{p}_2(t)w)$ must hold almost everywhere on $[0, t_f]$. It is easy to see that $\hat{p}_2(\cdot)$ cannot vanish on some subinterval, and it follows that the optimal control $\hat{u}_1(\cdot)$ is bang–bang, equal to $\hat{u}_1(t) = \text{sign}(\hat{p}_2(t))$. Applying a shooting method to problems (22)–(24) (with $p^0 = -1$), we




Fig. 12. Optimal trajectory.




Fig. 13. Optimal control.

determine the initial adjoint vector $\hat{p}(0) = (-0.5, 1)$, and observe that the trajectory has one switching time $\hat{\tau}_1 \simeq 3.26174615$ on $[0, t_f]$, that is, $\hat{u}_1(\cdot)$ is given by

$$\hat{u}_1(t) = \begin{cases} +1 & \text{for } 0 \leq t \leq \hat{\tau}_1, \\ -1 & \text{for } \hat{\tau}_1 \leq t \leq t_f, \end{cases}$$

with a final time $t_f \simeq 4.07756604$ (see Figs. 12 and 13). Furthermore, $\hat{x}(\cdot)$ is the unique minimal time solution and has a unique extremal lift (up to a multiplicative scalar), which is moreover normal.

Prolongating the trajectory $\hat{x}(\cdot)$ to the interval $[0, 11]$, we observe a second switching time at $\hat{\tau}_2 \simeq 6.21787838$, and a third one at $\hat{\tau}_3 \simeq 10.46930198$. Considering as in the previous example the extremal field emanating from the vertical manifold, we observe on Figs. 14 and 15 that the extremal field crosses transversally the second switching surface, but reflects off the third switching surface, and it follows from Theorem 2 that the first conjugate time t_c is equal to $\hat{\tau}_3$.

We propose the following regularization. Consider the control system

$$\begin{aligned} \dot{x}_1^\varepsilon(t) &= \sin(x_2^\varepsilon(t)) + \varepsilon u_2^\varepsilon(t), \\ \dot{x}_2^\varepsilon(t) &= -\sin(x_1^\varepsilon(t)) + u_1^\varepsilon(t), \end{aligned} \quad (25)$$

with the control constraint

$$(u_1^\varepsilon(\cdot))^2 + (u_2^\varepsilon(\cdot))^2 \leq 1, \quad (26)$$




Fig. 14. Extremal field, $t \in [0, 11]$.




Fig. 15. Zoom on the overlap of the flow at the third switching time.

and the initial and final conditions (24). Any optimal solution $\hat{x}^\varepsilon(\cdot)$ of (24)–(26) is the projection of an extremal $(\hat{x}^\varepsilon(\cdot), \hat{p}^\varepsilon(\cdot), p^{0^\varepsilon}, \hat{u}^\varepsilon(\cdot))$ such that

$$\begin{aligned} \hat{p}_1^\varepsilon(t) &= \hat{p}_2^\varepsilon(t) \cos(\hat{x}_1^\varepsilon(t)), \\ \hat{p}_2^\varepsilon(t) &= -\hat{p}_1^\varepsilon(t) \cos(\hat{x}_2^\varepsilon(t)), \end{aligned}$$

and the maximization condition implies that the extremal controls are given by

$$\begin{aligned} \hat{u}_1^\varepsilon(t) &= \frac{\hat{p}_2^\varepsilon(t)}{\sqrt{(\hat{p}_2^\varepsilon(t))^2 + \varepsilon^2(\hat{p}_1^\varepsilon(t))^2}}, \\ \hat{u}_2^\varepsilon(t) &= \frac{\varepsilon \hat{p}_1^\varepsilon(t)}{\sqrt{(\hat{p}_2^\varepsilon(t))^2 + \varepsilon^2(\hat{p}_1^\varepsilon(t))^2}}. \end{aligned} \quad (27)$$

Applying a shooting method to this problem, we determine the optimal trajectory of the regularized problem, and we indeed observe the expected convergence of $(\hat{x}^\varepsilon(\cdot), \hat{p}^\varepsilon(\cdot), -1, \hat{u}^\varepsilon)$ towards $(\hat{x}(\cdot), \hat{p}(\cdot), -1, \hat{u}_1)$, as ε tends to 0, in agreement with Theorem 4 (see Figs. 16–18).

As in the previous example, the controls (27) are smooth functions of t , and we apply the algorithm described in [35], computing as before the determinant $\det(\delta x_1^\varepsilon(t) \delta x_2^\varepsilon(t), f_1^\varepsilon(t) f_2^\varepsilon(t))$ (see Fig. 19).




Fig. 16. Trajectory.




Fig. 17. Adjoint vector.




Fig. 18. Control.

We report on Table 2 the values of the first geometric conjugate time of the optimal trajectory $\hat{x}^\varepsilon(\cdot)$, for different values of ε . We observe that, as expected, t_c^ε converges to t_c as ε tends to 0.

Remark 3.1. We observe on both previous examples that it is not needed to consider very small values of ε to estimate the first




Fig. 19. Determinant ($\varepsilon = 0.1$).

Table 2

Values of t_c^ε .

ε	t_c^ε
0.1	10.01593283
0.01	10.3164905
0.001	10.41858121
0.0001	10.45291892
0.00001	10.46419119

conjugate time t_c . Indeed, a conjugate time of a locally bang–bang trajectory can only occur at a switching time (see Remark 1.2) and, under our assumptions, switching times are isolated (see Remark 1.1). From Theorem 6, the first geometric conjugate time t_c^ε converges to t_c , when ε tends to 0. Therefore, as soon as ε is small enough so that t_c^ε is in a (not necessarily so small) neighborhood of some switching time \hat{t}_s of the bang–bang trajectory $\hat{x}(\cdot)$, this means that the bang–bang conjugate time t_c is equal to that switching time \hat{t}_s .

3.3. Conclusion

In this article we focused on the problem of determining an efficient procedure to compute the first conjugate time t_c for the minimal time problem for single-input control-affine systems $\dot{x} = X(x) + u_1 Y_1(x)$ in \mathbb{R}^n with the control constraint $|u_1(\cdot)| \leq 1$. We used the asymptotic approach developed in [36] which consists in adding new smooth vector fields Y_2, \dots, Y_m and a small parameter $\varepsilon > 0$, so as to come up with the minimal time problem for the system $\dot{x} = X(x) + u_1^\varepsilon Y_1(x) + \varepsilon \sum_{i=2}^m u_i^\varepsilon Y_i(x)$, under the control constraint $\sum_{i=1}^m (u_i^\varepsilon(\cdot))^2 \leq 1$, with the same boundary conditions as the initial problem. Under appropriate assumptions, the optimal controls of the latter problem, depending on ε , are smooth functions of t , and the theoretical and practical results for the conjugate time theory that are well known in the smooth case can be applied to the regularized problem. We proved that the first conjugate time of regularized problem converges to the first conjugate time initial problem, when ε tends to 0. We thus get as a byproduct an efficient way to compute conjugate times in the bang–bang case. We provided examples to illustrate our result.

Note that our results still hold if the control-affine system is considered on a manifold. In this article we considered \mathbb{R}^n for the sake of simplicity.

An open question is to extend our results to multi-input control-affine systems $\dot{x} = X(x) + \sum_{i=1}^p u_i Y_i(x)$ in \mathbb{R}^n , where $u = (u_1, \dots, u_p) \in L^\infty([0, t_f], \Delta)$ and Δ is a polyhedron (see [26]), or a convex polyhedron (see [21]), or a convex compact polyhedron

(see [20]) of \mathbb{R}^p . For $p > 1$, it would be interesting to consider the case where multiple switching times may occur, that is, when at least two control functions switch at the same time. A more general open question concerns the generalization to general cost functions, and/or more general dynamics.

Acknowledgements

We warmly thank Laura Poggiolini and Gianna Stefani for their help and remarks.

C. Silva is supported by FCT through the Ph.D. fellowship SFRH/DB/27272/2006.

References

- [1] L. Cesari, *Optimization – Theory and Applications*. Problems with Ordinary Differential Equations, in: *Applications of Mathematics*, vol. 17, Springer-Verlag, New York, 1983.
- [2] L. Pontryagin, V. Boltyanskii, R. Gramkrelidze, E. Mischenko, *The Mathematical Theory of Optimal Processes*, Wiley Interscience, 1962.
- [3] A. Bressan, A high order test for optimality of bang–bang controls, *SIAM J. Control Optim.* 23 (1985) 38–48.
- [4] G.A. Bliss, *Lectures on the Calculus of Variations*, The University of Chicago Press, Chicago, 1968.
- [5] A.V. Sarychev, The index of second variation of a control system, *Mat. Sb.* 113 (1980) 464–486. English transl. in *Math. USSR Sbornik* 41 (1982) 383–401.
- [6] H.J. Sussmann, Envelopes, conjugate points and optimal bang–bang extremals, in: M. Fliess, M. Hazewinkel (Eds.), *Proc. 1985 Paris Conf. on Nonlinear Systems*, Reidel Publishers, Dordrecht, the Netherlands, 1987.
- [7] A.J. Krener, H. Schättler, The structure of small-time reachable sets in low dimensions, *SIAM J. Control Optim.* 27 (1) (1989) 120–147.
- [8] V.G. Boltyanskii, Sufficient conditions for optimality and the justification of the dynamic programming method, *SIAM J. Control* 4 (1966) 326–361.
- [9] J.C. Dunn, Second-order optimality conditions in sets of L^∞ functions with range in a polyhedron, *SIAM J. Control Optim.* 33 (1995) 1603–1635.
- [10] K. Malanowski, H. Maurer, Sensitivity analysis for parametric control problems with control-state constraints, *Comput. Optim. Appl.* 5 (1996) 253–283.
- [11] H. Maurer, First and second order sufficient optimality conditions in mathematical programming and optimal control, *Math. Program. Study* 14 (1981) 163–177.
- [12] H. Maurer, D. Augustin, Sensitivity analysis and real-time control of parametric optimal control problems using boundary value methods, in: M. Grötschel, et al. (Eds.), *On-line Optimization of Large Scale Systems*, Springer-Verlag, Berlin, 2001, pp. 17–55.
- [13] H. Maurer, H.J. Oberle, Second order sufficient conditions for optimal control problems with free final time: The Riccati approach, *SIAM J. Control Optim.* 41 (2002) 380–403.
- [14] H. Maurer, S. Pickenhain, Second-order sufficient conditions for optimal control problems with mixed control-state constraints, *J. Optim. Theory Appl.* 86 (1995) 649–667.
- [15] A.A. Milyutin, N.P. Osmolovskii, *Calculus of Variations and Optimal Control*, in: *Transl. Math. Monogr.*, vol. 180, AMS, Providence, RI, 1998.
- [16] V. Zeidan, The Riccati equation for optimal control problems with mixed state-control constraints: Necessity and sufficiency, *SIAM J. Control Optim.* 32 (1994) 1297–1321.
- [17] D. Augustin, H. Maurer, Computational sensitivity analysis for state constrained optimal control problems, *Ann. Oper. Res.* 101 (2001) 75–99.
- [18] K. Malanowski, H. Maurer, Sensitivity analysis for optimal control problems subject to higher order state constraints, *Optimization with data perturbations, II*, *Ann. Oper. Res.* 101 (2001) 43–73.
- [19] N.P. Osmolovskii, F. Lempio, Transformation of quadratic forms to perfect squares for broken extremal, *Set-Valued Anal.* 10 (2002) 209–232.
- [20] A.V. Sarychev, First- and second-order sufficient optimality conditions for bang–bang controls, *SIAM J. Control Optim.* 35 (1997) 315–440.
- [21] H. Maurer, N.P. Osmolovskii, Second order sufficient conditions for time-optimal bang–bang control problems, *SIAM J. Control Optim.* 42 (2004) 2239–2263.
- [22] H. Maurer, C. Büskens, J.-H.R. Kim, C.Y. Kaya, Optimization methods for the verification of second order sufficient conditions for bang–bang controls, *Optim. Control Appl. Meth.* 26 (2005) 129–156.
- [23] A. Agrachev, G. Stefani, P. Zezza, Strong optimality for a bang–bang trajectory, *SIAM J. Control Optim.* 41 (4) (2002) 991–1014.
- [24] N.P. Osmolovskii, H. Maurer, Equivalence of second order optimality conditions for bang–bang control problems. Part 1: main results, *Control Cybernet.* 34 (3) (2005) 927–949.
- [25] N.P. Osmolovskii, H. Maurer, Equivalence of second order optimality conditions for bang–bang control problems. Part 2: Proofs, variational derivatives and representations, *Control Cybernet.*, 36 555–584.
- [26] L. Poggiolini, G. Stefani, State-local optimality of a bang–bang trajectory: a Hamiltonian approach, *Syst. Control Lett.* 53 (2004) 269–279.
- [27] A. Agrachev, M. Sigalotti, On the local structure of optimal trajectories in \mathbb{R}^3 , *SIAM J. Control Optim.* 42 (2003) 513–531.
- [28] J. Noble, H. Schättler, Sufficient conditions for relative minima of broken extremals in optimal control theory, *J. Math. Anal. Appl.* 269 (2002) 98–128.
- [29] U. Ledzewicz, H. Schättler, Optimal bang–bang controls for a two-compartment model in cancer chemotherapy, *J. Optim. Theory Appl.* 114 (2002) 609–637.
- [30] H.J. Sussmann, B. Piccoli, Regular synthesis and sufficient conditions for optimality, *SIAM J. Control Optim.* 39 (2) (2000) 359–410.
- [31] A. Agrachev, R. Gamkrelidze, Symplectic methods for optimization and control, in: B. Jajubczyk, W. Respondek (Eds.), *Geometry of Feedback and Optimal Control*, Marcel Dekker, 1998, pp. 19–78.
- [32] A. Agrachev, G. Stefani, P. Zezza, A Hamiltonian approach to strong minima in optimal control, in: *Proceedings of Symposia in Pure Mathematics*, vol. 64, American Mathematical Society, 1998, pp. 11–22.
- [33] M.R. Hestenes, Application of the theory of quadratic forms in Hilbert spaces to the calculus of variations, *Pacific J. Math.* 1 (1951) 525–582.
- [34] H. Maurer, N.P. Osmolovskii, Second order optimality conditions for bang–bang control problems, *Control Cybernet.* 32 (3) (2003) 555–584.
- [35] B. Bonnard, J.-B. Caillaud, E. Trélat, Second order optimality conditions in the smooth case and applications in optimal control, *ESAIM Control Optim. Calc. Var.* 13 (2) (2007) 207–236.
- [36] C.J. Silva, E. Trélat, Smooth regularization of bang–bang optimal control problems, *IEEE - Transactions on Automatic Control* (2010) (in press).
- [37] J.-P. Aubin, H. Frankowska, *Set-Valued Analysis*, Birkhäuser, Boston, 1990.
- [38] F.H. Clarke, R. Vinter, The relationship between the maximum principle and dynamic programming, *SIAM J. Control Optim.* 25 (5) (1987) 1291–1311.
- [39] P. Cannarsa, C. Sinestrari, *Semiconcave Functions, Hamilton–Jacobi Equations, and Optimal Control*, in: *Progress in Nonlinear Differential Equations and their Applications*, vol. 58, Birkhäuser Boston, Inc, Boston, MA, 2004.
- [40] L. Rifford, E. Trélat, Morse–Sard type results in sub-Riemannian geometry, *Math. Ann.* 332 (1) (2005) 145–159.
- [41] L. Rifford, E. Trélat, On the stabilization problem for nonholonomic distributions, *J. Eur. Math. Soc.* 11 (2) (2009) 223–255.
- [42] G. Stefani, Regularity properties of the minimum-time map, in: *Nonlinear synthesis (Sopron, 1989)*, in: *Progr. Systems Control Theory*, vol. 9, Birkhäuser Boston, Boston, MA, 1991, pp. 270–282.
- [43] A. Agrachev, Y. Sachkov, *Control theory from the geometric viewpoint*, in: *Control Theory and Optimization, II*, in: *Encyclopaedia of Mathematical Sciences*, vol. 87, Springer-Verlag, Berlin, 2004.
- [44] M.D. Kiefer, H. Schättler, Parametrized families of extremals and singularities in solutions to the Hamilton–Jacobi–Bellman equation, *SIAM J. Control Optim.* 37 (1999) 1346–1371.
- [45] H. Maurer, D. Augustin, Second order sufficient conditions and sensitivity analysis for the controlled Rayleigh problem, in: J. Guddat, et al. (Eds.), *Parametric Optimization and Related Topics, IV*, Lang, Frankfurt, 1997, pp. 245–259.