

Stability of periodic stationary solutions of scalar conservation laws with space-periodic flux

Anne-Laure Dalibard

Abstract

This article investigates the long-time behaviour of parabolic scalar conservation laws of the type $\partial_t u + \operatorname{div}_y A(y, u) - \Delta_y u = 0$, where $y \in \mathbb{R}^N$ and the flux A is periodic in y . More specifically, we consider the case when the initial data is an L^1 disturbance of a stationary periodic solution. We show, under polynomial growth assumptions on the flux, that the difference between u and the stationary solution vanishes for large times in L^1 norm. The proof uses a self-similar change of variables which is well-suited for the analysis of the long time behaviour of parabolic equations. Then, convergence in self-similar variables follows from arguments from dynamical systems theory. One crucial point is to obtain compactness in L^1 on the family of rescaled solutions; this is achieved by deriving uniform bounds in weighted L^2 spaces.

Keywords. Long time asymptotics, parabolic scalar conservation law, asymptotic expansion, moment estimates, homogenization

1 Introduction

The goal of this article is to study the long time limit of solutions of the equation

$$\partial_t u + \operatorname{div}_y A(y, u) - \Delta_y u = 0, \quad t > 0, y \in \mathbb{R}^N, \quad (1.1)$$

where the flux $A : \mathbb{R}^N \times \mathbb{R} \rightarrow \mathbb{R}^N$ is assumed to be \mathbb{T}^N -periodic with respect to its first variable. Here and in the rest of the article, \mathbb{T}^N denotes the N -dimensional torus, i.e. $\mathbb{T}^N = (\mathbb{R}/\mathbb{Z})^N$.

Classical results on scalar conservation laws (see for instance [20, 15]) ensure that the semi-group associated with equation (1.1) is well-defined in $L^1(\mathbb{R}^N) + L^\infty(\mathbb{R}^N)$. The

CNRS, UMR 8553-DMA, Ecole Normale Supérieure, 45 rue d'Ulm, 75005 Paris, FRANCE. e-mail: Anne-Laure.Dalibard@ens.fr

Mathematics Subject Classification (2010): 35B35, 35B40, 35B27

case when the initial data belongs to $U(y) + L^1(\mathbb{R})$ (when $N = 1$), where U is a viscous shock profile of equation (1.1) has already been dealt with in a previous article, see [8]. In the present paper, we restrict our study to the case when the initial data belongs to $v(y) + L^1(\mathbb{R}^N)$, where v is a given periodic stationary solution of (1.1). The choice of such initial data stems from the fact that the semi-group associated with (1.1) is non-expansive in the L^1 norm; hence it is natural to investigate stability in $L^1(\mathbb{R}^N)$.

When the flux A is linear, say

$$A(y, u) = \alpha(y)u,$$

this study coincides (at least for some particular functions α) with the one led by Adrien Blanchet, Jean Dolbeault and Michal Kowalczyk in [5] on the large time behaviour of Brownian ratchets, as we will explain in Remark 1.3. It is proved in [5] that if the flux A is linear and if

$$\limsup_{t \rightarrow \infty} \frac{1}{(1 + 2t)^2} \int_{\mathbb{R}^N} |u(t, y) - v(y)| (y - ct)^4 dy < \infty \quad (1.2)$$

for some velocity $c \in \mathbb{R}^N$ which will be defined later on (see (1.7)), then there exists a constant C and a number $\kappa \in (0, 1/2)$ such that

$$\int_{\mathbb{R}^N} \left| u(t, y) - v(y) - \frac{M f_0(y)}{(1 + 2t)^{N/2}} F\left(\frac{y - ct}{\sqrt{1 + 2t}}\right) \right| dy \leq Ct^{-\kappa}, \quad (1.3)$$

where f_0 is the solution of an elliptic equation in \mathbb{T}^N (see (1.6)), F is a Gaussian profile, and M is the mass of the initial disturbance, i.e.

$$M = \int_{\mathbb{R}^N} (u|_{t=0} - v).$$

(1.3) is a technical assumption which is expected to hold for a large class of initial data (but such a fact is not proved in [5]).

Unfortunately, as we explained in [8], the above result does not imply that the same convergence holds in the nonlinear case. Moreover, the proof of [5], which is based on entropy dissipation methods together with Log-Sobolev Poincaré inequalities, can hardly be transposed as such to a nonlinear setting, although attempts in this direction have been made: for instance, in [10] M. Di Francesco and P. Markowich prove convergence towards diffusive waves for the Burgers equation thanks to entropy dissipation methods. However, their strategy relies crucially on the Hopf-Cole formula, and therefore could not be transposed in the present general setting. Hence we have chosen here a slightly different approach, which enables us to extend some of the results of [5] to a nonlinear context. Additionally, we recover a weaker version of the convergence (1.3), but without the need for assumption (1.2). In fact, we prove that (1.2) holds for a class of initial data which is dense in $v + L^1(\mathbb{R}^N)$.

The present work is also embedded in the broader study of the long time behaviour of conservation laws. We refer the interested reader to the review paper by D. Serre [21] (and the references therein) for a thorough description of the homogeneous case, in which the author investigates the stability of stationary solutions of scalar conservation laws in various models (parabolic and hyperbolic settings, relaxation models...)

Before stating the main results of this paper, let us now recall a few properties of equation (1.1). First, according to a result of [7], periodic stationary solutions of (1.1) exist, provided the flux A satisfies some growth assumptions. In fact, several different growth regimes were studied in [7]; we only recall one of them here, which is the most relevant with regards to our purposes. In the rest of the article, we assume that A belongs to $W_{\text{loc}}^{1,\infty}(\mathbb{T}^N \times \mathbb{R}^N)$, and that

$$\exists p_0 \in \mathbb{R}, \forall y \in \mathbb{T}^N, \operatorname{div}_y A(y, p_0) = 0. \quad (1.4)$$

We also assume that there exists $n \in (0, (N+2)/N)$ such that

$$\begin{aligned} \forall P > 0, \exists C_P > 0, \forall (p, q) \in \mathbb{R}^2, |p| \leq P, \\ \begin{cases} |\partial_p A(y, p+q) - \partial_p A(y, p)| \leq C_P(|q| + |q|^n), \\ |\operatorname{div}_y A(y, p+q) - \operatorname{div}_y A(y, p)| \leq C_P(|q| + |q|^n). \end{cases} \end{aligned} \quad (1.5)$$

These assumptions were introduced in [8, 7]. They ensure that for any $r \in \mathbb{R}$, there exists a unique periodic stationary solution of (1.1) with mean value r ; we refer to [7] for a discussion of the optimality of conditions (1.4), (1.5). Moreover, if u is a solution of (1.1) with initial data $u|_{t=0} \in v + L^1 \cap L^\infty(\mathbb{R}^N)$, where $v \in W^{1,\infty}(\mathbb{R}^N)$ is any stationary solution of (1.1), then $u \in L^\infty([0, \infty) \times \mathbb{R}^N)$. This result will be used several times in the article, and its proof is recalled in Appendix A.

We now introduce the profiles which characterize the asymptotic behaviour of the function u :

- First, the function f_0 occurring in (1.3) is the unique solution in $H^1(\mathbb{T}^N)$ of the equation

$$-\Delta_y f_0 + \operatorname{div}_y(\alpha_1 f_0) = 0, \quad \langle f_0 \rangle = 1, \quad (1.6)$$

where

$$\alpha_1(y) := (\partial_p A)(y, v(y)) \in L^\infty(\mathbb{T}^N)^N.$$

Above and in the rest of the article, the notation $\langle \cdot \rangle$ stands for the average on the torus \mathbb{T}^N , that is

$$\langle f \rangle := \int_{\mathbb{T}^N} f.$$

- The drift velocity c is then defined by

$$c = N \langle \alpha_1 f_0 \rangle. \quad (1.7)$$

- The last function which will appear in the asymptotic profile of u is the equivalent, in the non-linear case, of the Gaussian profile F occurring in (1.3); it is the unique solution, in a suitable functional space, of an elliptic equation of the form

$$-\sum_{1 \leq i, j \leq N} \eta_{i,j} \partial_i \partial_j F_M - \operatorname{div}_x(x F_M) + a \cdot \nabla_x F_M^2 = 0 \text{ in } \mathbb{R}^N, \text{ with } \int_{\mathbb{R}^N} F_M = M \in \mathbb{R}, \quad (1.8)$$

where the coefficients $\eta_{i,j}$ and a are constant, and the matrix $(\eta_{i,j})_{1 \leq i, j \leq N}$ is coercive. Unfortunately, giving the precise definition of $\eta_{i,j}$ and a would take us too far at this stage. Let us merely mention that $a = 0$ if $N \geq 2$. We also recall that thanks to a result of J. Aguirre, M. Escobedo, and E. Zuazua (see [1]), the above equation has a unique solution for all $M \in \mathbb{R}$, and we refer to the next section for more details. In particular, $\eta_{i,j}$, a are defined in (2.14), (2.16) respectively.

The main result of this paper is the following:

Theorem 1.1. *Let $A \in W_{loc}^{5,\infty}(\mathbb{T}^N \times \mathbb{R})^N$, and assume that A satisfies (1.4), (1.5).*

Let v be a periodic stationary solution of (1.1), and let $u_{ini} \in v + L^1(\mathbb{R}^N)$. Let u be the unique solution of (1.1) with initial condition $u|_{t=0} = u_{ini}$. Set

$$M := \int_{\mathbb{R}^N} (u_{ini} - v) dy.$$

Then as $t \rightarrow \infty$,

$$\int_{\mathbb{R}^N} \left| u(t, y) - v(y) - \frac{1}{(1+2t)^{N/2}} f_0(y) F_M \left(\frac{y-ct}{\sqrt{1+2t}} \right) \right| dy \rightarrow 0.$$

Remark 1.1. In fact, the regularity assumptions on the flux A are not as stringent as stated in the Theorem above. In particular, the conditions on the derivatives with respect to the space variable y can be considerably reduced. When looking closely at the proof, the correct regularity assumptions on A are

$$\begin{aligned} \partial_p^k A &\in L_{loc}^\infty(\mathbb{T}^N \times \mathbb{R})^N \quad \forall k \in \{0, 1, \dots, 4\}, \\ \operatorname{div}_y A, \operatorname{div}_y \partial_p^4 A &\in L_{loc}^\infty(\mathbb{T}^N \times \mathbb{R}). \end{aligned}$$

Remark 1.2. Notice that even in the linear case, Theorem 1.1 does not yield any convergence rate, in contrast with (1.3). This is due to the method of proof, which relies on abstract arguments from dynamical systems theory. The convergence rate obtained by Blanchet, Dolbeault and Kowalczyk in [5] is based on more “constructive” techniques, namely entropy dissipation methods and Log Sobolev inequalities. The counterpart of such techniques lies in the necessity for assumption (1.2). We emphasize that such an assumption is not necessary here.

Remark 1.3. Let us now make precise the link between brownian ratchets and equation (1.1) in the linear case. In [5], A. Blanchet, J. Dolbeault and M. Kowalczyk study the long time behaviour of the solution $f = f(t, y)$ of the equation

$$\partial_t f = \Delta_y f + \operatorname{div}_y(\nabla \psi(y - \omega t) f), \quad t > 0, y \in \mathbb{R}^N, \quad (1.9)$$

with $\psi \in \mathcal{C}^2(\mathbb{T}^N)$, $\omega \in \mathbb{R}^N$. Setting

$$u(t, y) = f(t, y + \omega t) \quad \forall t > 0, \forall y \in \mathbb{R}^N,$$

we see that u satisfies

$$\partial_t u + \operatorname{div}_y(\alpha(y)u) - \Delta_y u = 0,$$

where the drift coefficient α is given by

$$\alpha(y) = -\omega - \nabla_y \psi(y). \quad (1.10)$$

Hence the study of (1.9) and that of (1.1) in the linear case are closely related; they are strictly equivalent in dimension one, since any function $\alpha \in \mathcal{C}^1(\mathbb{T})$ can be decomposed as

$$\alpha = \int_{\mathbb{T}} \alpha + \left(\alpha - \int_{\mathbb{T}} \alpha \right) = \int_{\mathbb{T}} \alpha + \partial_y \phi, \quad \text{for some } \phi \in \mathcal{C}^2(\mathbb{T}).$$

The equivalence does not hold when $N \geq 2$, but in fact, all the results of [5] remain true for an arbitrary drift $\alpha \in \mathcal{C}^1(\mathbb{T}^N)$ (using exactly the same techniques as the ones developed in [5]). This will be a consequence of the analysis we will perform in the next sections. The choice for a function α with the structure (1.10) stems from physical considerations (see [4]): equation (1.9) describes the evolution of the density of particles in a traveling potential, moving with constant speed ω .

In the course of the proof of Theorem 1.1, we will also prove that condition (1.2) holds for a large class of initial data. The precise result is the following:

Proposition 1.1. *Assume that the flux A is linear, and that $u_{ini} \in v + L^1(\mathbb{R}^N)$ is such that*

$$\exists m > N + 8, \quad \int_{\mathbb{R}^N} |u_{ini}(y) - v(y)|^2 (1 + |y|^2)^{m/2} dy < \infty.$$

Let u be the unique solution of (1.1) with initial data u_{ini} . Then (1.2) is satisfied. As a consequence (see [5]), (1.3) holds.

Hence for linear fluxes and for a large range of initial data, a rate of convergence can be given. The derivation of convergence rates in the non-linear case goes beyond the scope of this article; in fact, the standard methods to derive convergence rates rely on the use of entropy-entropy dissipation inequalities (see [10] in the case of the Burgers equation), which we have chosen not to use here.

Another consequence of Theorem 1.1 is the stability of stationary shock profiles of equation (1.1) (see [8]) in dimension one:

Definition 1.1. Assume that $N = 1$. A *stationary shock profile* is a stationary solution U of (1.1) such that there exists periodic stationary solutions v_l, v_r of (1.1) such that

$$\begin{aligned}\lim_{y \rightarrow +\infty} [U(y) - v_r(y)] &= 0, \\ \lim_{y \rightarrow -\infty} [U(y) - v_l(y)] &= 0.\end{aligned}$$

It was proved in [8] that the stability of shock profiles is a consequence of the stability of periodic stationary solutions. Thus we have the following

Corollary 1.1. Assume that $N = 1$, and that the hypotheses of Theorem 1.1 are satisfied. Let $U \in L^\infty(\mathbb{R})$ be a stationary shock profile of (1.1). Let $u_{ini} \in U + L^1(\mathbb{R})$ such that

$$\int_{\mathbb{R}} (u_{ini} - U) = 0,$$

and let u be the unique solution of (1.1) with initial data u_{ini} . Then

$$\lim_{t \rightarrow \infty} \|u(t) - U\|_{L^1(\mathbb{R})} = 0.$$

The strategy of proof of Theorem 1.1 is close to the one developed in [12], in which M. Escobedo and E. Zuazua study the long time behaviour of a homogeneous version of (1.1); we also refer the interested reader to [11], in which M. Escobedo, J.L. Vazquez and E. Zuazua extend the analysis performed in [12] to the case when the flux has sub-critical growth, and to [23], in which E. Zuazua extends the results of [12] to more general situations. The first step of the analysis consists in a self-similar change of variables, which helps us to focus on the appropriate length scales; this will be done in the next section, in which we also derive the equations on the limit profiles f_0 and F_M . Then, in section 3, we obtain some compactness on the rescaled sequence by deriving some uniform L^2 bounds in weighted spaces. Eventually, we conclude the proof in Section 4 by using semi-group arguments inherited from dynamical systems theory.

Throughout the article, we will use the following notation: if $\psi \in L^\infty_{loc}(\mathbb{R}^N)$, we set, for all $p \in [1, \infty)$,

$$\begin{aligned}L^p(\psi) &:= \left\{ u \in L^p_{loc}(\mathbb{R}^N), \int_{\mathbb{R}^N} |u|^p \psi < +\infty \right\}, \\ \text{and } \|u\|_{L^p(\psi)} &= \left(\int_{\mathbb{R}^N} |u|^p \psi \right)^{1/p}, \\ H^1(\psi) &:= \{ u \in L^2(\psi), \nabla u \in L^2(\psi) \}, \\ \text{and } \|u\|_{H^1(\psi)}^2 &= \|u\|_{L^2(\psi)}^2 + \|\nabla u\|_{L^2(\psi)}^2.\end{aligned}$$

Sobolev spaces of the type $W^{s,p}(\psi)$, $H^s(\psi)$, with $s \in \mathbb{N}$ arbitrary and $p \in [1, \infty)$, are defined in a similar fashion. When we write $\|u\|_{L^p(\mathbb{R}^N)}$, without specifying a weight function, we always refer to the usual L^p norm in \mathbb{R}^N , with respect to the Lebesgue measure (i.e. $\psi \equiv 1$).

2 The homogenized system

The goal of this section is to analyze the expected asymptotic behaviour of the solution $u(t)$ of equation (1.1); to that end, we change the space and time variables and introduce a parabolic scaling, which is appropriate for the study of the long time behaviour of diffusion equations. Then, using a two-scale Ansatz in space and time which was introduced in [5], we construct an approximate solution of the rescaled system. Eventually, we recall and derive several properties of the limit system.

2.1 Parabolic scaling

Consider the solution $u \in L_{\text{loc}}^\infty([0, \infty) \times \mathbb{R}^N)$ of (1.1), with $u|_{t=0} = u_{\text{ini}} \in v + L^1 \cap L^\infty(\mathbb{R}^N)$. It is a classical feature of scalar conservation laws that the semi-group associated with (1.1) is contractant in $L^1(\mathbb{R}^N)$. Hence, for all $t \geq 0$, $u(t) \in v + L^1(\mathbb{R}^N)$, and

$$\|u(t) - v\|_1 \leq \|u_{\text{ini}} - v\|_1.$$

Thus it is natural to compute the equation satisfied by $f(t) = u(t) - v \in L^1(\mathbb{R}^N)$: since v is a stationary solution of (1.1), there holds

$$\partial_t f + \text{div}_y B(y, f) - \Delta_y f = 0, \quad t > 0, y \in \mathbb{R}^N,$$

where the flux B is defined by

$$B(y, f) = A(y, v(y) + f) - A(y, v(y)), \quad \forall (y, f) \in \mathbb{T}^N \times \mathbb{R}.$$

The flux $B(y, f)$ vanishes at $f = 0$, for all y . Moreover, if the flux A satisfies the assumptions of Theorem 1.1, there exists $\alpha_1 \in C^1(\mathbb{T}^N)$ and $\tilde{B}_1 \in \mathcal{C}(\mathbb{T}^N \times \mathbb{R})$ such that

$$B(y, f) = \alpha_1(y)f + \tilde{B}_1(y, f), \tag{2.1}$$

and the flux \tilde{B}_1 is such that

$$\forall X > 0, \exists C_X > 0, \forall f \in [-X, X], \forall y \in \mathbb{T}^N, \left| \tilde{B}_1(y, f) \right| \leq C_X |f|^2.$$

At some point in the proof, we will need a more refined approximation of B in a neighbourhood of $f = 0$; we thus also introduce $\alpha_2, \alpha_3 \in L^\infty(\mathbb{T}^N)$, $\tilde{B}_3 \in L^\infty(\mathbb{T}^N \times \mathbb{R})$ such that

$$B(y, f) = \alpha_1(y)f + \alpha_2(y)f^2 + \alpha_3(y)f^3 + \tilde{B}_3(y, f),$$

and the flux \tilde{B}_3 is such that for all $X > 0$, there exists a constant $C_X > 0$, such that for all $f \in [-X, X]$, for all $y \in \mathbb{T}^N$,

$$\begin{aligned} \left| \tilde{B}_3(y, f) \right| &\leq C_X |f|^4, \\ \left| \text{div}_y \tilde{B}_3(y, f) \right| &\leq C_X |f|^4, \\ \left| \partial_f \tilde{B}_3(y, f) \right| &\leq C_X |f|^3. \end{aligned}$$

The existence of α_i ($i = 1, 2, 3$) and the bounds on \tilde{B}_1, \tilde{B}_3 are ensured by the assumption that $A \in W^{5,\infty}(\mathbb{T}^N \times \mathbb{R})$. Notice in particular that

$$\alpha_1(y) = \partial_f B(y, f)|_{f=0} = (\partial_p A)(y, v(y)), \quad \forall y \in \mathbb{T}^N.$$

As explained in [5], the interplay between the diffusion and the drift α_1 induces a displacement of the center of mass. In the linear case, that is, when $\tilde{B}_1 = 0$, the evolution of the center of mass can be computed as follows: since the function f satisfies

$$\partial_t f + \operatorname{div}_y(\alpha_1 f) - \Delta_y f = 0,$$

there holds

$$\frac{d}{dt} \int_{\mathbb{R}^N} y f(t, y) dy = N \int_{\mathbb{R}^N} \alpha_1(y) f(t, y) dy.$$

Now, for $t \geq 0, y \in \mathbb{T}^N$, set

$$\tilde{f}(t, y) = \sum_{k \in \mathbb{Z}^N} f(t, y + k).$$

Since the function α_1 is periodic, \tilde{f} satisfies

$$\partial_t \tilde{f} + \operatorname{div}_y(\alpha_1 \tilde{f}) - \Delta_y \tilde{f} = 0, \quad t > 0, y \in \mathbb{T}^N,$$

and we have, for all $t \geq 0$,

$$\int_{\mathbb{R}^N} \alpha_1 f(t) = \int_{\mathbb{T}^N} \alpha_1 \tilde{f}(t).$$

Using Lemma 1.1 of [18] together with a Poincaré inequality on the torus \mathbb{T}^N , it can be easily proved that as $t \rightarrow \infty$, $\tilde{f}(t)$ converges with exponential speed in $L^1(\mathbb{T}^N)$ towards $\langle \tilde{f} \rangle f_0$, where f_0 is the unique solution of (1.6). Additionally, notice that

$$\langle \tilde{f} \rangle = \sum_{k \in \mathbb{Z}^N} \langle f(\cdot + k) \rangle = \int_{\mathbb{R}^N} f = M.$$

Consequently, setting

$$c := N \langle \alpha_1 f_0 \rangle$$

we infer that in the linear case,

$$\frac{d}{dt} \int_{\mathbb{R}} (y - ct) f \rightarrow 0 \quad \text{exponentially fast.}$$

In fact, it turns out that the nonlinearity has no effect on this displacement, although this is not quite clear if we try to include the quadratic term \tilde{B}_1 in the above calculation.

We will justify this result by formal calculations in the next paragraph. Nonetheless, it can be proved in the case $N = 1$ (see for instance [8]) that when $\|f_0\|_1$ is not too large,

$$\|f(t)\|_{L^2(\mathbb{R})} \leq C \frac{\|f_0\|_1}{t^{1/4}} \quad \forall t > 0,$$

and more generally, the L^p norm of $f(t)$ vanishes for long times for all $p \in (1, \infty]$. This somehow explains why the quadratic term does not modify the motion of the center of mass for large times: the term $\tilde{B}_1(\cdot, f(t, \cdot))$ vanishes in $L^1(\mathbb{R})$ as $t \rightarrow \infty$. Hence, hereinafter, we choose to make in the general case the same change of variables as the one dictated by the linear case. Precisely, let $U \in L_{\text{loc}}^\infty([0, \infty) \times \mathbb{R}^N)$ such that

$$f(t, y) = \frac{1}{(1+2t)^{N/2}} U \left(\log \sqrt{1+2t}, \frac{y-ct}{\sqrt{1+2t}} \right), \quad t \geq 0, y \in \mathbb{R}^N. \quad (2.2)$$

This change of variables is classical in the study of long-time parabolic dynamics, see for instance [12]. In the present case, our change of variables is exactly the same as in [5]; straightforward calculations lead to

$$\partial_\tau U - \operatorname{div}_x(xU) + R \operatorname{div}_x((\alpha_1(z) - c)U) - \Delta_x U = -R^{N+1} \operatorname{div}_x \tilde{B}_1 \left(z, \frac{U}{R^N} \right), \quad (2.3)$$

with $\tau > 0$, $x \in \mathbb{R}^N$, and where

$$R = e^\tau \quad \text{and} \quad z = Rx + c \frac{R^2 - 1}{2}.$$

Studying the long time behaviour of f amounts to studying the long time behaviour of U . Now, as $\tau \rightarrow \infty$, the quantity R becomes very large, and thus the variable z is highly oscillating. Hence, as emphasized in [5], the asymptotic study of equation (2.3) somehow falls into the scope of homogenization theory; the small parameter measuring the period of the oscillations is then $\varepsilon = R^{-1} = e^{-\tau}$. However, one substantial difference with classical homogenization problems is that the small parameter depends on time, which sometimes makes the proofs much more technical. We refer to [5] for more details.

Let us also mention that the homogenization of equation (2.3) with a ‘‘fixed’’ small parameter, and when the quadratic flux \tilde{B}_1 vanishes, has been performed by Thierry Goudon and Frédéric Poupaud in [14]. As a consequence, the formal asymptotic expansions which will be performed in the next section are in fact very close to the ones of [14].

2.2 Formal derivation of the limit system

As usual in homogenization problems (see [3] for instance), the idea is now to assume that the solution U of (2.3) admits an asymptotic development in powers of the small parameter measuring the period of the oscillations; in the present case, the small parameter

is $e^{-\tau}$, so that we expect the approximation to be valid for large times only. Hence, assume that when $\tau \gg 1$,

$$U(\tau, x) \approx U_0(\tau, x, z) + e^{-\tau}U_1(\tau, x, z) + e^{-2\tau}U_2(\tau, x, z) + \dots \quad (2.4)$$

where $z = e^\tau x + c(e^{2\tau} - 1)/2$ stands for the fast variable and where for all $(\tau, x) \in \mathbb{R}_+ \times \mathbb{R}^N$, the function

$$z \mapsto U_i(\tau, x, z)$$

is \mathbb{T}^N -periodic. Plugging the Ansatz (2.4) into equation (2.3) and identifying the powers of $R = e^\tau$ leads to a cascade of equations on the terms U_0, U_1 , etc. Notice that according to Lemma A.1 in the Appendix, $f \in L^\infty([0, \infty) \times \mathbb{R}^N)$, and thus U/R^N is bounded in L^∞ .

• **Terms of order R^2 :** Identifying the highest order terms in equation (2.3) when U is given by (2.4) leads to

$$c \cdot \nabla_z U_0 + \operatorname{div}_z((\alpha_1 - c)U_0) - \Delta_z U_0 = -\Delta_z U_0 + \operatorname{div}_z(\alpha_1 U_0) = 0, \quad z \in \mathbb{T}^N.$$

We recall the following result, which is a straightforward consequence of the Krein-Rutman Theorem (see [9]):

Lemma 2.1. *Let $\alpha \in L^\infty(\mathbb{T}^N)^N$. Consider the vector space*

$$E[\alpha] := \{w \in H^1(\mathbb{T}^N), -\Delta_z w + \operatorname{div}_z(\alpha w) = 0\}.$$

Then $\dim E[\alpha] = 1$, and there exists a unique function $m \in E[\alpha]$ such that $\langle m \rangle = 1$.

Moreover, $m \in W^{1,p}(\mathbb{T}^N)$ for all $p < \infty$, and

$$\inf_{z \in \mathbb{T}^N} m > 0.$$

In the present case, $E[\alpha_1] = \mathbb{R}f_0$, where f_0 is defined by (1.6). Hence there exists a function $F = F(\tau, x)$ such that

$$U_0(\tau, x, z) = f_0(z)F(\tau, x) \quad \forall (\tau, x, z) \in [0, \infty) \times \mathbb{R}^N \times \mathbb{T}^N. \quad (2.5)$$

• **Terms of order R^1 :** Concerning the terms of order $R^1 = e^\tau$, the case when the space dimension is equal to one has to be treated separately. Indeed,

$$\begin{aligned} R^{N+1} \operatorname{div}_x \tilde{B}_1 \left(z, \frac{U}{R^N} \right) &= R^{1-N} \operatorname{div}_x(\alpha_2 U^2) + R^{1-2N} \operatorname{div}_x(\alpha_3 U^3) \\ &\quad + R^{N+1} \operatorname{div}_x \left(\tilde{B}_3 \left(z, \frac{U}{R^N} \right) \right), \end{aligned}$$

and using the bounds on \tilde{B}_3 ,

$$\begin{aligned} R^{N+1} \operatorname{div}_x \left(\tilde{B}_3 \left(z, \frac{U}{R^N} \right) \right) &= R^N (\operatorname{div}_z \tilde{B}_3) \left(z, \frac{U}{R^N} \right) + R \nabla_x U \cdot (\partial_U \tilde{B}_3) \left(z, \frac{U}{R^N} \right) \\ &= \mathcal{O}(R^{-3N}) + \mathcal{O}(R^{-3N+2}) = \mathcal{O}(R^{-1}). \end{aligned}$$

We infer that if U is given by (2.4), then

$$\begin{aligned} R^{N+1} \operatorname{div}_x \tilde{B}_1 \left(z, \frac{U}{R^N} \right) &= R^{2-N} \operatorname{div}_z (\alpha_2 U_0^2) \\ &+ R^{1-N} [\operatorname{div}_x (\alpha_2 U_0^2) + 2 \operatorname{div}_z (\alpha_2 U_0 U_1)] \\ &+ R^{2-2N} \operatorname{div}_z (\alpha_3 U_0^3) \\ &+ \mathcal{O}(R^{-1}). \end{aligned} \quad (2.6)$$

Consequently, we obtain that when $N \geq 2$, the term U_1 solves the equation

$$-\Delta_z U_1 + \operatorname{div}_z (\alpha_1 U_1) = -\operatorname{div}_x ((\alpha_1 - c) U_0) + 2 \sum_{i=1}^N \frac{\partial^2 U_0}{\partial x_i \partial z_i}. \quad (2.7)$$

Since $U_0(t, x, z) = f_0(z)F(t, x)$, we have

$$\langle (\alpha_1 - c) U_0(t, x, \cdot) \rangle = F(t, x) (\langle \alpha_1 f_0 \rangle - c) = 0$$

by definition of c . Hence the right-hand side of (2.7) has zero mean value, and the compatibility condition is satisfied. Thus for all $(t, x) \in [0, \infty) \times \mathbb{R}^N$, (2.7) has a unique solution in $H^1(\mathbb{T}^N)$. Moreover, using the linearity of (2.7) together with the expression (2.5), we infer that U_1 can be written as

$$U_1(t, x, z) = f_1(z) \cdot \nabla_x F(t, x), \quad (2.8)$$

where $f_1 \in H^1(\mathbb{T}^N)^N$ satisfies

$$-\Delta_z f_{1,i} + \operatorname{div}_z (\alpha_1 f_{1,i}) = -f_0 (\alpha_{1,i} - c_i) + 2 \partial_{z_i} f_0, \quad \forall i \in \{1, \dots, N\}. \quad (2.9)$$

Notice that according to the regularity assumptions on the flux A , the function $\operatorname{div}_y \alpha_1$ belongs to $L^\infty(\mathbb{T}^N)$. Using boot-strap arguments for equation (1.6), we deduce that $f_0 \in W^{2,p}(\mathbb{T}^N)$ for all $p < \infty$, and therefore $f_1 \in W^{2,p}(\mathbb{T}^N)$ for all $p < \infty$. In particular, $f_1, f_0 \in W^{1,\infty}(\mathbb{T}^N)$.

If $N = 1$, on the other hand, the corrector U_1 solves the equation

$$-\partial_{zz} U_1 + \partial_z (\alpha_1 U_1) = -\partial_x ((\alpha_1 - c) U_0) + 2 \frac{\partial^2 U_0}{\partial x \partial z} - \partial_z (\alpha_2 U_0^2). \quad (2.10)$$

Notice that the compatibility condition is satisfied, for the same reason as before. Hence in this case,

$$U_1(t, x, z) = f_1(z) \partial_x F(t, x) + g_1(z) F(t, x)^2, \quad (2.11)$$

where $g_1 \in H^1(\mathbb{T})$ solves

$$-\Delta_z g_1 + \partial_z(\alpha_1 g_1) = -\partial_z(\alpha_2 (f_0)^2).$$

The fact that the compatibility condition is satisfied in all cases justifies *a posteriori* the use of the change of variables (2.2) in the nonlinear case. This means that, at least on a formal level, the displacement of the center of mass of the function f is unaffected by the presence of the quadratic term \tilde{B}_1 .

• **Terms of order R^0 :** As we identify the terms of order one in equation (2.3), we obtain

$$\begin{aligned} & -\Delta_z U_2 + \operatorname{div}(\alpha_1 U_2) \\ = & -\partial_\tau U_0 + \operatorname{div}_x(xU_0) + \Delta_x U_0 - \operatorname{div}_x((\alpha_1 - c)U_1) + 2 \sum_{i=1}^N \frac{\partial^2 U_1}{\partial x_i \partial z_i} + \mathcal{A}_{NL}, \end{aligned} \quad (2.12)$$

where the term \mathcal{A}_{NL} stems from the expansion of the nonlinear term \tilde{B}_1 . According to (2.6), we have

$$\begin{aligned} \mathcal{A}_{NL} &= \partial_x(\alpha_2 U_0^2) + 2\partial_z(\alpha_2 U_0 U_1) + \partial_z(\alpha_3 U_0^3) \quad \text{if } N = 1, \\ \mathcal{A}_{NL} &= \operatorname{div}_z(\alpha_2 U_0^2) \quad \text{if } N = 2, \\ \mathcal{A}_{NL} &= 0 \quad \text{if } N \geq 3. \end{aligned}$$

The evolution equation for the function F follows from the compatibility condition; precisely, we obtain

$$\partial_\tau F - \operatorname{div}_x(xF) - \Delta_x F + \operatorname{div}_x \langle (\alpha_1 - c)U_1 \rangle - \langle \mathcal{A}_{NL} \rangle = 0.$$

We now distinguish between the cases $N \geq 2$ and $N = 1$.

▷ If $N \geq 2$, $\langle \mathcal{A}_{NL} \rangle = 0$; using (2.8), we infer that F satisfies

$$\partial_\tau F - \operatorname{div}_x(xF) - \sum_{1 \leq i, j \leq N} \eta_{i,j} \frac{\partial^2 F}{\partial x_i \partial x_j} = 0, \quad \tau > 0, \quad x \in \mathbb{R}^N \text{ with } N \geq 2, \quad (2.13)$$

where the coefficients $(\eta_{i,j})_{1 \leq i, j \leq N}$ are given by

$$\eta_{i,j} = \delta_{i,j} - \langle (\alpha_{1,i} - c_i) f_{1,j} \rangle. \quad (2.14)$$

The following Lemma entails that equation (2.13) is well-posed:

Lemma 2.2. *The matrix $\eta := (\eta_{i,j})_{1 \leq i, j \leq N}$ is coercive.*

Lemma 2.2 is proved in [14] for any $N \geq 1$, and its proof is recalled in [5] when $N = 1$. For the reader's convenience, we sketch the main steps of the proof here, and we refer to [14], Proposition 4.6 for details.

Proof. Let L be the differential operator

$$L\phi = -\Delta_z\phi + \operatorname{div}_z(\alpha_1\phi).$$

The idea is to introduce, for all $j \in \{1, \dots, N\}$, the function χ_j which solves the adjoint problem

$$L^*\chi_j = -\Delta_z\chi_j - \alpha_1 \cdot \nabla_z\chi_j = \alpha_{1,j} - c_j, \quad \langle \chi_j \rangle = 0.$$

Since the right-hand side satisfies $\langle (\alpha_{1,j} - c_j)\psi \rangle = 0$ for all $\psi \in \ker L = E[\alpha_1] = \mathbb{R}f_0$, the function χ_j is well-defined. For all $\xi \in \mathbb{R}^N$, we have

$$\begin{aligned} \sum_{i,j} \langle (\alpha_{1,i} - c_i)f_{1,j} \rangle \xi_i \xi_j &= \langle L^*(\chi \cdot \xi)f_1 \cdot \xi \rangle \\ &= \langle \chi \cdot \xi (-f_0(\alpha_1 - c) \cdot \xi + 2\nabla_z f_0 \cdot \xi) \rangle \\ &= -\langle f_0 \chi \cdot \xi L^*(\chi \cdot \xi) \rangle + \langle 2\chi \cdot \xi \nabla_z f_0 \cdot \xi \rangle \\ &= -\langle L(f_0 \chi \cdot \xi) \chi \cdot \xi \rangle - 2 \langle f_0 \xi \cdot \nabla_z(\chi \cdot \xi) \rangle. \end{aligned}$$

Expanding $L(f_0 \chi \cdot \xi)$ and using the identity $Lf_0 = 0$ leads to

$$\langle L(f_0 \chi \cdot \xi) \chi \cdot \xi \rangle = \langle f_0 |\nabla_z(\chi \cdot \xi)|^2 \rangle.$$

Hence

$$\begin{aligned} \sum_{1 \leq i,j \leq N} \eta_{i,j} \xi_i \xi_j &= |\xi|^2 + \langle f_0 |\nabla_z(\chi \cdot \xi)|^2 \rangle + 2 \langle f_0 \xi \cdot \nabla_z(\chi \cdot \xi) \rangle \\ &= \langle f_0 |\xi + \nabla_z(\chi \cdot \xi)|^2 \rangle. \end{aligned}$$

We deduce that

$$\sum_{1 \leq i,j \leq N} \eta_{i,j} \xi_i \xi_j \geq 0 \quad \forall \xi \in \mathbb{R}^N.$$

Now, let $\xi \in \mathbb{R}^N$ such that $\sum \eta_{i,j} \xi_i \xi_j = 0$. Since $f_0(z) > 0$ for all z , we infer that

$$\xi + \nabla_z(\chi \cdot \xi) = 0 \quad \forall z \in \mathbb{T}^N.$$

Taking the average of the above inequality on \mathbb{T}^N leads to $\xi = 0$. Hence the matrix $(\eta_{i,j})$ is coercive. \square

▷ If $N = 1$, we have

$$\langle \mathcal{A}_{NL} \rangle = \partial_x \langle \alpha_2 U_0^2 \rangle = \langle \alpha_2 f_0^2 \rangle \partial_x F^2.$$

Moreover, in this case U_1 is given by (2.11); hence

$$\langle \operatorname{div}_x((\alpha_1 - c)U_1) \rangle = \langle (\alpha_1 - c)f_1 \rangle \partial_{xx} F + \langle (\alpha_1 - c)g_1 \rangle \partial_x F^2.$$

Consequently, the compatibility condition reads

$$\partial_\tau F - \partial_x(xF) + a\partial_x F^2 - \eta\partial_{xx}F = 0, \quad \tau > 0, \quad x \in \mathbb{R}, \quad (2.15)$$

where the coefficients a, η are given by

$$\begin{aligned} a &:= \langle \alpha_2 f_0^2 \rangle + \langle (\alpha_1 - c)g_1 \rangle, \\ \eta &:= 1 - \langle (\alpha_1 - c)f_1 \rangle. \end{aligned} \quad (2.16)$$

Lemma 2.2 states that the diffusion coefficient η is positive.

Remark 2.1. (i) The fact that the coefficient a appearing in (1.8) is zero for all $N \geq 2$ is consistent with the results of M. Escobedo and E. Zuazua in [12]: indeed, the authors of [12] consider the convection-diffusion equation

$$v_t - \Delta v = \alpha \cdot \nabla(|v|^{q-1}v) \quad \text{in } (0, \infty) \times \mathbb{R}^N$$

for some exponent $q \geq 1 + 1/N$ and for $\alpha \in \mathbb{R}^N$. They prove that if $q > 1 + 1/N$, the asymptotic behaviour of v is given by the heat kernel, i.e. the convective term does not play any role. On the contrary, if $q = 1 + 1/N$, v behaves asymptotically like the self-similar solution of a nonlinear equation.

In the present case, because of the Taylor expansion (2.1), the flux B consists of a linear drift term and a quadratic term. Hence the study performed in this paper is related to the case $q = 2$ of [12]; we infer that $q = 1 + 1/N$ if and only if $N = 1$, and we check that this is the only case when a nonlinear behaviour can be observed at the limit.

- (ii) In order to have a better understanding of the large time behaviour of equation (1.1), it would be interesting to compute the next term of the development, in the spirit of the paper by E. Zuazua [22]. In particular, the nonlinear behaviour of the equation is expected to appear at the next order for all $N \in \mathbb{N}$. Notice that this implies that the term U_1 is not, in fact, the next term in the development. However, such an analysis is beyond the scope of the present article.

The formal derivation of an approximate solution is now complete. In the following paragraphs, we recall or prove several results concerning the well-posedness and the long time behaviour of equations (2.13) and (2.15). We will often refer to the equation on F as the ‘‘homogenized equation’’; this term refers to equation (2.13) when $N \geq 2$, and (2.15) when $N = 1$.

2.3 Existence and uniqueness of stationary solutions of the homogenized equation

This paragraph is concerned with the existence and uniqueness (in suitable functional spaces) of stationary solutions of the homogenized equations (2.13) and (2.15). In the case when $N = 1$, or when $(\eta_{i,j})_{1 \leq i,j \leq N} = \lambda I$ for some $\lambda > 0$, such results are stated in [1]. In the general case, we merely use a linear change of variables, and the problem is then reduced to the case of an isotropic diffusion.

Lemma 2.3. *Assume that $N \geq 2$. For $\gamma > 0$, set $\psi_\gamma : x \in \mathbb{R}^N \mapsto \exp(\gamma|x|^2)$. Then there exists $\gamma > 0$ such that for all $M \in \mathbb{R}$, there exists a unique function $F_M \in H^1(\psi_\gamma)$ satisfying*

$$-\sum_{1 \leq i,j \leq N} \eta_{i,j} \partial_i \partial_j F_M - \operatorname{div}_x(x F_M) = 0, \quad \int_{\mathbb{R}^N} F_M = M. \quad (2.17)$$

Furthermore, the following properties hold:

- (i) For all $M \in \mathbb{R}$, $F_M = M F_1$;
- (ii) $F_1 \in W^{2,p} \cap C^\infty(\mathbb{R}^N)$ for all $p \in [1, \infty)$, and $F_1 \in H^2(\psi_\gamma)$;
- (iii) $F_1(x) > 0$ for all $x \in \mathbb{R}^N$.

Proof. The idea is to perform an affine change of variables in order to transform the diffusion term into a laplacian. Precisely, set

$$s_{i,j} = \frac{\eta_{i,j} + \eta_{j,i}}{2}, \quad 1 \leq i, j \leq N.$$

Then the matrix $S = (s_{i,j})$ is symmetric and positive definite (see Lemma 2.2); hence there exists an orthogonal matrix $O \in \mathcal{M}_n(\mathbb{R})$ and positive numbers λ_j such that

$$S = O^T \operatorname{Diag}(\lambda_1, \dots, \lambda_N) O.$$

Let us change the variables by setting

$$x = Py, \quad \text{with } P := O^T \operatorname{Diag}(\lambda_1^{1/2}, \dots, \lambda_N^{1/2}), \quad (2.18)$$

and for any function $F \in L^1(\mathbb{R}^N)$, define

$$\tilde{F}(y) = F(Py).$$

It can be readily checked that for all $x \in \mathbb{R}^N$,

$$\sum_{i,j} \eta_{i,j} \frac{\partial^2 F}{\partial x_i \partial x_j}(x) = \sum_{k,l} \tilde{\eta}_{k,l} \frac{\partial^2 \tilde{F}}{\partial y_k \partial y_l}(P^{-1}x),$$

where the coefficients $\tilde{\eta}_{k,l}$ are given by

$$\tilde{\eta}_{k,l} = \sum_{i,j} (P^{-1})_{k,i} (P^{-1})_{l,j} s_{i,j} = (P^{-1} S (P^{-1})^T)_{k,l}.$$

Using the definitions of the matrices P and S , we infer that

$$\tilde{\eta} = P^{-1} S (P^{-1})^T = I_N.$$

Thus the diffusion term is transformed into a laplacian with this change of variables.

Let us now compute the drift term. We have

$$x_i = (Py)_i,$$

and, denoting by (e_1, \dots, e_N) the canonical basis of \mathbb{R}^N ,

$$\begin{aligned} \frac{\partial F(x)}{\partial x_i} &= \frac{\partial}{\partial x_i} \tilde{F}(P^{-1}x) \\ &= (P^{-1}e_i) \cdot \nabla_y \tilde{F}(P^{-1}x). \end{aligned}$$

Thus, always setting $x = Py$,

$$\begin{aligned} x \cdot \nabla_x F(x) &= \sum_{i=1}^N (Py)_i (P^{-1}e_i) \cdot \nabla_y \tilde{F}(y) \\ &= \left[P^{-1} \left(\sum_{i=1}^N (Py)_i e_i \right) \right] \cdot \nabla_y \tilde{F}(y) \\ &= (P^{-1}Py) \cdot \nabla_y \tilde{F}(y) = y \cdot \nabla_y \tilde{F}(y). \end{aligned}$$

Notice that this property is in fact independent of the definition of the matrix P . Consequently, F_M is a solution of (2.17) if and only if \tilde{F}_M satisfies

$$-\Delta_y \tilde{F}_M - \operatorname{div}_y (y \tilde{F}_M) = 0, \quad \int_{\mathbb{R}^N} \tilde{F}_M = (\det S)^{-1/2} M.$$

The only solutions of the above equation in $H^1(\mathbb{R}^N)$ are Gaussian functions. Hence there exists a unique solution of (2.17) in $H^1(\mathbb{R}^N)$ for all M , and this solution is given by

$$F_M(x) = CM \exp\left(-\frac{1}{2}|P^{-1}x|^2\right),$$

where the positive constant C is a normalization factor. Moreover,

$$|P^{-1}x|^2 = \left| \operatorname{Diag}\left(\lambda_1^{-1/2}, \dots, \lambda_N^{-1/2}\right) Ox \right|^2,$$

and thus, since $|Ox|^2 = |x|^2$,

$$\left(\max_{1 \leq i \leq N} \lambda_i \right)^{-1} |x|^2 \leq |P^{-1}x|^2 \leq \left(\min_{1 \leq i \leq N} \lambda_i \right)^{-1} |x|^2.$$

All the properties of the lemma follow, with

$$\gamma < \left(2 \max_{1 \leq i \leq N} \lambda_i \right)^{-1}.$$

□

In the case when $N = 1$, the existence of a stationary solution is treated in [1]. Hence we merely recall the main results of [1] in that regard.

Lemma 2.4 (Aguirre, Escobedo, Zuazua). *Let $M \in \mathbb{R}$ be arbitrary, and let $a \in \mathbb{R}$, $\eta > 0$. Let $\gamma := (2\eta)^{-1}$.*

Then there exists a unique function $F_M \in H^1(\psi_\gamma)$ which satisfies

$$-\eta \partial_{xx} F_M - \partial_x(x F_M) + a \partial_x F_M^2 = 0, \quad \int_{\mathbb{R}} F_M = M.$$

Moreover, F_M enjoys the following properties:

- (i) $F_M \in W^{2,p} \cap C^\infty(\mathbb{R})$ for all $p \in [1, \infty)$, and $F_M \in H^2(\psi_\gamma)$;
- (ii) If $M > 0$, then $F_M(x) > 0$ for all $x \in \mathbb{R}^N$.

We deduce from the above Lemma that if $\gamma' < \gamma$, then there exists a constant $C_{\gamma'}$ such that

$$|F_M(x)|, |\partial_x F_M(x)| \leq C_{\gamma'} \exp(-\gamma' x^2) \quad \forall x \in \mathbb{R}.$$

Indeed, since $F_M \in H^2(\psi_\gamma)$, it can be easily proved that $F_M \psi_{\gamma'} \in H^2(\mathbb{R})$ for all $\gamma' < \gamma$. Sobolev embeddings then imply that $F_M \psi_{\gamma'} \in W^{1,\infty}(\mathbb{R})$.

The existence of stationary solutions of (2.13) and (2.15) is thus ensured. We now tackle the study of the properties of equations (2.13) and (2.15), focusing in particular on the long-time behaviour and on regularity issues.

2.4 Long time behaviour of the homogenized equation

We begin with a definition of the weight function $K \in C^\infty(\mathbb{R}^N)$, which plays a central role in the theory of existence. For $N \geq 2$, we use the change of variables (2.18), which was introduced in the proof of Lemma 2.3. This allows us to transform the matrix $(\eta_{i,j})$ into the identity matrix. For $t > 0$, $y \in \mathbb{R}^N$, set $\tilde{F}(t, y) = F(t, Py)$. If F is a solution of (2.13), then \tilde{F} solves

$$\partial_t \tilde{F} - \operatorname{div}_y(y \tilde{F}) - \Delta_y \tilde{F} = 0.$$

Consequently, the results of [12] can be directly applied to \tilde{F} , for which existence is proved in the functional space $L^2(K_0)$, where $K_0(y) = \exp(y^2/2)$. Performing the inverse change of variables, it is clear that the relevant weight function is given by

$$K(x) := K_0(P^{-1}x) = \exp\left(\frac{|P^{-1}x|^2}{2}\right). \quad (2.19)$$

Notice that by definition of the matrix P , there exist positive constants γ, γ' such that

$$\exp(\gamma'x^2) \leq K(x) \leq \exp(\gamma x^2) \quad \forall x \in \mathbb{R}^N.$$

When $N = 1$, the weight function K is given by

$$K(x) := \exp\left(\frac{|x|^2}{2\eta}\right).$$

We immediately deduce from [12] the following Proposition:

Proposition 2.1. *Let $F_{ini} \in L^\infty(\mathbb{R}^N) \cap L^2(K)$. Then the homogenized problem has a unique solution*

$$F \in \mathcal{C}([0, \infty), L^2(K)) \cap \mathcal{C}((0, \infty), H^2(K)) \cap \mathcal{C}^1((0, \infty), L^2(K))$$

such that $F|_{t=0} = F_{ini}$.

Moreover,

$$\lim_{t \rightarrow \infty} \|F(t) - F_M\|_{L^1(\mathbb{R}^N)} = 0,$$

where F_M is the unique stationary solution of the homogenized problem with mass $M = \int_{\mathbb{R}^N} F_{ini}$.

Consequently, the homogenized equations (2.15) and (2.13) are well posed. We conclude this section by stating a result on the construction of an approximate solution:

Definition 2.1. Let $F \in \mathcal{C}([0, \infty), L^2(K)) \cap \mathcal{C}((0, \infty), H^2(K))$. We define the approximate solution of (2.3) associated with F by

$$U^{\text{app}}[F](\tau, x; R) = U_0(\tau, x, z) + R^{-1}U_1(\tau, x, z) + R^{-2}V_2(\tau, x, z),$$

with $\tau \geq 0$, $x \in \mathbb{R}^N$, $R > 0$ and $z := Rx + c\frac{R^2-1}{2}$, and where

- U_0 is defined by (2.5);
- U_1 is defined by (2.8) if $N \geq 2$ and by (2.11) if $N = 1$;

- V_2 is defined by

$$\begin{aligned} -\Delta_z V_2 + \operatorname{div}(\alpha_1 V_2) &= (f_0(z) - 1) [-\partial_\tau F + \operatorname{div}_x(xF) + \Delta_x F] \\ &\quad + \langle \operatorname{div}_x((\alpha_1 - c)U_1) \rangle - \operatorname{div}_x((\alpha_1 - c)U_1) \\ &\quad + 2 \sum_{i=1}^N \frac{\partial^2 U_1}{\partial x_i \partial z_i} + \mathcal{A}_{NL} - \langle \mathcal{A}_{NL} \rangle. \end{aligned}$$

Notice that we do not require, in the above definition, that F is a solution of (2.13) or (2.15); hence the right-hand side in the equation on the term of order R^{-2} is slightly modified, so that the compatibility condition is satisfied and V_2 is well-defined. Of course, if F is a solution of (2.15) or (2.13), the equation on V_2 becomes (2.12) and $V_2 = U_2$.

We then have the following result:

Lemma 2.5. *1. Let $M \in \mathbb{R}$ be arbitrary. Define the function $U \in L^\infty([0, \infty) \cap \mathbb{R}^N)$ by*

$$U(\tau, x) := U^{app}[F_M](\tau, x; e^\tau).$$

Then U is a solution of

$$\begin{aligned} \partial_\tau U - \operatorname{div}_x(xU) - \Delta_x U + R \operatorname{div}_x((\alpha_1(z) - c)U) &= \\ &= -R^{N+1} \operatorname{div}_x \tilde{B}_1 \left(z, \frac{U}{R^N} \right) + U^{rem}, \end{aligned}$$

where the remainder term U^{rem} is such that there exist $C > 0$, $\gamma > 0$ such that

$$\|U^{rem}(\tau)\|_{L^1(e^{\gamma|x|^2})} + \|U^{rem}(\tau)\|_{L^\infty(\mathbb{R}^N)} \leq C e^{-\tau} \quad \forall \tau \geq 0.$$

2. *Let $F_{ini} \in L^\infty(\mathbb{R}^N) \cap L^2(K)$, and let $F \in \mathcal{C}([0, \infty), L^2(K))$ be the unique solution of the homogenized equation such that $F|_{t=0} = F_{ini}$. Let $\rho \in \mathcal{C}_0^\infty(\mathbb{R}^N)$ be a mollifying kernel ($\rho \geq 0$, $\int \rho = 1$), and let $F_\delta := F *_x \rho_\delta$, where $\rho_\delta = \delta^{-N} \rho(\cdot/\delta)$, for $\delta > 0$.*

Let $(\tau_n)_{n \geq 0}$ be a sequence of positive numbers such that $\lim_{n \rightarrow \infty} \tau_n = +\infty$. For $n \in \mathbb{N}$, $\delta > 0$, define the function u_n^δ by

$$u_n^\delta(\tau, x) = U^{app}[F_\delta](\tau, x; e^{\tau_n + \tau}), \quad x \in \mathbb{R}^N, \tau \geq 0.$$

Then u_n^δ satisfies, with $R_n = e^{\tau_n + \tau}$ and $z_n = R_n x + c \frac{R_n^2 - 1}{2}$,

$$\begin{aligned} \partial_\tau u_n^\delta - \operatorname{div}_x(xu_n^\delta) + R_n \operatorname{div}_x((\alpha_1(z_n) - c)u_n^\delta) - \Delta_x u_n^\delta &= \\ &= -R_n^{N+1} \operatorname{div}_x \tilde{B}_1 \left(z, \frac{u_n^\delta}{R_n^N} \right) + r_n^\delta, \end{aligned}$$

where the remainder term r_n^δ satisfies, for all $T > 0$,

$$\|r_n^\delta\|_{L^\infty([0,T],L^1(\mathbb{R}^N))} \leq \omega_T(\delta) + C_{\delta,T}e^{-\tau_n},$$

where $\omega_T : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a function depending only on T such that $\lim_{\delta \rightarrow 0^+} \omega_T(\delta) = 0$.

The proof of the above Lemma follows the calculations of the first paragraph; the proof is lengthy but straightforward, and is therefore left to the reader. The fact that U^{rem} has exponential decay is a consequence of Lemmas 2.3, 2.4.

3 Weighted L^2 bounds for the rescaled equation

As explained in the previous section, we choose to work with the rescaled equation (2.3) rather than with the original one (1.1). In fact, it can be easily checked that Theorem 1.1 is equivalent to the following Proposition:

Proposition 3.1. *Let $U_{ini} \in L^1(\mathbb{R}^N)$, and let $M := \int_{\mathbb{R}^N} U_{ini}$.*

Let $U \in \mathcal{C}([0, \infty), L^1(\mathbb{R}^N))$ be the unique solution of (2.3) with initial data $U|_{\tau=0} = U_{ini}$. Then

$$\lim_{\tau \rightarrow \infty} \int_{\mathbb{R}^N} \left| U(\tau, x) - f_0 \left(e^\tau x + c \frac{e^{2\tau} - 1}{2} \right) F_M(x) \right| dx = 0,$$

where the speed c is defined by (1.7), and $F_M \in L^1(\mathbb{R}^N)$ is the unique stationary solution of the homogenized equation (2.13), (2.15) with total mass M .

In turn, since the function $f_0 \in L^\infty(\mathbb{T}^N)$ is such that $\inf_{\mathbb{T}^N} f_0 > 0$, the above statement is equivalent to

$$\lim_{\tau \rightarrow \infty} \|V(\tau) - F_M\|_{L^1(\mathbb{R}^N)} = 0,$$

where the function $V = V(\tau, x)$ is defined by

$$V(\tau, x) := \frac{U(\tau, x)}{f_0 \left(e^\tau x + c \frac{e^{2\tau} - 1}{2} \right)}, \quad \tau \geq 0, \quad x \in \mathbb{R}^N. \quad (3.1)$$

The proof of Proposition 3.1 consists of essentially two steps: first, we prove compactness properties in $L^1(\mathbb{R}^N)$ for the family $(V(\tau))_{\tau \geq 0}$. To that end, we derive uniform bounds with respect to τ in weighted L^2 spaces; this step will be achieved in the current section. Then, we prove in the next section, using techniques inherited from dynamical systems theory, that the limit of any converging sequence $V(\tau_n)$ is equal to F_M . As emphasized in the introduction, the proof of convergence relies on rather abstract arguments, and thus does not yield any rate of convergence in general. However, when the flux A

is linear, the weighted L^2 bounds allow us to prove that the family $U(\tau)$ has uniformly bounded moments of order four, and thus (1.2) holds. As proved in [5], the convergence stated in Theorem 1.1 then takes place with algebraic rate.

The main result of this section is the following:

Proposition 3.2. *Let $U_{ini} \in L^1 \cap L^\infty(\mathbb{R}^N)$, and let $U \in \mathcal{C}([0, \infty), L^1(\mathbb{R}^N))$ be the unique solution of (2.3) with initial data $U|_{\tau=0} = U_{ini}$. Let $m > 2(N + 1)$ be arbitrary, and assume that*

$$\int_{\mathbb{R}^N} |U_{ini}(x)|^2 (1 + |x|^2)^{m/2} dx < +\infty.$$

Then there exists a constant $C_m > 0$ (depending only on m, N , and on the flux A) such that if $\|U_{ini}\|_{L^1(\mathbb{R}^N)} \leq C_m$, then

$$\begin{aligned} \sup_{\tau \geq 0} \int_{\mathbb{R}^N} |V(\tau, x)|^2 (1 + |x|^2)^{m/2} dx &< +\infty, \\ \sup_{\tau \geq 0} \int_{\tau}^{\tau+1} \int_{\mathbb{R}^N} |\nabla_x V(s, x)|^2 dx ds &< +\infty. \end{aligned} \tag{3.2}$$

As a consequence, there exists a sequence (τ_n) of positive numbers such that $\tau_n \in [n, n + 1]$ for all n , and such that the sequence $(V(\tau_n, x))_{n \geq 0}$ is compact in $L^1(\mathbb{R}^N)$.

Moreover, if the flux A is linear, then $C_m = +\infty$ for all $m > 2(N + 1)$.

Before proving the bounds (3.2), we explain how they entail the existence of a converging sequence. Thus we admit that (3.2) holds for the time being. First, for any $X \geq 1$, $\tau \geq 0$, we have

$$\begin{aligned} &\int_{|x| \geq X} |V(\tau, x)| dx \\ &\leq \left(\int_{|x| \geq X} |V(\tau, x)|^2 (1 + |x|^2)^{m/2} dx \right)^{1/2} \left(\int_{|x| \geq X} (1 + |x|^2)^{-m/2} dx \right)^{1/2} \\ &\leq CX^{(N-m)/2} \left(\sup_{\tau \geq 0} \int_{\mathbb{R}^N} |V(\tau, x)|^2 (1 + |x|^2)^{m/2} dx \right)^{1/2}. \end{aligned}$$

Since $m > N$, we infer that the family $\{V(\tau, x)\}_{\tau \geq 0}$ is equi-integrable.

Moreover, let $\mathcal{K} \subset \mathbb{R}^N$ be an arbitrary compact set, and let $h \in \mathbb{R}^N$ be arbitrary, with $|h| \leq 1$. Let

$$\tilde{\mathcal{K}} := \{x \in \mathbb{R}^N, d(x, \mathcal{K}) \leq 1\}.$$

The set $\tilde{\mathcal{K}}$ is clearly compact. Then

$$\begin{aligned} &\int_{\mathcal{K}} |V(\tau, x + h) - V(\tau, x)| dx \\ &\leq |h| \int_{\mathcal{K}} \int_0^1 |\nabla V|(\tau, x + \lambda h) d\lambda dx \\ &\leq |h| \int_{\tilde{\mathcal{K}}} |\nabla V(\tau, z)| dz \leq |h| |\tilde{\mathcal{K}}|^{1/2} \left(\int_{\mathbb{R}^N} |\nabla_x V(\tau, x)|^2 dx \right)^{1/2}. \end{aligned}$$

Now, for all $n \in \mathbb{N}$, there exists $\tau_n \in [n, n + 1]$ such that

$$\int_{\mathbb{R}^N} |\nabla_x V(\tau_n, x)|^2 dx \leq \int_n^{n+1} \int_{\mathbb{R}^N} |\nabla_x V(s, x)|^2 dx ds.$$

Consequently, there exists a constant C , depending only on \mathcal{K} and on the bounds on V in $L^2_{\text{loc}}([0, \infty), H^1)$, such that

$$\forall n \in \mathbb{N}, \quad \int_{\mathcal{K}} |V(\tau_n, x + h) - V(\tau_n, x)| dx \leq C|h|.$$

Hence the sequence $\{V(\tau_n, x)\}_{n \geq 0}$ is equi-continuous in $L^1(\mathbb{R}^N)$.

Notice also that

$$\sup_{n \geq 0} \|V(\tau_n)\|_{L^1(\mathbb{R}^N)} \leq \frac{1}{\inf_{\mathbb{T}^N} f_0} \sup_{n \geq 0} \|U(\tau_n)\|_{L^1(\mathbb{R}^N)} \leq \frac{\|U_{\text{ini}}\|_{L^1(\mathbb{R}^N)}}{\inf_{\mathbb{T}^N} f_0}.$$

Thus the sequence $\{V(\tau_n, x)\}_{n \geq 0}$ is bounded in $L^1(\mathbb{R}^N)$.

According to classical results of functional analysis (see for instance [6]), we infer that the sequence $(V(\tau_n))_{n \geq 0}$ is compact in L^1 .

The rest of the section is devoted to the proof of the bounds (3.2). We first prove that $V \in L^\infty_{\text{loc}}([0, \infty), L^2((1 + |x|^2)^{m/2}))$. Then, using the construction of approximate solutions of (2.3) performed in the previous section, we derive an energy inequality on the function V . Carefully controlling the non-linear terms appearing in this energy inequality, we are led to (3.2).

Before addressing the proof, we recall a result which will play a key role in several arguments: since $U_{\text{ini}} \in L^\infty \cap L^1(\mathbb{R}^N)$, there exists a positive constant C , depending only on the flux A and on $\|U_{\text{ini}}\|_1, \|U_{\text{ini}}\|_\infty$, such that

$$\|U(\tau)\|_{L^\infty(\mathbb{R}^N)} \leq C e^{N\tau}. \quad (3.3)$$

Indeed, performing backwards the parabolic scaling (2.2), it turns out that this inequality is equivalent to the boundedness of u in $L^\infty([0, \infty) \times \mathbb{R}^N)$, where u is the solution of (1.1) with initial data $v + U_{\text{ini}}$. And the L^∞ bound on u follows from Lemma A.1 in the Appendix.

First step: the family $V(\tau)$ is locally bounded in $L^2((1 + |x|^2)^{m/2})$.

This amounts in fact to proving that $U \in L^\infty_{\text{loc}}([0, \infty), L^2((1 + |x|^2)^{m/2}))$. Hence, multiply (2.3) by $U(\tau, x)(1 + |x|^2)^{m/2}$ and integrate with respect to the variable x . Always

with the notation $R = e^\tau$, $z = Rx + c\frac{R^2-1}{2}$, this leads to

$$\begin{aligned}
& \frac{1}{2} \frac{d}{d\tau} \int_{\mathbb{R}^N} |U(\tau, x)|^2 (1 + |x|^2)^{m/2} dx \\
&= - \int_{\mathbb{R}^N} |\nabla_x U|^2 (1 + |x|^2)^{m/2} dx - m \int_{\mathbb{R}^N} (x \cdot \nabla_x U) U (1 + |x|^2)^{-1+m/2} dx \\
&\quad - \frac{1}{2} \int_{\mathbb{R}^N} (x \cdot \nabla_x |U|^2) (1 + |x|^2)^{m/2} dx - m \int_{\mathbb{R}^N} |U|^2 |x|^2 (1 + |x|^2)^{-1+m/2} dx \\
&\quad - R^{N+1} \int_{\mathbb{R}^N} \left[B \left(z, \frac{U(\tau, x)}{R^N} \right) - c \frac{U(\tau, x)}{R^N} \right] \cdot \nabla_x U (1 + |x|^2)^{m/2} dy \\
&\quad - m R^{N+1} \int_{\mathbb{R}^N} \left[B \left(z, \frac{U(\tau, x)}{R^N} \right) - c \frac{U(\tau, x)}{R^N} \right] \cdot x U (1 + |x|^2)^{-1+m/2} dx.
\end{aligned} \tag{3.4}$$

Since $U(\tau)/R^N$ is bounded (see (3.3)), there exists a constant C such that

$$\left| B \left(Rx + c \frac{R^2-1}{2}, \frac{U(\tau, x)}{R^N} \right) - c \frac{U(\tau, x)}{R^N} \right| \leq C \frac{|U(\tau, x)|}{R^N}.$$

Moreover,

$$|x(1 + |x|^2)^{-1+m/2}|, \quad ||x|^2(1 + |x|^2)^{-1+m/2}| \leq (1 + |x|^2)^{m/2} \quad \forall x \in \mathbb{R}^N.$$

Hence, using the Cauchy-Schwarz inequality, we infer that the last two terms in (3.4) are bounded by

$$\frac{1}{4} \int_{\mathbb{R}^N} |\nabla_x U|^2 (1 + |x|^2)^{m/2} dx + CR^2 \int_{\mathbb{R}^N} |U|^2 (1 + |x|^2)^{m/2} dx.$$

On the other hand,

$$\begin{aligned}
& \left| \int_{\mathbb{R}^N} (x \cdot \nabla_x |U|^2) (1 + |x|^2)^{m/2} dx \right| \\
&= \left| \int_{\mathbb{R}^N} |U|^2 (N(1 + |x|^2)^{m/2} + mx^2(1 + |x|^2)^{-1+m/2}) dx \right| \\
&\leq C \int_{\mathbb{R}^N} |U|^2 (1 + |x|^2)^{m/2} dx.
\end{aligned}$$

Gathering all the terms, we deduce that there exists $c \in (0, 1)$, $C > 0$ such that

$$\begin{aligned}
& \frac{d}{d\tau} \int_{\mathbb{R}^N} |U(\tau, x)|^2 (1 + |x|^2)^{m/2} dx \\
&\leq -c \int_{\mathbb{R}^N} |\nabla_x U(\tau, x)|^2 (1 + |x|^2)^{m/2} dx + Ce^{2\tau} \int_{\mathbb{R}^N} |U(\tau, x)|^2 (1 + |x|^2)^{m/2} dx.
\end{aligned}$$

Using Gronwall's Lemma, we infer that

$$U \in L_{\text{loc}}^\infty([0, \infty), L^2((1 + |x|^2)^{m/2})), \quad \nabla_x U \in L_{\text{loc}}^2([0, \infty), L^2((1 + |x|^2)^{m/2})).$$

Second step: The energy inequality.

The idea here is the following: assume momentarily that the flux B is linear, that is, $\tilde{B}_1 = 0$. Let $\psi \in L^\infty([0, \infty) \times \mathbb{R}^N)$ be a solution of (2.3) such that $\psi(\tau, x) > 0$ for all τ, x . Then, according to [18], for any convex function $H \in \mathcal{C}^2(\mathbb{R})$, we have

$$\frac{d}{dt} \int_{\mathbb{R}^N} \psi(\tau, x) H \left(\frac{U(\tau, x)}{\psi(\tau, x)} \right) dx = - \int_{\mathbb{R}^N} H'' \left(\frac{U(\tau, x)}{\psi(\tau, x)} \right) \left| \nabla_x \left(\frac{U(\tau, x)}{\psi(\tau, x)} \right) \right|^2 dx.$$

Taking $H : x \in \mathbb{R} \mapsto x^2$, we infer that

$$\sup_{\tau \geq 0} \int_{\mathbb{R}^N} |U(\tau, x)|^2 \frac{dx}{\psi(\tau, x)} < +\infty.$$

Hence, if $\psi(\tau, x)$ behaves like $(1 + |x|^2)^{-m/2}$ for $|x|$ large, the L^2 bound in (3.2) is proved.

Thus the goal of this step is to build a positive function \tilde{U} , which behaves like $(1 + |x|^2)^{-m/2}$ for $|x|$ large, and which is an approximate solution of the linear part of (2.3), with remainder terms of order one. Using calculations similar to the ones led in [18], we then derive an inequality on the energy

$$\int_{\mathbb{R}^N} \left| \frac{U(\tau, x)}{\tilde{U}(\tau, x)} \right|^2 \tilde{U}(\tau, x).$$

From now on, we no longer assume that $\tilde{B}_1 = 0$.

The definition of \tilde{U} is inspired from the construction of an approximate solution in the previous paragraph. Precisely, we set

$$\tilde{U}(\tau, x) = f_0(z)h_m(x) + e^{-\tau} f_1(z) \cdot \nabla_x h_m(x), \quad \tau \geq 0, \quad x \in \mathbb{R}^N, \quad z = e^\tau x + c \frac{e^{2\tau} - 1}{2},$$

where the function $f_1 \in W^{1,\infty}(\mathbb{T}^N)^N$ is defined by (2.9), and where

$$h_m(x) := (1 + |x|^2)^{-m/2}.$$

Notice that even when $N = 1$, the structure of the function \tilde{U} is inspired from the linear case: there is no term of the form h_m^2 in \tilde{U} . The nonlinear term in the flux B will be treated independently.

Remember that $\inf_{\mathbb{T}^N} f_0 > 0$; since

$$\nabla_x h_m(x) = -m \frac{x}{1 + |x|^2} h_m, \quad x \in \mathbb{R}^N,$$

we deduce that there exists $\tau_0 > 0$ (depending on m), such that

$$0 < \frac{1}{2} f_0(z) h_m(x) \leq \tilde{U}(\tau, x) \leq 2 f_0(z) h_m(x) \quad \forall y \in \mathbb{R}^N, \quad \tau \geq \tau_0. \quad (3.5)$$

We now compute, for $\tau \geq \tau_0$, the rate of growth (or decay) of the energy $\int |U|^2 \tilde{U}^{-1}$. Using equation (2.3) and performing several integrations by parts, we obtain

$$\begin{aligned} & \frac{d}{d\tau} \int_{\mathbb{R}^N} \left| \frac{U(\tau, x)}{\tilde{U}(\tau, x)} \right|^2 \tilde{U}(\tau, x) dx \\ &= -2 \int_{\mathbb{R}^N} \left| \nabla_x \left(\frac{U(\tau, x)}{\tilde{U}(\tau, x)} \right) \right|^2 \tilde{U}(\tau, x) dx \\ &+ \int_{\mathbb{R}^N} \left| \frac{U(\tau, x)}{\tilde{U}(\tau, x)} \right|^2 \left[-\partial_\tau \tilde{U} + \Delta_x \tilde{U} + \operatorname{div}_x(x\tilde{U}) - e^\tau \operatorname{div}_x \left((\alpha_1(z) - c)\tilde{U} \right) \right] dx \\ &+ 2e^{(N+1)\tau} \int_{\mathbb{R}^N} \tilde{B}_1 \left(z, \frac{U(\tau, x)}{e^{N\tau}} \right) \cdot \nabla_x \left(\frac{U(\tau, x)}{\tilde{U}(\tau, x)} \right) dx. \end{aligned}$$

By definition of \tilde{U} , we have

$$\begin{aligned} & -\partial_\tau \tilde{U} + \Delta_x \tilde{U} + \operatorname{div}_x(x\tilde{U}) - e^\tau \operatorname{div}_x \left((\alpha_1(z) - c)\tilde{U} \right) \\ &= \operatorname{div}_x(xh_m)f_0(z) + f_0(z)\Delta_x h_m(x) + 2 \sum_{1 \leq i, j \leq N} \frac{\partial f_{1,i}(z)}{\partial z_j} \frac{\partial^2 h_m(x)}{\partial x_i \partial x_j} \\ &- \sum_{1 \leq i, j \leq N} [(\alpha_{1,i} - c_i)f_{1,j}](z) \frac{\partial^2 h_m(x)}{\partial x_i \partial x_j} \\ &+ e^{-\tau} \sum_{1 \leq i, j \leq N} f_{1,i}(z) \left[\frac{\partial}{\partial x_j} (x_j \partial_{x_i} h_m(x)) + \frac{\partial^3 h_m(x)}{\partial x_i \partial x_j^2} \right], \end{aligned}$$

where

$$z = e^\tau x + c \frac{e^{2\tau} - 1}{2}.$$

Notice that

$$\operatorname{div}_x(xh_m(x)) = (N - m)h_m(x) + \frac{m}{(1 + |x|^2)^{1 + \frac{m}{2}}},$$

and there exists a constant C (depending on m and N) such that for all $i, j \in \{1, \dots, N\}$,

$$\begin{aligned} & \left| \frac{\partial^2 h_m(x)}{\partial x_i \partial x_j} \right| \leq C \frac{1}{(1 + |x|^2)^{1 + \frac{m}{2}}}, \\ & |\nabla_x h_m(x)| + \left| |x| \frac{\partial^2 h_m(x)}{\partial x_i \partial x_j} \right| + \left| \frac{\partial^3 h_m(x)}{\partial x_i \partial x_j^2} \right| \leq Ch_m(x). \end{aligned}$$

Remember that $N - m < 0$ and that inequality (3.5) holds; hence we infer that for $\tau \geq \tau_0$ with τ_0 large enough,

$$\begin{aligned} & -\partial_\tau \tilde{U} + \Delta_x \tilde{U} + \operatorname{div}_x(x\tilde{U}) - e^\tau \operatorname{div}_x \left((\alpha_1(z) - c)\tilde{U} \right) \\ &\leq (N - m)f_0(z)h_m(x) + Ce^{-\tau}h_m(x) + C \frac{1}{(1 + |x|^2)^{1 + \frac{m}{2}}} \\ &\leq \frac{N - m}{4} \tilde{U}(\tau, x) + C \frac{1}{(1 + |x|^2)^{1 + \frac{m}{2}}}. \end{aligned}$$

On the other hand, since the flux \tilde{B}_1 is quadratic near the origin and $U/e^{N\tau}$ is bounded, we have

$$\begin{aligned} & \left| e^{(N+1)\tau} \int_{\mathbb{R}^N} \tilde{B}_1 \left(z, \frac{U(\tau, x)}{e^{N\tau}} \right) \cdot \nabla_x \left(\frac{U(\tau, x)}{\tilde{U}(\tau, x)} \right) dx \right| \\ & \leq C e^{(1-N)\tau} \int_{\mathbb{R}^N} |U(\tau, x)|^2 \left| \nabla_x \left(\frac{U(\tau, x)}{\tilde{U}(\tau, x)} \right) \right| dx. \end{aligned}$$

Gathering all the terms, we obtain

$$\frac{d}{d\tau} \int_{\mathbb{R}^N} \left| \frac{U}{\tilde{U}} \right|^2 \tilde{U} + \frac{m-N}{4} \int_{\mathbb{R}^N} \left| \frac{U}{\tilde{U}} \right|^2 \tilde{U} + 2 \int_{\mathbb{R}^N} \left| \nabla \left(\frac{U}{\tilde{U}} \right) \right|^2 \tilde{U} \quad (3.6)$$

$$\leq C \int_{\mathbb{R}^N} \left(\frac{U(\cdot, x)}{\tilde{U}(\cdot, x)} \right)^2 \frac{dx}{(1+|x|^2)^{1+\frac{m}{2}}} \quad (3.7)$$

$$+ C e^{(1-N)\tau} \int_{\mathbb{R}^N} |U|^2 \left| \nabla_x \left(\frac{U}{\tilde{U}} \right) \right|. \quad (3.8)$$

Notice that when the flux A is linear, the term (3.8) is zero.

Third step: control of the term (3.7).

Set $\phi := U/\tilde{U}$; then according to the first step,

$$\phi \in L_{\text{loc}}^\infty([\tau_0, \infty), L^2(h_m)) \cap L_{\text{loc}}^2([\tau_0, \infty), H^1(h_m)).$$

Moreover,

$$\nabla(\phi^2 h_m) = 2\phi h_m \nabla \phi + \phi^2 \nabla h_m;$$

since $|\nabla h_m| \leq m h_m$, we deduce that $\phi^2 h_m \in L_{\text{loc}}^1([\tau_0, \infty), W^{1,1}(\mathbb{R}^N))$, and thus, using Sobolev embeddings, $\phi^2 h_m \in L_{\text{loc}}^1([\tau_0, \infty), L^{p^*}(\mathbb{R}^N))$, where $p^* := N/(N-1)$ if $N \geq 2$, and $p^* = +\infty$ if $N = 1$. Additionally, the following inequality holds: there exists a constant C , depending only on N and m , such that for all $\tau \geq \tau_0$

$$\begin{aligned} \|\phi^2 h_m(\tau)\|_{L^{p^*}(\mathbb{R}^N)} & \leq C \|\nabla(\phi^2 h_m(\tau))\|_{L^1(\mathbb{R}^N)} \\ & \leq C \|\phi(\tau)\|_{L^2(h_m)} \|\nabla \phi(\tau)\|_{L^2(h_m)} + C \|\phi(\tau)\|_{L^2(h_m)}^2. \end{aligned}$$

We use the above inequality in order to control the term (3.7). First, let us write

$$\begin{aligned} & \int_{\mathbb{R}^N} |\phi(\tau, x)|^2 (1+|x|^2)^{-(1+\frac{m}{2})} dx \\ & = \int_{\mathbb{R}^N} (\phi^2(\tau) h_m)^a (|\phi(\tau)| h_m)^b, \end{aligned}$$

where the exponents a, b satisfy

$$\begin{cases} 2a + b = 2 \\ a \frac{m}{2} + b \frac{m}{2} = 1 + \frac{m}{2}, \end{cases}$$

which leads to $a = 1 - \frac{2}{m}$, $b = \frac{4}{m}$. Notice that $a, b \in (0, 1)$, provided m is large enough ($m > 4$, which is always satisfied if $m > 2(N + 1)$).

Then, using Hölder's inequality, we infer

$$\begin{aligned} & \int_{\mathbb{R}^N} |\phi(\tau, x)|^2 (1 + |x|^2)^{-(1+\frac{m}{2})} dx \\ & \leq \|\phi^2(\tau)h_m\|_{L^p(\mathbb{R}^N)}^a \|\phi(\tau)h_m\|_{L^1(\mathbb{R}^N)}^b, \end{aligned}$$

where the parameter p is given by

$$p = a(1 - b)^{-1} = \frac{1 - \frac{2}{m}}{1 - \frac{4}{m}}.$$

Notice that p is always larger than one. In order to be able to interpolate L^p between L^1 and L^{p^*} , p must also be smaller than p^* ; if $N = 1$, $p^* = \infty$, and thus we always have $p < p^*$. If $N \geq 2$, this condition amounts to $m > 2(N + 1)$; we assume that m always satisfies this assumption in the sequel.

Now, let $\theta \in (0, 1)$ such that

$$\frac{1}{p} = \frac{\theta}{1} + \frac{1 - \theta}{p^*};$$

using once again Hölder's inequality, we obtain

$$\begin{aligned} & \int_{\mathbb{R}^N} |\phi(\cdot, x)|^2 (1 + |x|^2)^{-(1+\frac{m}{2})} dx \\ & \leq \|\phi^2 h_m\|_{L^1(\mathbb{R}^N)}^{a\theta} \|\phi^2 h_m\|_{L^{p^*}(\mathbb{R}^N)}^{a(1-\theta)} \|\phi h_m\|_{L^1(\mathbb{R}^N)}^b \\ & \leq C \|\phi^2 h_m\|_{L^1(\mathbb{R}^N)}^{a\theta + a\frac{1-\theta}{2}} \|\nabla \phi\|_{L^2(h_m)}^{a(1-\theta)} \|\phi h_m\|_{L^1(\mathbb{R}^N)}^b \\ & \quad + C \|\phi^2 h_m\|_{L^1(\mathbb{R}^N)}^a \|\phi h_m\|_{L^1(\mathbb{R}^N)}^b. \end{aligned}$$

If $N = 1$, then $\theta = p^{-1}$, and straightforward computations lead to

$$a\theta + a\frac{1-\theta}{2} = 1 - \frac{3}{m}, \quad a(1-\theta) = \frac{2}{m}.$$

Hence, using Young's inequality, we deduce that for all $\lambda > 0$, there exists a constant C_λ such that

$$\begin{aligned} & \int_{\mathbb{R}^N} |\phi(\tau, x)|^2 (1 + |x|^2)^{-(1+\frac{m}{2})} dx \\ & \leq \lambda \|\phi^2(\tau)h_m\|_{L^1(\mathbb{R})} + \lambda \|\nabla \phi(\tau)\|_{L^2(h_m)}^2 + C_\lambda \|\phi(\tau)h_m\|_{L^1(\mathbb{R})}^2. \end{aligned} \quad (3.9)$$

If $N \geq 2$, the calculations are similar and lead to

$$a\theta + a\frac{1-\theta}{2} = 1 - \frac{N+2}{m}, \quad a(1-\theta) = \frac{2N}{m}.$$

Hence (3.9) is also valid in this case.

Using inequality (3.5) and choosing the parameter λ small enough leads eventually to

$$(3.7) \leq \frac{m-N}{16} \int_{\mathbb{R}^N} \left| \frac{U(\tau)}{\tilde{U}(\tau)} \right|^2 \tilde{U}(\tau) + \frac{1}{2} \int_{\mathbb{R}^N} \left| \nabla \left(\frac{U(\tau)}{\tilde{U}(\tau)} \right) \right|^2 \tilde{U}(\tau) \quad (3.10)$$

$$+ C \left(\int_{\mathbb{R}^N} |U(\tau, x)| dx \right)^2$$

for all $\tau \geq \tau_0$.

Fourth step: control of the term (3.8).

Remark 3.1. We recall that (3.8)=0 if the flux A is linear. Hence this step is required only in the nonlinear case.

Using inequality (3.3), we infer that there exists a constant C such that

$$(3.8) \leq C \int_{\mathbb{R}^N} |U(\tau, x)|^{1+\frac{1}{N}} \left| \nabla_x \left(\frac{U(\tau, x)}{\tilde{U}(\tau, x)} \right) \right| dx.$$

From now on, we treat the cases $N = 1$, $N = 2$, and $N \geq 3$ separately, and we set $\phi = U/\tilde{U}$.

- If $N = 1$, we have, for all $\tau \geq \tau_0$,

$$\begin{aligned} & \int_{\mathbb{R}^N} |U(\tau, x)|^{1+\frac{1}{N}} \left| \nabla_x \left(\frac{U(\tau, x)}{\tilde{U}(\tau, x)} \right) \right| dx \\ & \leq C \int_{\mathbb{R}} (\partial_x \phi(\tau) h_m^{1/2}) (\phi(\tau) h_m)^{1/2} (\phi^{3/2}(\tau) h_m) \\ & \leq C \|\partial_x \phi(\tau)\|_{L^2(h_m)} \|U(\tau)\|_{L^1(\mathbb{R})}^{1/2} \|\phi^{3/2}(\tau) h_m\|_{L^\infty(\mathbb{R})} \\ & \leq C \|\partial_x \phi(\tau)\|_{L^2(h_m)} \|U(\tau)\|_{L^1(\mathbb{R})}^{1/2} \|\partial_x (\phi^{3/2}(\tau) h_m)\|_{L^1(\mathbb{R})}. \end{aligned}$$

Moreover,

$$\partial_x (\phi^{3/2} h_m) = \frac{3}{2} \phi^{1/2} h_m \partial_x \phi + \phi^{3/2} \partial_x h_m,$$

and thus

$$\|\partial_x (\phi^{3/2} h_m)\|_{L^1(\mathbb{R})} \leq C \|\phi h_m\|_{L^1}^{1/2} (\|\partial_x \phi\|_{L^2(h_m)} + \|\phi\|_{L^2(h_m)}).$$

Eventually, we obtain, using once again (3.5),

$$(3.8) \leq C \|U(\tau)\|_{L^1(\mathbb{R})} \left[\int_{\mathbb{R}} \left| \nabla \left(\frac{U(\tau)}{\tilde{U}(\tau)} \right) \right|^2 \tilde{U}(\tau) + \int_{\mathbb{R}} \left| \frac{U(\tau)}{\tilde{U}(\tau)} \right|^2 \tilde{U}(\tau) \right].$$

Since

$$\|U(\tau)\|_{L^1(\mathbb{R})} \leq \|U_{ini}\|_{L^1(\mathbb{R})} \quad \forall \tau \geq 0,$$

we infer that if $\|U_{ini}\|_{L^1(\mathbb{R})}$ is sufficiently small, then

$$(3.8) \leq \frac{m-N}{16} \int_{\mathbb{R}^N} \left| \frac{U(\tau)}{\tilde{U}(\tau)} \right|^2 \tilde{U}(\tau) + \frac{1}{2} \int_{\mathbb{R}^N} \left| \nabla \left(\frac{U(\tau)}{\tilde{U}(\tau)} \right) \right|^2 \tilde{U}(\tau). \quad (3.11)$$

- If $N = 2$, using the Sobolev embedding $W^{1,1}(\mathbb{R}^2) \subset L^2(\mathbb{R}^2)$, we obtain, for $\tau \geq \tau_0$,

$$\begin{aligned} \int_{\mathbb{R}^N} |U(\cdot, x)|^{1+\frac{1}{N}} \left| \nabla_x \left(\frac{U(\cdot, x)}{\tilde{U}(\cdot, x)} \right) \right| dx &\leq C \int_{\mathbb{R}^2} (|\phi|^{3/2} h_m) (|\nabla_x \phi| h_m^{1/2}) \\ &\leq C \| |\phi|^{3/2} h_m \|_{L^2(\mathbb{R}^2)} \| \nabla_x \phi \|_{L^2(h_m)} \\ &\leq C \| \nabla (|\phi|^{3/2} h_m) \|_{L^1(\mathbb{R}^2)} \| \nabla_x \phi \|_{L^2(h_m)}. \end{aligned}$$

As is the case $N = 1$, we have

$$\| \nabla (|\phi|^{3/2} h_m) \|_{L^1(\mathbb{R}^2)} \leq C \| \phi h_m \|_{L^1(\mathbb{R}^2)}^{1/2} (\| \partial_x \phi \|_{L^2(h_m)} + \| \phi \|_{L^2(h_m)}).$$

Hence we are led to

$$(3.8) \leq C \| U(\tau) \|_{L^1(\mathbb{R}^2)}^{1/2} \left[\int_{\mathbb{R}^2} \left| \nabla \frac{U(\tau)}{\tilde{U}(\tau)} \right|^2 \tilde{U}(\tau) + \int_{\mathbb{R}^2} \left| \frac{U(\tau)}{\tilde{U}(\tau)} \right|^2 \tilde{U}(\tau) \right].$$

Following exactly the same argument as in the case $N = 1$, we deduce that if $\|U_{ini}\|_{L^1(\mathbb{R}^2)}$ is sufficiently small, then (3.11) holds.

- If $N \geq 3$, we have, for $\tau \geq \tau_0$,

$$\begin{aligned} &\int_{\mathbb{R}^N} |U(\cdot, x)|^{1+\frac{1}{N}} \left| \nabla_x \left(\frac{U(\cdot, x)}{\tilde{U}(\cdot, x)} \right) \right| dx \\ &\leq C \int_{\mathbb{R}^N} |\phi|^{1+\frac{1}{N}} |\nabla_x \phi| h_m^{1+\frac{1}{N}} \\ &\leq C \int_{\mathbb{R}^N} (|\phi| h_m)^{1/N} (|\nabla_x \phi| h_m^{1/2}) (|\phi| h_m^{1/2}) \\ &\leq C \| \phi h_m \|_{L^1(\mathbb{R}^N)}^{1/N} \| \nabla_x \phi \|_{L^2(h_m)} \| |\phi| h_m^{1/2} \|_{L^p(\mathbb{R}^N)}, \end{aligned}$$

where the parameter p is such that

$$\frac{1}{N} + \frac{1}{2} + \frac{1}{p} = 1,$$

i.e. $p = (2N)/(N-2)$. Using the Sobolev embedding $H^1(\mathbb{R}^N) \subset L^p(\mathbb{R}^N)$, we have

$$\begin{aligned} \| |\phi| h_m^{1/2} \|_{L^p(\mathbb{R}^N)} &\leq C \| \nabla_x (|\phi| h_m^{1/2}) \|_{L^2(\mathbb{R}^N)} \\ &\leq C (\| \nabla \phi \|_{L^2(h_m)} + \| \phi \|_{L^2(h_m)}). \end{aligned}$$

Thus, once again, we obtain

$$(3.8) \leq C \|U(\tau)\|_{L^1(\mathbb{R}^N)}^{1/N} \left[\int_{\mathbb{R}^N} \left| \nabla \left(\frac{U(\tau)}{\tilde{U}(\tau)} \right) \right|^2 \tilde{U}(\tau) + \int_{\mathbb{R}^2} \left| \frac{U(\tau)}{\tilde{U}(\tau)} \right|^2 \tilde{U}(\tau) \right],$$

and thus (3.11) holds as long as $\|U_{ini}\|_{L^1(\mathbb{R}^N)}$ is not too large.

Gathering inequalities (3.6), (3.10) and (3.11), we infer that if $\|U_{ini}\|_{L^1(\mathbb{R}^N)}$ is sufficiently small, then for all $\tau \geq \tau_0$,

$$\begin{aligned} \frac{d}{d\tau} \int_{\mathbb{R}^N} \left| \frac{U(\tau)}{\tilde{U}(\tau)} \right|^2 \tilde{U}(\tau) + \frac{m-N}{8} \int_{\mathbb{R}^N} \left| \frac{U(\tau)}{\tilde{U}(\tau)} \right|^2 \tilde{U}(\tau) + \int_{\mathbb{R}^N} \left| \nabla \left(\frac{U(\tau)}{\tilde{U}(\tau)} \right) \right|^2 \tilde{U}(\tau) \\ \leq C \left(\int_{\mathbb{R}^N} |U(\tau, x)| dx \right)^2 \leq C \|U_{ini}\|_{L^1(\mathbb{R}^N)}^2. \end{aligned} \quad (3.12)$$

Fifth step: Conclusion.

Let $C_1 := (m-N)/8$, $C_2 := C \|U_{ini}\|_{L^1(\mathbb{R}^N)}^2$. Using a Gronwall type argument, we deduce that for all $\tau \geq \tau_0$, we have

$$\begin{aligned} \int_{\mathbb{R}^N} \left| \frac{U(\tau)}{\tilde{U}(\tau)} \right|^2 \tilde{U}(\tau) + \int_{\tau_0}^{\tau} e^{-C_1(\tau-s)} \left(\int_{\mathbb{R}^N} \left| \nabla \left(\frac{U(s)}{\tilde{U}(s)} \right) \right|^2 \tilde{U}(s) \right) ds \\ \leq e^{-C_1(\tau-\tau_0)} \int_{\mathbb{R}^N} \left| \frac{U(\tau_0)}{\tilde{U}(\tau_0)} \right|^2 \tilde{U}(\tau_0) + \frac{C_2}{C_1} \\ \leq C \int_{\mathbb{R}^N} |U(\tau_0, x)|^2 (1 + |x|^2)^{m/2} + \frac{C_2}{C_1}. \end{aligned}$$

Using (3.5), we infer

$$\begin{aligned} \sup_{\tau \geq \tau_0} \int_{\mathbb{R}^N} |U(\tau, x)|^2 (1 + |x|^2)^{m/2} \leq C \int_{\mathbb{R}^N} |U(\tau_0, x)|^2 (1 + |x|^2)^{m/2} + C \frac{C_2}{C_1}, \\ \sup_{\tau \geq \tau_0} \int_{\tau}^{\tau+1} \int_{\mathbb{R}^N} \left| \nabla \left(\frac{U(s)}{\tilde{U}(s)} \right) \right|^2 h_m ds \leq C \int_{\mathbb{R}^N} |U(\tau_0, x)|^2 (1 + |x|^2)^{m/2} + C \frac{C_2}{C_1}. \end{aligned}$$

Hence $U \in L^\infty([0, \infty), L^2((1 + |x|^2)^{m/2}))$. Since f_0 is bounded away from zero, the L^2 bound on V follows.

Concerning the bound on $\nabla_x V$, notice that

$$V(\tau, x) = \frac{U(\tau, x)}{\tilde{U}(\tau, x)} \left(h_m(x) + e^{-\tau} \frac{f_1(z)}{f_0(z)} \cdot \nabla_x h_m(x) \right),$$

and thus

$$|\nabla_x V(\tau, x)| \leq C h_m(x) \left(\left| \nabla_x \left(\frac{U(\tau, x)}{\tilde{U}(\tau, x)} \right) \right| + \left| \frac{U(\tau, x)}{\tilde{U}(\tau, x)} \right| \right).$$

Consequently, for all $\tau \geq 0$

$$\int_{\mathbb{R}^N} |\nabla_x V(\tau, x)|^2 (1 + |x|^2)^{m/2} dx \leq C \left\| \frac{U(\tau)}{\tilde{U}(\tau)} \right\|_{H^1(h_m)}^2,$$

which leads to the bound on $\nabla_x V$. Notice that in fact, we recover

$$\nabla_x V \in L_{\text{loc}}^2([0, \infty), L^2((1 + |x|^2)^{m/2})).$$

Hence Proposition 3.2 is proved.

Let us now conclude this section by explaining how the bound (1.2) on the moments of order four follows from (3.2). Let $U_{ini} \in L^2(h_m^{-1})$, with $m > 2(N + 2)$ sufficiently large. Then we have proved that $U \in L^\infty([0, \infty), L^2(h_m^{-1}))$, with no restriction on the size of $\|U_{ini}\|_{L^1(\mathbb{R}^N)}$ in the linear case. Now, for all $\tau \geq 0$, using a simple Hölder inequality, we infer that

$$\int_{\mathbb{R}^N} |U(\tau, x)| |x|^4 dx \leq \|U(\tau)\|_{L^2(h_m^{-1})} \left(\int_{\mathbb{R}^N} |x|^8 (1 + |x|^2)^{-m/2} dx \right)^{1/2}.$$

Hence, if $m > N + 8$, we deduce that $U \in L^\infty([0, \infty), L^1(|x|^4))$; going back to the original variables, this entails that (1.2) is satisfied. Thus the convergence result (1.3) holds if the flux A is linear, and Proposition 1.1 is proved.

4 Long-time behaviour

This section is devoted to the rest of the proof of Theorem 1.1. The idea is to use the L^1 compactness proved in the previous section (see Proposition 3.2) together with techniques from dynamical systems theory. This type of proof was initiated by S. Osher and J. Ralston in [19], in which the authors proved the L^1 stability of travelling waves for a quasilinear parabolic equation. Their arguments were then adapted successfully to various kinds of problems in the context of scalar conservation laws (see for instance the review in [21]).

In the present study, our scheme of proof is in fact closely related to the one of M. Escobedo and E. Zuazua in [12]; indeed, the idea is to apply the dynamical systems tools to the rescaled parabolic system (2.3) rather than the original conservation law (1.1). The main difference with [12] lies in the presence of highly oscillating coefficients in (2.3); thus it is necessary to work simultaneously with the homogenized equation (2.13)-(2.15) and with the oscillating one.

Let us now introduce some notation and definitions. First, we denote by S_τ ($\tau \geq 0$) the semi-group associated with the homogenized equation, that is equation (2.15) if $N = 1$, and equation (2.13) if $N \geq 2$. According to Proposition 2.1, the semi-group S_τ is

well-defined in $L^\infty(\mathbb{R}^N) \cap L^2(K)$, where K is the weight function defined by (2.19); additionally, the L^1 contraction property holds, namely

$$\|S_\tau F_1 - S_\tau F_2\|_{L^1(\mathbb{R}^N)} \leq \|F_1 - F_2\|_{L^1(\mathbb{R}^N)} \quad \forall \tau \geq 0, \forall F_1, F_2 \in L^\infty(\mathbb{R}^N) \cap L^2(K).$$

Hence S_τ can be extended on $L^1(\mathbb{R}^N)$.

We also define the ω -limit set associated with a given function $U_{ini} \in L^1(\mathbb{R}^N)$: recalling the definition of the function V (see (3.1)), we set

$$\Omega[U_{ini}] := \left\{ \bar{V} \in L^1(\mathbb{R}^N), \exists \tau_n \rightarrow \infty, V(\tau_n) \rightarrow \bar{V} \text{ in } L^1(\mathbb{R}^N) \right\}, \quad (4.1)$$

where the function U in (3.1) is the unique solution of (2.3) with initial data U_{ini} . When there is no ambiguity, we will simply write Ω instead of $\Omega[U_{ini}]$.

Notice that $V(\tau_n)$ converges towards \bar{V} in L^1 if and only if

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} \left| U(\tau_n, x) - f_0 \left(e^{\tau_n} x + c \frac{e^{2\tau_n} - 1}{2} \right) \bar{V}(x) \right| dx = 0.$$

This equivalence will be used repeatedly throughout the section.

The organisation of this section is the following: we first introduce a ‘‘quasi-Liapunov function’’ for the semi-group associated with equation (2.3). We then prove that Proposition 3.1 holds when the initial data U_{ini} has a sufficiently small L^1 norm. Eventually, we prove Proposition 3.1 in the general case.

4.1 A quasi-Liapunov function

Let us first recall the definition of a Liapunov function (see [17]):

Definition 4.1. Let \mathcal{X} be a Banach space, and let $\pi : [0, \infty) \times \mathcal{X} \rightarrow \mathcal{X}$ be a *dynamical system* on \mathcal{X} , i.e. π satisfies the following:

- (i) $\pi(0, x) = x$ for all $x \in \mathcal{X}$;
- (ii) **Semi-group property:** For all $s, t \geq 0$, for all $x \in \mathcal{X}$, $\pi(t + s, x) = \pi(t, \pi(s, x))$;
- (iii) π is continuous on $[0, \infty) \times \mathcal{X}$.

Let $\mathcal{V} : \mathcal{X} \rightarrow \mathbb{R}$. \mathcal{V} is said to be a *Liapunov function* if

- (i) \mathcal{V} is continuous;
- (ii) $\dot{\mathcal{V}}(x) \leq 0$ for all $x \in \mathcal{X}$, where

$$\dot{\mathcal{V}}(x) := \liminf_{t \rightarrow 0^+} \frac{\mathcal{V}(\pi(t, x)) - \mathcal{V}(x)}{t}.$$

Liapunov functions are crucial in the analysis of the stability of dynamical systems. In the present context, because of the different scales involved, it seems difficult to find a suitable Liapunov function, and we thus extend the previous definition:

Definition 4.2. Let \mathcal{X} be a Banach space, and let $t \in [0, \infty) \mapsto x_t \in \mathcal{X}$ be a continuous curve in \mathcal{X} . Let $\mathcal{H} : [0, \infty) \times \mathcal{X} \rightarrow \mathbb{R}$ be continuous. We say that \mathcal{H} is a *quasi-Liapunov function* for the trajectory $(x_t)_{t \geq 0}$ if the following properties hold:

- (i) The family $\mathcal{H}(t, x_t)$ ($t \geq 0$) is bounded in \mathbb{R} ;
- (ii) There exists a function $\psi : [0, \infty) \rightarrow [0, \infty)$ such that $\lim_{t \rightarrow \infty} \psi(t) = 0$ and

$$\forall t \geq 0, \quad \sup_{s \geq t} (\mathcal{H}(s, x_s) - \mathcal{H}(t, x_t)) \leq \psi(t). \quad (4.2)$$

In other words, we drop the semi-group property associated with the dynamical system π , and we focus on a particular trajectory; moreover, inequality (4.2) means that $\mathcal{H}(s, x_s)$ is “almost decreasing” with respect to s for large s . Notice that a Lyapunov function associated with a semi-group π is decreasing along any trajectory $\pi(s, x)$ ($s \geq 0$).

We then have the following result:

Lemma 4.1. *Let \mathcal{X} be a Banach space, and let $t \in [0, \infty) \mapsto x_t \in \mathcal{X}$ be a continuous curve in \mathcal{X} . Let $\mathcal{H} : [0, \infty) \times \mathcal{X} \rightarrow \mathbb{R}$ be a quasi-Liapunov function for $(x_t)_{t \geq 0}$. Then $\mathcal{H}(t, x_t)$ has a finite limit as $t \rightarrow \infty$.*

Proof. First, since $\mathcal{H}(t, x_t)$ is bounded for $t \in [0, \infty)$, the quantities

$$\underline{H} := \liminf_{t \rightarrow \infty} \mathcal{H}(t, x_t), \quad \overline{H} := \limsup_{t \rightarrow \infty} \mathcal{H}(t, x_t)$$

are well-defined and belong to \mathbb{R} , with $\underline{H} \leq \overline{H}$.

Let $\varepsilon > 0$ arbitrary. There exists $t_\varepsilon > 0$ such that

$$\psi(t) \leq \varepsilon \quad \forall t \geq t_\varepsilon.$$

By definition of \underline{H} , there exists $s_\varepsilon \geq t_\varepsilon$ such that

$$|\mathcal{H}(s_\varepsilon, x_{s_\varepsilon}) - \underline{H}| \leq \varepsilon.$$

Since \mathcal{H} is a quasi-Liapunov function, for all $s \geq s_\varepsilon$, we have

$$\begin{aligned} \mathcal{H}(s, x_s) &\leq \mathcal{H}(s_\varepsilon, x_{s_\varepsilon}) + \psi(s_\varepsilon) \\ &\leq \underline{H} + 2\varepsilon. \end{aligned}$$

Hence

$$\overline{H} \leq \underline{H} + 2\varepsilon \quad \forall \varepsilon > 0,$$

and $\overline{H} = \underline{H}$. Thus the quantity $\mathcal{H}(t, x_t)$ has a finite limit as $t \rightarrow \infty$.

□

We now apply this notion to the present context:

Lemma 4.2. *Let $M \in \mathbb{R}$ be arbitrary, and let $U_{ini} \in L^1(\mathbb{R}^N)$. For $\tau \geq 0$ and $u \in L^1(\mathbb{R}^N)$, define the function \mathcal{H} by*

$$\mathcal{H}(\tau, u) := \int_{\mathbb{R}^N} |u(x) - U^{app}[F_M](\tau, x; e^\tau)| dx,$$

where the function U^{app} was introduced in Definition 2.1.

Let $U \in \mathcal{C}([0, \infty), L^1(\mathbb{R}^N))$ be the solution of (2.3) with initial data $U_{\tau=0} = U_{ini}$.

Then \mathcal{H} is a quasi-Liapunov function for the trajectory $\{U(\tau)\}_{\tau \geq 0}$ in $L^1(\mathbb{R}^N)$. As a consequence, the function

$$\tau \in [0, \infty) \mapsto \int_{\mathbb{R}^N} |U(\tau, x) - f_0(z)F_M(x)| dx, \quad \text{with } z = e^\tau x + c \frac{e^{2\tau} - 1}{2},$$

converges as $\tau \rightarrow \infty$.

Proof. This property is an easy consequence of the first point in Lemma 2.5; indeed, according to Lemma 2.5, there exists a constant C , depending only on N and M , such that

$$\frac{d}{d\tau} \mathcal{H}(\tau, U(\tau)) = \frac{d}{d\tau} \|U(\tau) - U^{app}[F_M](\tau)\|_{L^1(\mathbb{R}^N)} \leq C e^{-\tau}.$$

Consequently, for all $\tau' \geq \tau \geq 0$, we have

$$\mathcal{H}(\tau', U(\tau')) - \mathcal{H}(\tau, U(\tau)) \leq C(e^{-\tau} - e^{-\tau'}) \leq C e^{-\tau}.$$

Thus property (ii) of Definition 4.2 is satisfied. Additionally, notice that

$$\begin{aligned} 0 \leq \mathcal{H}(\tau, U(\tau)) &\leq \|U(\tau)\|_{L^1(\mathbb{R}^N)} + \|f_0\|_{L^\infty(\mathbb{T}^N)} \|F_M\|_{L^1(\mathbb{R}^N)} \\ &\quad + C e^{-\tau} \left(\|\nabla F_M\|_{L^1(\mathbb{R}^N)} + \|F_M\|_{L^2(\mathbb{R}^N)}^2 \right) \\ &\quad + C e^{-2\tau} \left(\|F_M\|_{W^{2,1}(\mathbb{R}^N)} + \|F_M\|_{L^3(\mathbb{R}^N)}^3 + \|\nabla F_M\|_{L^2(\mathbb{R}^N)}^2 \right) \\ &\leq \|U_{ini}\|_{L^1(\mathbb{R}^N)} + C. \end{aligned}$$

Whence $\mathcal{H}(\tau, U(\tau))$ is bounded for $\tau \in [0, \infty)$. Additionally the continuity of \mathcal{H} on $[0, \infty) \times L^1$ is obvious. Consequently \mathcal{H} is a quasi-Liapunov function for the trajectory $U(\tau)$. According to Lemma 4.1, $\mathcal{H}(\tau, U(\tau))$ admits a finite limit as $\tau \rightarrow \infty$. Furthermore, we have

$$\|U(\tau) - f_0(z)F_M\|_{L^1(\mathbb{R}^N)} = \|U(\tau) - U^{app}[F_M](\tau) + e^{-\tau}U_1 + e^{-2\tau}U_2\|_{L^1(\mathbb{R}^N)},$$

where U_1 and U_2 are defined by (2.8)-(2.11) and (2.12) respectively. Hence for all $\tau \geq 0$, there holds

$$\mathcal{H}(\tau, U(\tau)) - C e^{-\tau} \leq \|U(\tau) - f_0(z)F_M\|_{L^1(\mathbb{R}^N)} \leq \mathcal{H}(\tau, U(\tau)) + C e^{-\tau},$$

where the constant C depends only on $W^{s,p}$ bounds on F_M . Thus the function

$$\tau \mapsto \|U(\tau) - f_0(z)F_M\|_{L^1(\mathbb{R}^N)}$$

converges as $\tau \rightarrow \infty$, and

$$\lim_{\tau \rightarrow \infty} \|U(\tau) - f_0(z)F_M\|_{L^1(\mathbb{R}^N)} = \lim_{\tau \rightarrow \infty} \mathcal{H}(\tau, U(\tau)).$$

□

Definition 4.3. Let $U_{ini} \in L^1(\mathbb{R}^N)$ be arbitrary, and let $M := \int_{\mathbb{R}^N} U_{ini}$. Let U be the solution of (2.3) with initial data $U|_{t=0} = U_{ini}$.

We define the number $\ell(U_{ini})$ by

$$\ell(U_{ini}) := \lim_{\tau \rightarrow \infty} \int_{\mathbb{R}^N} |U(\tau, x) - f_0(z)F_M(x)| dx, \quad \text{with } z = e^\tau x + c \frac{e^{2\tau} - 1}{2}.$$

Notice that Proposition 3.1 is equivalent to

$$\ell(U_{ini}) = 0 \quad \forall U_{ini} \in L^1(\mathbb{R}^N).$$

Classically, we now derive a continuity property for the function ℓ :

Lemma 4.3. *The function*

$$U \in L^1(\mathbb{R}^N) \mapsto \ell(U) \in \mathbb{R}$$

is Lipschitz continuous.

Proof. Let $U_{ini}^{(1)}, U_{ini}^{(2)} \in L^1(\mathbb{R}^N)$, and let $M^{(i)} = \int_{\mathbb{R}^N} U_{ini}^{(i)}$ for $i = 1, 2$. We denote by $U^{(i)} \in \mathcal{C}([0, \infty), L^1(\mathbb{R}^N))$ the solution of (2.3) with initial data $U_{ini}^{(i)}$. Then for all $\tau \geq 0$, the L^1 contraction principle ensures that

$$\|U^{(1)}(\tau) - U^{(2)}(\tau)\|_{L^1(\mathbb{R}^N)} \leq \|U_{ini}^{(1)} - U_{ini}^{(2)}\|_{L^1(\mathbb{R}^N)}.$$

Hence, for all $\tau \geq 0$, we have

$$\begin{aligned} & \left| \int_{\mathbb{R}^N} |U^{(1)}(\tau, x) - f_0(z)F_{M^{(1)}}(x)| dx - \int_{\mathbb{R}^N} |U^{(2)}(\tau, x) - f_0(z)F_{M^{(2)}}(x)| dx \right| \\ & \leq \|U^{(1)}(\tau) - U^{(2)}(\tau)\|_{L^1(\mathbb{R}^N)} + \|f_0\|_{L^\infty(\mathbb{T}^N)} \|F_{M^{(1)}} - F_{M^{(2)}}\|_{L^1(\mathbb{R}^N)}. \end{aligned}$$

According to Lemma A.2 in the Appendix,

$$\|F_{M^{(1)}} - F_{M^{(2)}}\|_{L^1(\mathbb{R}^N)} = |M^{(1)} - M^{(2)}| \leq \|U_{ini}^{(1)} - U_{ini}^{(2)}\|_{L^1(\mathbb{R}^N)}.$$

Eventually, we obtain, for all $\tau \geq 0$,

$$\begin{aligned} \left| \int_{\mathbb{R}^N} |U^{(1)}(\tau, x) - f_0(z)F_{M^{(1)}}(x)| dx - \int_{\mathbb{R}^N} |U^{(2)}(\tau, x) - f_0(z)F_{M^{(2)}}(x)| dx \right| \leq \\ \leq \left(1 + \|f_0\|_{L^\infty(\mathbb{T}^N)}\right) \left\| U_{ini}^{(1)} - U_{ini}^{(2)} \right\|_{L^1(\mathbb{R}^N)}, \end{aligned}$$

and thus, passing to the limit,

$$\left| \ell \left(U_{ini}^{(1)} \right) - \ell \left(U_{ini}^{(2)} \right) \right| \leq \left(1 + \|f_0\|_{L^\infty(\mathbb{T}^N)}\right) \left\| U_{ini}^{(1)} - U_{ini}^{(2)} \right\|_{L^1(\mathbb{R}^N)}.$$

Hence ℓ is a Lipschitz continuous function. □

4.2 Analysis of the ω -limit set

Proposition 4.1. *Let $U_{ini} \in L^1(\mathbb{R}^N)$, and set*

$$M := \int_{\mathbb{R}^N} U_{ini}.$$

Assume that the ω -limit set Ω associated with U_{ini} is non-empty (see (4.1)). Then the following properties hold:

(i) *For all $\bar{V} \in \Omega$,*

$$\int_{\mathbb{R}^N} \bar{V} = M;$$

(ii) *$S_\tau \Omega \subset \Omega$ for all $\tau \geq 0$;*

(iii) *For all $\bar{V} \in \Omega$, we have*

$$\left\| \bar{V} - F_M \right\|_{L^1(\mathbb{R}^N)} = \ell(U_{ini}).$$

Proof. Throughout the proof, we denote by U the unique solution of equation (2.3) with initial data U_{ini} .

Property (i) is quite straightforward: indeed, conservation of mass for the equation (2.3) implies that

$$\int_{\mathbb{R}^N} U(\tau) = M \quad \forall \tau \geq 0.$$

If $\bar{V} \in \Omega$, then there exists a sequence $(\tau_n)_{n \geq 0}$ such that

$$\lim_{n \rightarrow \infty} \tau_n = \infty \text{ and } \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} |U(\tau_n, x) - f_0(z_n) \bar{V}(x)| dx = 0,$$

where $z_n = e^{\tau_n} x + c \frac{e^{2\tau_n} - 1}{2}$. According to a result of G. Allaire (see [2]),

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} f_0(z_n) \bar{V}(x) dx = \langle f_0 \rangle \int_{\mathbb{R}^N} \bar{V} = \int_{\mathbb{R}^N} \bar{V};$$

gathering the three equalities, we obtain property (i).

We now address the proof of property (ii), which relies on the second point in Lemma 2.5; let $\bar{V} \in \Omega$ be arbitrary, and for all $\varepsilon > 0$, let $\bar{V}_\varepsilon \in L^2(K) \cap L^\infty(\mathbb{R}^N)$ such that

$$\|\bar{V}_\varepsilon - \bar{V}\|_{L^1(\mathbb{R}^N)} \leq \varepsilon.$$

Let $(\tau_n)_{n \geq 0}$ be a sequence of positive numbers such that $\tau_n \rightarrow \infty$ and

$$\int_{\mathbb{R}^N} |U(\tau_n, x) - f_0(z_n) \bar{V}(x)| dx \rightarrow 0,$$

where $z_n = e^{\tau_n} x + c \frac{e^{2\tau_n} - 1}{2}$.

Let $\rho \in \mathcal{C}_0^\infty(\mathbb{R}^N)$ be a mollifying kernel; for $\delta > 0$, set $\rho_\delta := \delta^{-N} \rho(\cdot/\delta)$, and define the function $U_n^{\delta, \varepsilon}$ by

$$U_n^{\delta, \varepsilon}(\tau, x) := U^{\text{app}} [(S_\tau \bar{V}_\varepsilon) *_x \rho_\delta](\tau, x; e^{\tau_n + \tau}).$$

Then Lemma 2.5 ensures that $U_n^{\delta, \varepsilon}$ satisfies equation (2.3) with an error term, the latter being bounded for all $T > 0$ in $L^\infty([0, T], L^1(\mathbb{R}^N))$ by

$$\omega_{T, \varepsilon}(\delta) + C_{T, \varepsilon, \delta} e^{-\tau_n}$$

where $\omega_{T, \varepsilon} : [0, \infty) \rightarrow [0, \infty)$ is such that $\lim_{0+} \omega_{T, \varepsilon} = 0$, and where the constant $C_{T, \varepsilon, \delta}$ depends only on ε, δ, N and T .

Using the L^1 contraction principle for scalar conservation laws, we infer that for all $T > 0$, and for all $\tau \in [0, T]$,

$$\begin{aligned} & \int_{\mathbb{R}^N} |U(\tau_n + \tau, x) - U_n^{\delta, \varepsilon}(\tau, x)| dx \\ & \leq \omega_{T, \varepsilon}(\delta) + C_{T, \varepsilon, \delta} e^{-\tau_n} + \int_{\mathbb{R}^N} |U(\tau_n, x) - U_{n|\tau=0}^{\delta, \varepsilon}(x)| dx \\ & \leq \omega_{T, \varepsilon}(\delta) + C_{T, \varepsilon, \delta} e^{-\tau_n} + \int_{\mathbb{R}^N} |U(\tau_n, x) - f_0(z_n) \bar{V}(x)| dx \\ & \quad + \int_{\mathbb{R}^N} |U_{n|\tau=0}^{\delta, \varepsilon}(x) - f_0(z_n) \bar{V}(x)| dx. \end{aligned}$$

Now, according to Definition 2.1,

$$\begin{aligned} U_n^{\delta, \varepsilon}(\tau, x) &= (S_\tau \bar{V}_\varepsilon) *_x \rho_\delta(x) f_0 \left(e^{\tau_n + \tau} x + c \frac{e^{2(\tau_n + \tau)} - 1}{2} \right) \\ &+ e^{-(\tau_n + \tau)} U_1 \left(\tau, x, e^{\tau_n + \tau} x + c \frac{e^{2(\tau_n + \tau)} - 1}{2} \right) \\ &+ e^{-2(\tau_n + \tau)} V_2 \left(\tau, x, e^{\tau_n + \tau} x + c \frac{e^{2(\tau_n + \tau)} - 1}{2} \right). \end{aligned}$$

Hence for all $\tau \in [0, T]$, we have

$$\begin{aligned}
& \int_{\mathbb{R}^N} \left| U_n^{\delta, \varepsilon}(\tau, x) - S_\tau \bar{V} f_0 \left(e^{\tau_n + \tau} x + c \frac{e^{2(\tau_n + \tau)} - 1}{2} \right) \right| dx \\
& \leq \|f_0\|_\infty \sup_{\tau \in [0, T]} \|S_\tau \bar{V} - (S_\tau \bar{V}_\varepsilon) *_x \rho_\delta\|_{L^1(\mathbb{R}^N)} \\
& \quad + e^{-\tau_n} \left(\|U_1\|_{L^\infty([0, T] \times \mathbb{T}_z^N, L^1(\mathbb{R}_x^N))} + \|U_2\|_{L^\infty([0, T] \times \mathbb{T}_z^N, L^1(\mathbb{R}_x^N))} \right) \\
& \leq \|f_0\|_\infty \left[\|\bar{V} - \bar{V}_\varepsilon\|_{L^1} + \sup_{\tau \in [0, T]} \|S_\tau \bar{V}_\varepsilon - (S_\tau \bar{V}_\varepsilon) *_x \rho_\delta\|_{L^1(\mathbb{R}^N)} \right] \\
& \quad + e^{-\tau_n} \left(\|U_1\|_{L^\infty([0, T] \times \mathbb{T}_z^N, L^1(\mathbb{R}_x^N))} + \|U_2\|_{L^\infty([0, T] \times \mathbb{T}_z^N, L^1(\mathbb{R}_x^N))} \right) \\
& \leq C\varepsilon + \omega_{T, \varepsilon}(\delta) + C_{T, \varepsilon, \delta} e^{-\tau_n}.
\end{aligned}$$

Gathering the two inequalities, we deduce that for all n, δ, ε ,

$$\begin{aligned}
& \sup_{\tau \in [0, T]} \int_{\mathbb{R}^N} \left| U(\tau_n + \tau, x) - S_\tau \bar{V}(x) f_0 \left(e^{\tau_n + \tau} x + c \frac{e^{2(\tau_n + \tau)} - 1}{2} \right) \right| dx \\
& \leq \omega_{T, \varepsilon}(\delta) + C_{T, \varepsilon, \delta} e^{-\tau_n} + \int_{\mathbb{R}^N} |U(\tau_n, x) - f_0(z_n) \bar{V}(x)| dx + C\varepsilon.
\end{aligned}$$

In the right-hand side of the above inequality, we first choose ε sufficiently small, then δ so that $\omega_{T, \varepsilon}(\delta)$ is sufficiently small, and eventually n large enough so that the two remaining terms are small as well; hence

$$\lim_{n \rightarrow \infty} \inf_{\varepsilon > 0, \delta > 0} \left(\omega_{T, \varepsilon}(\delta) + C_{T, \varepsilon, \delta} e^{-\tau_n} + \int_{\mathbb{R}^N} |U(\tau_n, x) - f_0(z_n) \bar{V}(x)| dx + C\varepsilon \right) = 0.$$

Thus we have proved that for all $T > 0$,

$$\lim_{n \rightarrow \infty} \sup_{\tau \in [0, T]} \int_{\mathbb{R}^N} \left| U(\tau_n + \tau, x) - (S_\tau \bar{V}) f_0 \left(e^{\tau_n + \tau} x + c \frac{e^{2(\tau_n + \tau)} - 1}{2} \right) \right| dx = 0.$$

The above convergence entails immediately that $S_\tau \bar{V} \in \Omega$ for all $\tau \in [0, T]$. Since $T > 0$ was arbitrary, property (ii) is proved.

There remains to prove property (iii), which is a variant of the LaSalle invariance principle; let $\bar{V} \in \Omega$ be arbitrary, and let τ_n be a sequence of positive numbers such that $\lim_{n \rightarrow \infty} \tau_n = +\infty$ and

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} |U(\tau_n, x) - f_0(z_n) \bar{V}(x)| dx = 0,$$

where $z_n = e^{\tau_n} x + c \frac{e^{2\tau_n} - 1}{2}$. According to a result of G. Allaire (see [2]), we have, since

$\langle f_0 \rangle = 1$ and $f_0 \in \mathcal{C}(\mathbb{T}^N)$,

$$\begin{aligned}
\|\bar{V} - F_M\|_{L^1(\mathbb{R}^N)} &= \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} f_0(z_n) |\bar{V}(x) - F_M(x)| dx \\
&= \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} f_0(z_n) \left| \bar{V}(x) - \frac{U(\tau_n, x)}{f_0(z_n)} + \frac{U(\tau_n, x)}{f_0(z_n)} - F_M(x) \right| dx \\
&= \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} f_0(z_n) \left| \frac{U(\tau_n, x)}{f_0(z_n)} - F_M(x) \right| dx \\
&= \ell(U_{ini}).
\end{aligned}$$

Consequently,

$$\|\bar{V} - F_M\|_{L^1(\mathbb{R}^N)} = \ell(U_{ini}) \quad \forall \bar{V} \in \Omega.$$

□

Corollary 4.1. *Let $U_{ini} \in L^1(\mathbb{R}^N)$, and set*

$$M := \int_{\mathbb{R}^N} U_{ini}.$$

Assume that the ω -limit set $\Omega[U_{ini}]$ is non-empty. Then $\ell(U_{ini}) = 0$, and thus the result of Proposition 3.1 holds.

Proof. Let $\bar{V} \in \Omega$ be arbitrary. Then

$$\lim_{\tau \rightarrow \infty} \|S_\tau \bar{V} - F_M\|_{L^1(\mathbb{R}^N)} = 0;$$

this property is stated in Proposition 2.1 in the case when $\bar{V} \in L^\infty(\mathbb{R}^N) \cap L^2(K)$, but can be in fact easily generalized to an arbitrary function $\bar{V} \in L^1$ by using the contractivity of the semi-group S_τ : indeed, let $\varepsilon > 0$, and let $\bar{V}_\varepsilon \in L^\infty(\mathbb{R}^N) \cap L^2(K)$ such that $\int \bar{V}_\varepsilon = \int \bar{V} = M$, and

$$\|\bar{V}_\varepsilon - \bar{V}\|_{L^1(\mathbb{R}^N)} \leq \varepsilon.$$

Then for all $\tau \geq 0$,

$$\begin{aligned}
\|S_\tau \bar{V} - F_M\|_{L^1(\mathbb{R}^N)} &\leq \|S_\tau \bar{V} - S_\tau \bar{V}_\varepsilon\|_{L^1(\mathbb{R}^N)} + \|S_\tau \bar{V}_\varepsilon - F_M\|_{L^1(\mathbb{R}^N)} \\
&\leq \|\bar{V} - \bar{V}_\varepsilon\|_{L^1(\mathbb{R}^N)} + \|S_\tau \bar{V}_\varepsilon - F_M\|_{L^1(\mathbb{R}^N)}.
\end{aligned}$$

Hence, using Proposition 2.1, we infer that

$$\limsup_{\tau \rightarrow \infty} \|S_\tau \bar{V} - F_M\|_{L^1(\mathbb{R}^N)} \leq \varepsilon \quad \forall \varepsilon > 0,$$

and thus $\|S_\tau \bar{V} - F_M\|_{L^1(\mathbb{R}^N)}$ vanishes as $\tau \rightarrow \infty$.

On the other hand, property (ii) in Proposition 4.1 ensures that $S_\tau \bar{V} \in \Omega$ for all $\tau \geq 0$, and thus, using (iii),

$$\|S_\tau \bar{V} - F_M\|_{L^1(\mathbb{R}^N)} = \ell(U_{ini}) \quad \forall \tau \geq 0.$$

Consequently, $\ell(U_{ini}) = 0$. Going back to the definition of $\ell(U_{ini})$, we deduce that

$$\lim_{\tau \rightarrow \infty} \int_{\mathbb{R}^N} \left| U(\tau, x) - f_0 \left(e^\tau x + c \frac{e^{2\tau} - 1}{2} \right) F_M(x) \right| dx = 0.$$

□

Thus the proof of Proposition 3.1 is complete provided we are able to show that the set $\Omega[U_{ini}]$ is non-empty for a sufficiently large class of functions $U_{ini} \in L^1(\mathbb{R}^N)$. In the case when $\|U_{ini}\|_1$ is small, this result follows from Proposition 3.2 and from a contraction principle. The proof in the general case is more involved, and in fact, an analysis similar to the one performed in Section 3 has to be conducted once more.

4.3 Proof of Proposition 3.1 when $\|U_{ini}\|_{L^1(\mathbb{R}^N)}$ is small

We now complete the proof of Theorem 1.1 when $\|U_{ini}\|_{L^1(\mathbb{R}^N)}$ is small. Let $U_{ini} \in L^1(\mathbb{R}^N)$. Assume that U_{ini} satisfies the following assumptions

$$\exists m > 2(N + 1), \quad U_{ini} \in L^2((1 + |x|^2)^{m/2}) \cap L^\infty(\mathbb{R}^N), \quad (4.3)$$

$$\text{and} \quad \|U_{ini}\|_{L^1(\mathbb{R}^N)} \leq C_m, \quad (4.4)$$

where the constant C_m was introduced in Proposition 3.2. Then according to Proposition 3.2, the ω -limit set $\Omega[U_{ini}]$ is non-empty, and consequently Proposition 3.1 is true (see Corollary 4.1).

Let us now prove that Proposition 3.1 holds when U_{ini} merely satisfies (4.4): this fact is a direct consequence of the density of $L^2((1 + |x|^2)^{m/2}) \cap L^\infty(\mathbb{R}^N)$ in $L^1(\mathbb{R}^N)$, together with the continuity of ℓ . Indeed, for all $\varepsilon > 0$, let $U_{ini}^\varepsilon \in L^2((1 + |x|^2)^{m/2}) \cap L^\infty(\mathbb{R}^N)$ such that

$$\|U_{ini} - U_{ini}^\varepsilon\|_{L^1(\mathbb{R}^N)} \leq \varepsilon, \quad \|U_{ini}^\varepsilon\|_{L^1(\mathbb{R}^N)} \leq C_m.$$

Then $\ell(U_{ini}^\varepsilon) = 0$. Since ℓ is Lipschitz continuous (see Lemma 4.3), there exists a constant C such that

$$\ell(U_{ini}) = |\ell(U_{ini}) - \ell(U_{ini}^\varepsilon)| \leq C \|U_{ini} - U_{ini}^\varepsilon\|_{L^1(\mathbb{R}^N)} \leq C\varepsilon.$$

Since the above inequality holds for all $\varepsilon > 0$, we deduce that $\ell(U_{ini}) = 0$. Recalling the definition of ℓ , we infer that Proposition 3.1 holds for all initial data $U_{ini} \in L^1(\mathbb{R}^N)$ satisfying (4.4).

4.4 Proof of Proposition 3.1 in the general case

The case when $\|U_{ini}\|_1$ is large follows from the following Lemma:

Lemma 4.4. *There exists a constant C_0 , depending only on N and on the flux A , such that for all $U_{ini} \in L^1(\mathbb{R}^N)$,*

$$\ell(U_{ini}) \leq C_0 \Rightarrow \ell(U_{ini}) = 0.$$

Before proving the above Lemma, let us explain why the result of Proposition 3.1 follows. Since ℓ is Lipschitz continuous, $\ell(L^1(\mathbb{R}^N))$ is a connected set of $[0, \infty)$, i.e. an interval. Moreover, $0 \in \ell(L^1(\mathbb{R}^N))$ according to the previous paragraph, and Lemma 4.4 entails that

$$\ell(L^1(\mathbb{R}^N)) \cap (0, C_0] = \emptyset.$$

Consequently, $\ell(L^1(\mathbb{R}^N)) = \{0\}$.

There remains to prove Lemma 4.4. According to Corollary 4.1 and using by now standard arguments, we only have to prove that there exists a set $\mathcal{A} \subset L^1(\mathbb{R}^N)$, which is dense in $L^1(\mathbb{R}^N)$, and such that

$$\exists C > 0, \forall U_{ini} \in L^1(\mathbb{R}^N) \cap \mathcal{A}, \quad \ell(U_{ini}) \leq C \Rightarrow \Omega[U_{ini}] \neq \emptyset. \quad (4.5)$$

In the following, we will take $\mathcal{A} = L^2((1 + |x|^2)^{m/2})$, for some $m > 0$ sufficiently large.

The scheme of proof of the implication (4.5) is very similar to the one of Proposition 3.2; indeed, we have to prove that if $\ell(U_{ini})$ is small enough, then there exists a sequence (τ_n) of positive numbers, with $\lim_{n \rightarrow \infty} \tau_n = +\infty$, such that $(V(\tau_n, \cdot))_{n \rightarrow \infty}$ is a compact sequence in $L^1(\mathbb{R}^N)$. Notice that this is obviously equivalent to the compactness of the sequence $V(\tau_n, \cdot) - F_M$, whose L^1 norm is of the order of $\ell(U_{ini})$ as $n \rightarrow \infty$. Thus our strategy is the following: rather than using directly the equation on U , we consider the equation on the function $U - U^{\text{app}}[F_M]$. We prove that for an appropriate function \tilde{U} , an inequality of the type (3.6) holds, with U replaced by $U - U^{\text{app}}[F_M]$. Then, all the occurrences of $\|U(\tau)\|_{L^1(\mathbb{R}^N)}$ in the proof of Proposition 3.2 are replaced by $\|(U - U^{\text{app}}[F_M])(\tau)\|_{L^1(\mathbb{R}^N)}$, which converges towards $\ell(U_{ini})$ as $\tau \rightarrow \infty$. Thus the same arguments which led us to compactness in the case when $\|U_{ini}\|_{L^1(\mathbb{R}^N)}$ is small show that compactness holds, provided $\ell(U_{ini})$ is small enough.

Let us now retrace the main lines of the proof: first, consider a function $U_{ini} \in L^1(\mathbb{R}^N)$ such that $U_{ini} \in L^2((1 + |x|^2)^{m/2})$ for some sufficiently large m (to be chosen later). Set $M = \int_{\mathbb{R}^N} U_{ini}$ and

$$W(\tau, x) = U(\tau, x) - U^{\text{app}}[F_M](\tau, x; e^\tau).$$

In the rest of the proof, for the sake of brevity, we will write $U^{\text{app}}(\tau, x)$ as a short-hand for $U^{\text{app}}[F_M](\tau, x; e^\tau)$. Then the following properties hold

$$\begin{aligned} W &\in L_{\text{loc}}^\infty([0, \infty), L^2((1 + |x|^2)^{m/2})) \cap L_{\text{loc}}^2([0, \infty), H^1((1 + |x|^2)^{m/2})), \\ &\exists C > 0, \forall \tau \geq 0, \quad \|W(\tau, \cdot)\|_{L^\infty(\mathbb{R}^N)} \leq C e^{N\tau}, \\ &\lim_{\tau \rightarrow \infty} \|W(\tau)\|_{L^1(\mathbb{R}^N)} = \ell(U_{ini}). \end{aligned}$$

Moreover, using Lemma 2.5, we deduce that W satisfies

$$\begin{aligned} \partial_\tau W &= \operatorname{div}_x(xW) + \Delta_x W - R \operatorname{div}_x((\alpha_1(z) - c)W) \\ &\quad - R^{N+1} \operatorname{div}_x \left[\tilde{B}_1 \left(z, \frac{U}{R^N} \right) - \tilde{B}_1 \left(z, \frac{U^{\text{app}}}{R^N} \right) \right] \\ &\quad + U^{\text{rem}}, \end{aligned}$$

with $R = e^\tau$, $z = Rx + c \frac{R^2 - 1}{2}$, and we recall that the remainder U^{rem} satisfies

$$\|U^{\text{rem}}(\tau)\|_{L^\infty(\mathbb{R}^N)} + \|U^{\text{rem}}(\tau)\|_{L^2(e^{\gamma|x|^2})} \leq C e^{-\tau} \quad (4.6)$$

for some $\gamma > 0$.

Then, using the bounds on U, U^{app} together with the regularity assumptions on \tilde{B} , it can be easily proved that

$$\tilde{B}_1 \left(z, \frac{U(\tau, x)}{R^N} \right) - \tilde{B}_1 \left(z, \frac{U^{\text{app}}(\tau, x)}{R^N} \right) = 2\alpha_2(z) f_0(z) \frac{F_M(x) W(\tau, x)}{R^{2N}} + b(\tau, x),$$

and the function b is such that there exists $C > 0$ such that

$$\forall (\tau, x) \in \mathbb{R}_+ \times \mathbb{R}^N, \quad |b(\tau, x)| \leq C \left(\left| \frac{W(\tau, x)}{R^N} \right|^2 + \frac{|W(\tau, x)|}{R^{2N+1}} + \frac{|U^{\text{app}}(\tau, x)|^3}{R^{3N}} \right).$$

We define a function \tilde{W} by

$$\tilde{W}(\tau, x) = W_0(x, z) + e^{-\tau} W_1(x, z),$$

with $W_0(x, z) = f_0(z) h_m(x)$ and

$$-\Delta_z W_1 + \operatorname{div}_z(\alpha_1 W_1) = 2\Delta_{yz} W_0 - \operatorname{div}_x((\alpha_1 - c)W_0) - 2\mathbf{1}_{N=1} \operatorname{div}_z(\alpha_2 f_0 F_M W_0).$$

Notice that by definition of f_0 and c , the compatibility condition is always satisfied, and

$$W_1(x, z) = f_1(z) \cdot \nabla_y h_m(x) + \mathbf{1}_{N=1} w_1(z) F_M(x) h_m(x),$$

with

$$-\Delta_z w_1 + \operatorname{div}_z(\alpha_1 w_1) = -2 \operatorname{div}_z(\alpha_2 f_0^2).$$

Let $\tau_0 > 0$ such that

$$\tilde{W}(\tau, x) \geq \frac{1}{2} f_0(z) h_m(x) \quad \forall \tau \geq \tau_0, \quad \forall y \in \mathbb{R}^N.$$

For further purposes, we also choose τ_0 such that

$$\|W(\tau, \cdot)\|_{L^1(\mathbb{R}^N)} \leq 2\ell(U_{\text{ini}}) \quad \forall \tau \geq \tau_0.$$

(Notice that if $\ell(U_{ini}) = 0$ there is nothing to prove).

Using calculations similar to the ones performed in the proof of Proposition 3.2, we infer that for $\tau \geq \tau_0$,

$$\begin{aligned}
& \frac{d}{d\tau} \int_{\mathbb{R}^N} \left| \frac{W}{\tilde{W}} \right|^2 \tilde{W}(\tau) + \frac{m-N}{4} \int_{\mathbb{R}^N} \left| \frac{W}{\tilde{W}} \right|^2 \tilde{W}(\tau) + 2 \int_{\mathbb{R}^N} \left| \nabla \left(\frac{W}{\tilde{W}} \right) \right|^2 \tilde{W}(\tau) \\
\leq & C \int_{\mathbb{R}^N} \left(\frac{W(\tau, x)}{\tilde{W}(\tau, x)} \right)^2 \frac{dx}{(1+|x|^2)^{1+\frac{m}{2}}} \\
& + C e^{(1-N)\tau} \int_{\mathbb{R}^N} |W(\tau, x)|^2 \left| \nabla \left(\frac{W(\tau, x)}{\tilde{W}(\tau, x)} \right) \right| dx \\
& + C e^{-N\tau} \int_{\mathbb{R}^N} |W(\tau, x)| \left| \nabla \left(\frac{W(\tau, x)}{\tilde{W}(\tau, x)} \right) \right| dx \\
& + C e^{(-2N+1)\tau} \int_{\mathbb{R}^N} |U^{\text{app}}(\tau, x)|^3 \left| \nabla \left(\frac{W(\tau, x)}{\tilde{W}(\tau, x)} \right) \right| dx \\
& + \int_{\mathbb{R}^N} \left| \frac{W(\tau, x)}{\tilde{W}(\tau, x)} \right| |U^{\text{rem}}(\tau, x)| dx.
\end{aligned}$$

Using the same arguments as in the third step of the proof of Proposition 3.2, we deduce that if $m > 2(N+1)$,

$$\begin{aligned}
C \int_{\mathbb{R}^N} \left(\frac{W(\tau, x)}{\tilde{W}(\tau, x)} \right)^2 \frac{dx}{(1+|x|^2)^{1+\frac{m}{2}}} & \leq \frac{m-N}{20} \int_{\mathbb{R}^N} \left| \frac{W(\tau)}{\tilde{W}(\tau)} \right|^2 \tilde{W}(\tau) \\
& + \frac{1}{3} \int_{\mathbb{R}^N} \left| \nabla \left(\frac{W(\tau)}{\tilde{W}(\tau)} \right) \right|^2 \tilde{W}(\tau) \\
& + C \ell(U_{ini})^2.
\end{aligned}$$

Similarly, the calculations of the fourth step in the proof of Proposition 3.2 yield

$$\begin{aligned}
& e^{(1-N)\tau} \int_{\mathbb{R}^N} |W(\tau, x)|^2 \left| \nabla_x \left(\frac{W(\tau, x)}{\tilde{W}(\tau, x)} \right) \right| dx \\
\leq & C \|W(\tau)\|_{L^1(\mathbb{R}^N)}^{1/N} \left[\int_{\mathbb{R}^N} \left| \frac{W(\tau)}{\tilde{W}(\tau)} \right|^2 \tilde{W}(\tau) + \int_{\mathbb{R}^N} \left| \nabla \left(\frac{W(\tau)}{\tilde{W}(\tau)} \right) \right|^2 \tilde{W}(\tau) \right] \\
\leq & C \ell(U_{ini})^{1/N} \left[\int_{\mathbb{R}^N} \left| \frac{W(\tau)}{\tilde{W}(\tau)} \right|^2 \tilde{W}(\tau) + \int_{\mathbb{R}^N} \left| \nabla \left(\frac{W(\tau)}{\tilde{W}(\tau)} \right) \right|^2 \tilde{W}(\tau) \right].
\end{aligned}$$

The two additional terms coming from the estimation of b can be easily bounded thanks to the Cauchy-Schwarz inequality. For $\tau \geq \tau_0$, and τ_0 large enough, we have

$$\begin{aligned}
& C e^{-N\tau} \int_{\mathbb{R}^N} |W(\tau)| \left| \nabla_y \left(\frac{W(\tau)}{\tilde{W}(\tau)} \right) \right| + C e^{(1-2N)\tau} \int_{\mathbb{R}^N} |U^{\text{app}}(\tau)|^3 \left| \nabla_y \left(\frac{W(\tau)}{\tilde{W}(\tau)} \right) \right| dx \\
\leq & \frac{m-N}{20} \int_{\mathbb{R}^N} \left| \frac{W(\tau)}{\tilde{W}(\tau)} \right|^2 \tilde{W}(\tau) + \frac{1}{3} \int_{\mathbb{R}^N} \left| \nabla \left(\frac{W(\tau)}{\tilde{W}(\tau)} \right) \right|^2 \tilde{W}(\tau) + C.
\end{aligned}$$

Eventually, using the Cauchy-Schwarz inequality together with the bound (4.6), we infer that

$$\begin{aligned}
& \int_{\mathbb{R}^N} \left| \frac{W(\tau, x)}{\tilde{W}(\tau, x)} \right| |U^{rem}(\tau, x)| dx \\
& \leq \|U^{rem}(\tau)\|_{L^2(\tilde{W}(\tau)^{-1})} \left(\int_{\mathbb{R}^N} \left| \frac{W(\tau)}{\tilde{W}(\tau)} \right|^2 \tilde{W}(\tau) \right)^{1/2} \\
& \leq C \|U^{rem}(\tau)\|_{L^2(e^{\gamma|x|^2})} \left(\int_{\mathbb{R}^N} \left| \frac{W(\tau)}{\tilde{W}(\tau)} \right|^2 \tilde{W}(\tau) \right)^{1/2} \\
& \leq C e^{-\tau} \left(\int_{\mathbb{R}^N} \left| \frac{W(\tau)}{\tilde{W}(\tau)} \right|^2 \tilde{W}(\tau) \right)^{1/2} \\
& \leq C + \frac{m-N}{20} \int_{\mathbb{R}^N} \left| \frac{W(\tau)}{\tilde{W}(\tau)} \right|^2 \tilde{W}(\tau).
\end{aligned}$$

Gathering all the terms, we deduce that there exists a constant C_m , depending only on N and m , such that if $\ell(U_{ini}) \leq C_m$, then for all $\tau \geq \tau_0$,

$$\frac{d}{d\tau} \int_{\mathbb{R}^N} \left| \frac{W(\tau)}{\tilde{W}(\tau)} \right|^2 \tilde{W}(\tau) + \frac{m-N}{20} \int_{\mathbb{R}^N} \left| \frac{W(\tau)}{\tilde{W}(\tau)} \right|^2 \tilde{W}(\tau) + \int_{\mathbb{R}^N} \left| \nabla \left(\frac{W(\tau)}{\tilde{W}(\tau)} \right) \right|^2 \tilde{W}(\tau) \leq C.$$

Compactness of a subsequence $W(\tau_n)$ follows. Hence the ω -limit set is non-empty, and thus $\ell(U_{ini}) = 0$.

Appendix A

Lemma A.1. *Assume that the flux A satisfies (1.4), (1.5). Let $v \in W^{1,\infty}(\mathbb{T}^N)$ be a periodic stationary solution of (1.1), and let $u \in L_{loc}^\infty([0, \infty), L^\infty(\mathbb{R}^N)) \cap \mathcal{C}([0, \infty), L_{loc}^1(\mathbb{R}^N))$ be the unique solution of (1.1) with initial data $u_{ini} \in v(y) + L^1 \cap L^\infty(\mathbb{R}^N)$. Then $u \in L^\infty([0, \infty) \times \mathbb{R}^N)$.*

Proof. This result was proved in [8] in the case $N = 1$. When $N \geq 2$, the proof goes along the same lines; the only difference lies in the use of the Sobolev embeddings, which depend on the dimension. Hence we merely recall here the main steps of the proof, with an emphasis on the case $N \geq 2$.

In the rest of the proof, we set $f(t, y) = u(t, y) - v(y)$. Then f solves the equation

$$\partial_t f + \operatorname{div}_y B(y, f) - \Delta_y f = 0, \tag{4.7}$$

and according to (1.5) the flux B is such that for all $f \in \mathbb{R}$,

$$\begin{aligned}
|\operatorname{div}_y B(y, f)| & \leq C(|f| + |f|^n), \\
|\partial_f B(y, f)| & \leq C(|f| + |f|^n),
\end{aligned}$$

where the exponent n is such that $n < (N + 2)/N$. Moreover,

$$\|f(t)\|_{L^1(\mathbb{R}^N)} \leq \|u_{ini} - v\|_{L^1(\mathbb{R}^N)} \quad \forall t \geq 0.$$

For $q \geq 1$ arbitrary, multiply (4.7) by $f|f|^{q-1}$, and integrate over \mathbb{R}^N . Using a few integrations by parts (see [8]), we are led to

$$\frac{d}{dt} \int_{\mathbb{R}^N} |f|^{q+1} + c_q \int_{\mathbb{R}^N} \left| \nabla_y |f|^{\frac{q+1}{2}} \right|^2 \leq C_q \left(\int_{\mathbb{R}^N} |f|^{q+1} + \int_{\mathbb{R}^N} |f|^{q+n} \right) \quad (4.8)$$

We then use Sobolev embeddings in order to control the L^{q+1} and L^{q+n} norms in the right-hand side. We distinguish between the cases $N = 2$ and $N \geq 3$, since the space H^1 is critical in dimension two.

• If $N = 2$, then $H^1(\mathbb{R}^2) \subset L^p(\mathbb{R}^2)$ for all $p \in [2, \infty)$. Interpolating L^{q+n} between L^1 and L^p for some p sufficiently large, we have, using the Gagliardo-Nirenberg-Sobolev inequality

$$\begin{aligned} \|f\|_{L^{q+n}(\mathbb{R}^2)} &\leq \|f\|_{L^1(\mathbb{R}^2)}^\theta \|f\|_{L^p(\mathbb{R}^2)}^{1-\theta} \quad \text{with} \quad \frac{1}{q+n} = \frac{\theta}{1} + \frac{1-\theta}{p} \\ &\leq \|f\|_{L^1(\mathbb{R}^2)}^\theta \left\| |f|^{\frac{q+1}{2}} \right\|_{L^{\frac{2p}{q+1}}(\mathbb{R}^2)}^{\frac{2(1-\theta)}{q+1}} \\ &\leq C_p \|f\|_{L^1(\mathbb{R}^2)}^\theta \|f\|_{L^{\frac{q+1}{2}}(\mathbb{R}^2)}^{\frac{(1-\theta)(q+1)}{2p}} \|\nabla |f|^{\frac{q+1}{2}}\|_{L^2(\mathbb{R}^2)}^{(1-\theta)\left(\frac{2}{q+1} - \frac{1}{p}\right)} \end{aligned}$$

Notice that

$$2\frac{q+n}{q+1}(1-\theta) = 2\frac{q+n-1}{(q+1)\left(1-\frac{1}{p}\right)},$$

and

$$\frac{q+n-1}{q+1} < 1 \quad \forall q \geq 1$$

since $n < (N + 2)/N$. Thus, we choose $p > 1$ such that

$$(q+n)(1-\theta) \left(\frac{2}{q+1} - \frac{1}{p} \right) < 2.$$

Young's inequality then implies that for all $\lambda > 0$, there exists a constant $C_{\lambda,q}$ and exponents q_1, q_2 such that

$$\int_{\mathbb{R}^N} |f|^{q+n} \leq \lambda \left\| \nabla |f|^{\frac{q+1}{2}} \right\|_{L^2(\mathbb{R}^2)}^2 + C_{\lambda,q} \left(\|f\|_{L^1(\mathbb{R}^2)}^{q_1} + \|f\|_{L^{\frac{q+1}{2}}(\mathbb{R}^2)}^{q_2} \right). \quad (4.9)$$

The other term in the right-hand side of (4.8) can be bounded in a similar fashion: we have, for all $\lambda > 0$,

$$\int_{\mathbb{R}^N} |f|^{q+1} \leq \lambda \left\| \nabla |f|^{\frac{q+1}{2}} \right\|_{L^2(\mathbb{R}^2)}^2 + C_{\lambda,q} \left(\|f\|_{L^1(\mathbb{R}^2)}^{q_3} + \|f\|_{L^{\frac{q+1}{2}}(\mathbb{R}^2)}^{q_4} \right), \quad (4.10)$$

for some exponents q_3, q_4 which can be explicitly computed. Choosing an appropriate parameter λ , we infer that there exist $q_1, q_2, q_3, q_4 > 0$ such that

$$\begin{aligned} & \frac{d}{dt} \int_{\mathbb{R}^N} |f|^{q+1} + c_q \int_{\mathbb{R}^N} \left| \nabla_y |f|^{\frac{q+1}{2}} \right|^2 \\ & \leq C_q \left(\|f\|_{L^1(\mathbb{R}^2)}^{q_1} + \|f\|_{L^{\frac{q+1}{2}}(\mathbb{R}^2)}^{q_2} + \|f\|_{L^1(\mathbb{R}^2)}^{q_3} + \|f\|_{L^{\frac{q+1}{2}}(\mathbb{R}^2)}^{q_4} \right). \end{aligned}$$

Using (4.10) one more time leads to

$$\begin{aligned} & \frac{d}{dt} \int_{\mathbb{R}^N} |f|^{q+1} + c_q \int_{\mathbb{R}^N} |f|^{q+1} \\ & \leq C_q \left(\|f\|_{L^1(\mathbb{R}^2)}^{q_1} + \|f\|_{L^{\frac{q+1}{2}}(\mathbb{R}^2)}^{q_2} + \|f\|_{L^1(\mathbb{R}^2)}^{q_3} + \|f\|_{L^{\frac{q+1}{2}}(\mathbb{R}^2)}^{q_4} \right). \end{aligned}$$

Using a Gronwall-type argument, we infer that for all $q \geq 1$,

$$f \in L^\infty([0, \infty), L^{\frac{q+1}{2}}(\mathbb{R}^2)) \Rightarrow f \in L^\infty([0, \infty), L^q(\mathbb{R}^2)).$$

Since $f \in L^\infty([0, \infty), L^1(\mathbb{R}^2))$, we infer by induction on q that $f \in L^\infty([0, \infty), L^q(\mathbb{R}^2))$ for all $q \geq 1$.

- When $N \geq 3$, we use the Sobolev embedding $H^1(\mathbb{R}^N) \subset L^{p^*}(\mathbb{R}^N)$, where

$$p^* = \frac{2N}{N-2}.$$

Interpolating L^{q+n} between L^1 and $L^{\frac{p^*(q+1)}{2}}$, we obtain

$$\begin{aligned} \|f\|_{L^{q+n}(\mathbb{R}^N)} & \leq \|f\|_{L^{\frac{p^*(q+1)}{2}}(\mathbb{R}^N)}^\theta \|f\|_{L^1(\mathbb{R}^N)}^{1-\theta} \leq \left\| |f|^{\frac{q+1}{2}} \right\|_{L^{p^*}(\mathbb{R}^N)}^{\frac{2\theta}{q+1}} \|f\|_{L^1(\mathbb{R}^N)}^{1-\theta} \\ & \leq C \left\| \nabla |f|^{\frac{q+1}{2}} \right\|_{L^2(\mathbb{R}^2)}^{\frac{2\theta}{q+1}} \|f\|_{L^1(\mathbb{R}^N)}^{1-\theta}, \end{aligned}$$

where the parameter $\theta \in (0, 1)$ is given by

$$\frac{1}{q+n} = \frac{2\theta}{p^*(q+1)} + \frac{1-\theta}{1}.$$

It can be checked that

$$n < \frac{N+2}{N} \Rightarrow \frac{\theta(q+n)}{q+1} < 1.$$

Hence (4.9) holds when $N \geq 3$. Inequality (4.10) is proved with similar arguments. As in the two-dimensional case, we deduce that $f \in L^\infty([0, \infty), L^q(\mathbb{R}^N))$ for all q . Using Theorem 8.1 in Chapter III of [16] (see [8] for details), we infer eventually that $f \in L^\infty([0, \infty) \times \mathbb{R}^N)$.

□

Appendix B

Lemma A.2. *Let $M > M'$ be arbitrary. Then*

$$F_M(y) > F_{M'}(y) \quad \forall y \in \mathbb{R}^N.$$

As a consequence,

$$\|F_M - F_{M'}\|_{L^1(\mathbb{R}^N)} = M - M'.$$

Proof. The arguments are exactly the ones which lead to the uniqueness of stationary solutions of (2.13), (2.15), and they can be found in [1]. We recall the main steps below for the reader's convenience.

Let $F := F_M - F_{M'}$. Then $F \in L^1 \cap C^2(\mathbb{R}^N)$, and $\int_{\mathbb{R}^N} F > 0$. Hence the set

$$\Theta := \{x \in \mathbb{R}^N, F(x) > 0\}$$

is non-empty. The idea is to prove that $F_+ = F \mathbf{1}_\Theta$ satisfies a linear elliptic equation; since $F_+ \geq 0$, the strong maximum principle entails that F_+ cannot vanish anywhere, and thus $F_+(x) > 0$ for all $x \in \mathbb{R}^N$.

Let us now derive an equation on F_+ . Subtracting the equations on F_M and $F_{M'}$, we have

$$-\sum_{1 \leq i, j \leq N} \eta_{i,j} \frac{\partial^2 F}{\partial x_i \partial x_j} + \operatorname{div}_x(bF) = 0,$$

where

$$b(x) = a(F_M(x) + F_{M'}(x)) - x, \quad x \in \mathbb{R}^N;$$

notice that $a = 0$ if $N \geq 2$. Since $F \in H^2(\mathbb{R}^N)$, we have

$$\operatorname{div}_x(bF) \mathbf{1}_\Theta = \operatorname{div}_x(bF_+)$$

almost everywhere. Thus, we obtain

$$-\sum_{1 \leq i, j \leq N} \eta_{i,j} \mathbf{1}_\Theta \frac{\partial^2 F}{\partial x_i \partial x_j} + \operatorname{div}_x(bF_+) = 0.$$

Integrating the above equation on \mathbb{R}^N leads to

$$\int_{\Theta} \sum_{1 \leq i, j \leq N} \eta_{i,j} \frac{\partial^2 F}{\partial x_i \partial x_j} = 0.$$

Let us now perform the change of variables (2.18), which changes the matrix η into identity: setting $\tilde{F}(y) = F(Py)$, and $\tilde{\Theta} := \{\tilde{F} > 0\}$, we infer

$$\int_{\tilde{\Theta}} \Delta_y \tilde{F} = C \int_{\tilde{\Theta}} \sum_{1 \leq i, j \leq N} \eta_{i,j} \frac{\partial^2 F}{\partial x_i \partial x_j} = 0.$$

Moreover, $\tilde{F} \in H^2 \cap W^{2,1}(\mathbb{R}^N)$, and thus Lemma 7 in [1] applies. We deduce that

$$\Delta_y(\tilde{F}\mathbf{1}_\Theta) = \mathbf{1}_\Theta \Delta_y \tilde{F},$$

and thus

$$\sum_{1 \leq i, j \leq N} \eta_{i,j} \mathbf{1}_\Theta \frac{\partial^2 F}{\partial x_i \partial x_j} = \sum_{1 \leq i, j \leq N} \eta_{i,j} \frac{\partial^2 F_+}{\partial x_i \partial x_j}.$$

Eventually, F_+ solves the elliptic equation

$$-\sum_{1 \leq i, j \leq N} \eta_{i,j} \frac{\partial^2 F_+}{\partial x_i \partial x_j} + \operatorname{div}_x(bF_+) = 0,$$

with $b \in L^\infty_{\text{loc}}(\mathbb{R}^N)$. Using either a unique continuation principle or Harnack's inequality (see [13], Theorem 8.20), we infer that if F_+ vanishes at some point x in \mathbb{R}^N , then F_+ is identically zero on \mathbb{R}^N , which is absurd. Hence $F_+(x) > 0$ for all $x \in \mathbb{R}^N$, and thus $\mathbb{R}^N \setminus \Theta = \emptyset$, which means that $F(x) > 0$ for all $x \in \mathbb{R}^N$.

□

Acknowledgements

I wish to thank Adrien Blanchet, Jean Dolbeault, and Michal Kowalczyk, for very fruitful and stimulating discussions.

References

- [1] J. Aguirre, M. Escobedo, and E. Zuazua, *Self-similar solutions of a convection diffusion equation and related semilinear elliptic problems*, Comm. Partial Differential Equations **15** (1990), no. 2, 139–157.
- [2] Grégoire Allaire, *Homogenization and two-scale convergence*, SIAM J. Math. Anal. **23** (1992), no. 6, 1482–1518.
- [3] Alain Bensoussan, Jacques-Louis Lions, and George Papanicolaou, *Asymptotic analysis for periodic structures*, Studies in Mathematics and its Applications, vol. 5, North-Holland Publishing Co., Amsterdam, 1978.
- [4] Adrien Blanchet, Jean Dolbeault, and Michal Kowalczyk, *Travelling fronts in stochastic Stokes' drifts*, Physica A: Statistical Mechanics and its Applications **387** (2008), no. 23, 5741–5751.
- [5] Adrien Blanchet, Jean Dolbeault, and Michal Kowalczyk, *Stochastic Stokes' drift, homogenized functional inequalities, and large time behavior of brownian ratchets*, SIAM Journal on Mathematical Analysis **41** (2009), no. 1, 46–76.

- [6] Haïm Brezis, *Analyse fonctionnelle*, Collection Mathématiques Appliquées pour la Maîtrise., Masson, Paris, 1983, Théorie et applications.
- [7] Anne-Laure Dalibard, *Homogenization of a quasilinear parabolic equation with vanishing viscosity*, J. Math. Pures Appl. (9) **86** (2006), no. 2, 133–154.
- [8] ———, *Long time behaviour of viscous scalar conservation laws*, to be published in the Indiana University Mathematics Journal, 2010.
- [9] Robert Dautray and Jacques-Louis Lions, *Analyse mathématique et calcul numérique pour les sciences et les techniques. Vol. 5*, INSTN: Collection Enseignement, Masson, Paris, 1988, Spectre des opérateurs, With the collaboration of Michel Artola, Michel Cessenat, Jean Michel Combes and Bruno Scheurer, Reprinted from the 1984 edition.
- [10] Marco Di Francesco and Peter A. Markowich, *Entropy dissipation and Wasserstein metric methods for the viscous Burgers' equation: convergence to diffusive waves*, Partial differential equations and inverse problems, Contemp. Math., vol. 362, Amer. Math. Soc., Providence, RI, 2004, pp. 145–165.
- [11] Miguel Escobedo, Juan Luis Vázquez, and Enrike Zuazua, *Asymptotic behaviour and source-type solutions for a diffusion-convection equation*, Arch. Rational Mech. Anal. **124** (1993), no. 1, 43–65.
- [12] Miguel Escobedo and Enrike Zuazua, *Large time behavior for convection-diffusion equations in \mathbf{R}^N* , J. Funct. Anal. **100** (1991), no. 1, 119–161.
- [13] David Gilbarg and Neil S. Trudinger, *Elliptic partial differential equations of second order*, Classics in Mathematics, Springer-Verlag, Berlin, 2001, Reprint of the 1998 edition.
- [14] Thierry Goudon and Frédéric Poupaud, *Homogenization of transport equations: weak mean field approximation*, SIAM J. Math. Anal. **36** (2004/05), no. 3, 856–881 (electronic).
- [15] S. N. Kruzhkov, *First order quasilinear equations in several independent variables*, Math. USSR Sb. (1970), no. 10, 217–243.
- [16] O. A. Ladyženskaja, V. A. Solonnikov, and N. N. Ural'ceva, *Linear and quasilinear equations of parabolic type*, Translated from the Russian by S. Smith. Translations of Mathematical Monographs, Vol. 23, American Mathematical Society, Providence, R.I., 1967.

- [17] J. P. LaSalle, *The stability of dynamical systems*, Society for Industrial and Applied Mathematics, Philadelphia, Pa., 1976, With an appendix: “Limiting equations and stability of nonautonomous ordinary differential equations” by Z. Artstein, Regional Conference Series in Applied Mathematics.
- [18] Philippe Michel, Stéphane Mischler, and Benoît Perthame, *General relative entropy inequality: an illustration on growth models*, J. Math. Pures Appl. (9) **84** (2005), no. 9, 1235–1260.
- [19] Stanley Osher and James Ralston, *L^1 stability of travelling waves with applications to convective porous media flow*, Comm. Pure Appl. Math. **35** (1982), no. 6, 737–749.
- [20] Denis Serre, *Systems of conservation laws. I*, Cambridge University Press, Cambridge, 1999, Hyperbolicity, entropies, shock waves, Translated from the 1996 French original by I. N. Sneddon.
- [21] ———, *L^1 -stability of nonlinear waves in scalar conservation laws*, Evolutionary equations. Vol. I, Handb. Differ. Equ., North-Holland, Amsterdam, 2004, pp. 473–553.
- [22] Enrique Zuazua, *Weakly nonlinear large time behavior in scalar convection-diffusion equations*, Differential Integral Equations **6** (1993), no. 6, 1481–1491.
- [23] ———, *A dynamical system approach to the self-similar large time behavior in scalar convection-diffusion equations*, J. Differential Equations **108** (1994), no. 1, 1–35.