

“Good Lie Brackets” for Control Affine Systems

Andrei Agrachev

We consider a control affine system

$$\dot{q} = f_0(q) + \sum_{i=1}^k u_i f_i(q), \quad q \in M, \quad u_i \in \mathbb{R}. \quad (1)$$

Here f_j , $j = 0, \dots, k$, are smooth vector fields on the manifold M . We assume that these fields and their iterated Lie brackets have at most linear growth for a complete Riemannian metric on M ; in particular, all these vector fields are complete.

We study controllability issues and do not impose constraints on the control parameters u_i to simplify the task. We however do not want to abuse this freedom too much and use only uniformly bounded trajectories to attain points of a bounded domain.

Example 1:

$$\begin{cases} \dot{x} = \varphi(x, y) + u \\ \dot{y} = \psi(x, y), \end{cases} \quad u, x \in \mathbb{R}, y \in \mathbb{R}^m,$$

where ψ is a degree n vector-polynomial w.r.t. x .

Extended system:

$$\begin{cases} \dot{x} = u \\ \dot{y} = \sum_{i=0}^n \frac{u_i}{i!} \frac{\partial^i \psi}{\partial x^i}(x, y), \end{cases} \quad u_0 = 1, \quad \sum_{i,j=0}^{\frac{n}{2}} u_{i+j} \xi_i \xi_j \geq 0, \quad \forall \xi. .$$

Any diffeomorphism produced by the extended system in time t can be C^∞ -approximated by a diffeomorphism produced by the original system in the same time.

Example 2:

Vector fields f_0, f_1, \dots, f_k generate a step 3 nilpotent Lie algebra.

Extended system:

$$\dot{q} = f_0 + \sum_{i=1}^k \left(u_i f_i + u_{i0} [f_i, f_0] + \frac{u_{ii}}{2} [f_i, [f_i, f_0]] \right) + \sum_{1 \leq i < j \leq k} \left(u_{ij} [f_i, [f_j, f_0]] + v_{ij} [[f_i, f_j], f_0] + w_{ij} [f_i, f_j] \right),$$

where $\sum_{i,j=0}^k u_{ij} \xi_i \xi_j \geq 0, \forall \xi.$, $u_{00} = 1$, $u_{ij} = u_{ji}$, and $v_{ij}, w_{ij} \in \mathbb{R}$ are free.

We use chronological notations. Let f_τ be a time-varying vector field, then

$$\overrightarrow{\exp} \int_0^t f_\tau d\tau : M \rightarrow M, \quad t \in \mathbb{R},$$

is the flow generated by the equation $\dot{q} = f_\tau(q)$ and $t \mapsto q_0 \circ \overrightarrow{\exp} \int_0^t f_\tau d\tau$ is a trajectory of this flow. Moreover,

$$\left(\overrightarrow{\exp} \int_0^t f_\tau d\tau \right)_*^{-1} g = \overrightarrow{\exp} \int_0^t \text{ad} f_\tau d\tau g,$$

where $(\text{ad} f)g = [f, g]$.

Let us start with small time control. To this end we take a sample control $u(t) = (u_1(t), \dots, u_k(t))$, $t \in [0, 2]$, a small parameter $\varepsilon > 0$ and cook a re-scaled control $\frac{1}{\varepsilon}u(\frac{t}{\varepsilon})$. Now re-scaling time in the equation

$$\dot{q} = f_0(q) + \frac{1}{\varepsilon} \sum_{i=1}^k u_i\left(\frac{t}{\varepsilon}\right) f_i(q),$$

we obtain the system

$$\frac{dq}{d\tau} = \varepsilon f_0(q) + \sum_{i=1}^k u_i(\tau) f_i(q),$$

where $\tau = \frac{t}{\varepsilon}$.

We set $f_u = \sum_{i=1}^k u_i f_i(q)$; then

$$q(\varepsilon) = q_0 \circ \overrightarrow{\exp} \int_0^2 \varepsilon f_0 + f_u d\tau =$$

$$q_0 \circ \overrightarrow{\exp} \int_0^2 \varepsilon \overrightarrow{\exp} \int_0^\tau \text{ad}_{f_u(\theta)} d\theta f_0 d\tau \circ \overrightarrow{\exp} \int_0^2 f_u d\tau.$$

Now assume that $u(t)$ has a form:

$$u(t) = \begin{cases} \frac{v(\tau)}{2}, & 0 \leq \tau \leq 1; \\ -\frac{v(1-\tau)}{2}, & 1 < \tau \leq 2; \end{cases}$$

then $\overrightarrow{\exp} \int_1^2 f_{u(\tau)} d\tau = \left(\overrightarrow{\exp} \int_0^1 f_{u(\tau)} d\tau \right)^{-1}$ and we get

$$q(\varepsilon) = q_0 \circ \overrightarrow{\exp} \int_0^2 \varepsilon \overrightarrow{\exp} \int_0^\tau \text{ad} f_{u(\theta)} d\theta f_0 d\tau =$$

$$q_0 \circ e^{\varepsilon \left(\int_0^2 \overrightarrow{\exp} \int_0^\tau \text{ad} f_{u(\theta)} d\theta f_0 d\tau + O(\varepsilon) \right)}.$$

Moreover, $\overrightarrow{\exp} \int_0^{1+s} f_{u(\tau)} d\tau = \overrightarrow{\exp} \int_0^{1-s} f_{u(\tau)} d\tau$, $0 \leq s \leq 1$, and we obtain:

$$\int_0^2 \overrightarrow{\exp} \int_0^t \text{ad} f_{u(\tau)} d\tau f_0 dt = \int_0^1 \overrightarrow{\exp} \int_0^t \text{ad} f_{v(\tau)} d\tau f_0 dt,$$

where $v(\cdot)$ is any function from $L_1([0, 1]; \mathbb{R}^k)$. We set:

$$\mathcal{V} = \left\{ \int_0^1 \overrightarrow{\exp} \int_0^t \text{ad} f_{v(\tau)} d\tau f_0 dt : v \in L_1([0, 1]; \mathbb{R}^k) \right\}$$

and we can re-write:

$$q(\varepsilon) = q_0 \circ e^{\varepsilon(V + O(\varepsilon))}, \quad V \in \mathcal{V}.$$

Let $t \mapsto V_t \in \mathcal{V}$ be a piecewise constant family of vector fields, $V_t = V_i, \forall t \in (\frac{i-1}{\varepsilon}, \frac{i}{\varepsilon}]$. We repeat our procedure n times with an appropriate choice of $v(\cdot)$ for any segment $[\frac{i-1}{\varepsilon}, \frac{i}{\varepsilon}]$ and obtain:

$$q(n\varepsilon) = q_0 \circ e^{\varepsilon(V_\varepsilon + O(\varepsilon))} \circ \dots \circ e^{\varepsilon(V_{n\varepsilon} + O(\varepsilon))} = q_0 \circ \overrightarrow{\exp} \int_0^{n\varepsilon} V_t + O(\varepsilon) dt.$$

It follows that we can arbitrarily well approximate any diffeomorphism of the form $\overrightarrow{\exp} \int_0^t V_\tau d\tau$ by the value at the time moment t of a flow generated by control system (1). Moreover, $t \mapsto V_t$ maybe of class L_1 , it is not obliged to be piecewise constant.

Of course, we can also approximate any diffeomorphism of the form

$$\overrightarrow{\exp} \int_0^t V_\tau d\tau \circ \overrightarrow{\exp} \int_0^1 f_{u(t)} dt, \quad u(t) \in \mathbb{R}^k, \quad V_\tau \in \mathcal{V}.$$

The second term in the product is generated by the control linear system (without drift).

Assume for the moment that the fields f_1, \dots, f_k generate a finite-dimensional Lie algebra, $\text{Lie}\{f_1, \dots, f_k\} = L$, $\dim L < \infty$. Let $\mathcal{L} \subset \text{Diff}M$ be the Lie group generated by L ; it is a finite-dimensional Lie subgroup of $\text{Diff}M$. Moreover,

$$\mathcal{L} = \left\{ \overrightarrow{\exp} \int_0^1 f_{u(t)} dt : u(\cdot) \in L_1([0, 1]; \mathbb{R}^k) \right\}.$$

Proposition 1. $\bar{\mathcal{V}} = \overline{\text{conv}\{p_*f_0 : p \in \mathcal{L}\}}$.

Proof. Recall that $p_* = \text{Ad}p^{-1}$ and $\text{Ad}\overrightarrow{\exp} \int_0^t f_\tau d\tau = \overrightarrow{\exp} \int_0^t \text{ad} f_\tau d\tau$.

A convex combination of the vector fields p_*f_0 , $p \in \mathcal{L}$, can be written as $\int_0^1 \text{Ad}p_t f_0 dt$, where $t \mapsto p_t$ is a piecewise constant family of diffeomorphisms from \mathcal{L} . We are interested in the closure of the convexification and we may substitute piecewise constant families by smooth families or by any other class of families that is dense in $L_1([0, 1]; \mathcal{L})$. The result now follows from the fact that any starting from I curve in \mathcal{L} can be uniformly approximated by the curve of the form $p_t = \overrightarrow{\exp} \int_0^t f_{u(\tau)} d\tau$. \square

It follows that attainable sets of the system

$$\dot{q} = U(q), \quad U \in \text{conv}\{p_*f_0 : p \in \mathcal{L}\} \quad (2)$$

are contained in the closure of the attainable sets of system (1). Now if $\text{conv}\{p_*f_0 : p \in \mathcal{L}\}$ is an affine subspace or it contains an affine subspace of vector fields than we can repeat the whole procedure and obtain more available vector fields in the right hand side.

Of course, Proposition 1 can be extended to many interesting families of vector fields which generate infinite dimensional Lie algebras but we do not plan to do these functional analytic exercises in the current paper. We are mainly interested in the structural questions.

Roughly speaking, the construction provides an extension of the original affine subspace of admissible vector fields to a convex set in the closure of the Lie subalgebra generated by this affine subspace. The system can be moved in the direction of any field from this convex set that is built of Lie bracket polynomials and series of the original fields f_j . Moreover, the structure of these polynomials and series does not depend on the choice of the original fields, the fields f_j serve just as variables.

In other words, we may speak about a universal convex set in the closure of the free Lie algebra with generators a_0, a_1, \dots, a_k . This is the set of “good” combinations of brackets and any control affine system can be moved in the direction of these combinations.

Let $\text{Ass}(a_0, \dots, a_k)$ be the free associated algebra over \mathbb{R} , its elements are linear combinations of words in the alphabet $\{a_0, \dots, a_k\}$. Then $\text{Ass}(a_0, \dots, a_k) = \bigoplus_{n=0}^{\infty} A_n$, where the space A_n consists of linear combinations of words with n letters and $A_0 = \mathbb{R}$ corresponds to the empty word. The *closure* of $\text{Ass}(a_0, \dots, a_k)$ is the algebra of formal series

$$\mathfrak{A} = \left\{ \sum_{n=0}^{\infty} x_n : x_n \in A_n \right\}$$

endowed with topology of the term-wise convergence; we write $\mathfrak{A} = \overline{\text{Ass}(a_0, \dots, a_k)}$

The universal control affine system with k -dimensional control is the system

$$\dot{x} = x \left(a_0 + \sum_{i=1}^k u_i a_i \right), \quad x \in \mathfrak{A}, \quad u_i \in \mathbb{R}. \quad (3)$$

Given control $u(\cdot) \in L_1([0, t]; \mathbb{R}^k)$ and initial condition $x(0)$, we can explicitly write the unique solution of (3) that is a curve in \mathfrak{A} whose homogeneous components are absolutely continuous vector functions.

Let W be the set of words in the alphabet $\{a_0, \dots, a_k\}$ and

$$\Delta^n(t) = \{(\tau_1, \dots, \tau_n) : 0 \leq \tau_n \leq \dots \leq \tau_1 \leq t\}$$

be the n -dimensional simplex. Given a word $w = a_{i_n} \dots a_{i_1}$, we set

$$S_u^w(t) = \int \dots \int_{\Delta^n(t)} u_{i_n}(\tau_n) \dots u_{i_1}(\tau_1) d\tau_1 \dots d\tau_n,$$

where $u_0(t) \equiv 1$. Solutions of (3) have a form:

$$x(t) = x(0) \sum_{w \in W} S_u^w(t) w.$$

We keep using chronological notations while working in \mathfrak{A} with the composition “ \circ ” substituted by the product in \mathfrak{A} . In what follows, we assume that $x(0) = 1$. More notations:

$$L = \text{Lie}(a_1, \dots, a_k) \subset \text{Ass}(a_1, \dots, a_k), \quad \mathcal{L} = \{e^V : V \in \bar{L}\}, \quad a_u = \sum_{i=1}^k u_i a_i.$$

Next statement easily follows from basic facts of Lie theory.

Theorem 1.

$$\mathcal{L} = \overline{\left\{ \overrightarrow{\exp} \int_0^1 a_{v(t)} dt : v(\cdot) \in L_1([0, 1]; \mathbb{R}^k) \right\}}.$$

Moreover,

$$\overline{\text{conv}\{(Adx)a_0 : x \in \mathcal{L}\}} = \overline{\left\{ \int_0^1 \overrightarrow{\exp} \int_0^t \text{ad}_{a_{v(\tau)}} d\tau a_0 dt : v \in L_1([0, 1]; \mathbb{R}^k) \right\}}.$$

□

We may translate all the computations to the universal setting by the substitution of f_j with a_j and we obtain:

Theorem 2. *For any $t > 0$, the product of \mathcal{L} and the attainable set of the system*

$$\dot{x} = xV, \quad V \in \text{conv}\{(\text{Ad}z)a_0 : z \in \mathcal{L}\}$$

at t is contained in the closure of the attainable set of system (3) at t .

□

Let V_α , $\alpha = 1, 2, \dots$, be a linearly ordered homogeneous additive basis of L . It may be a Hall basis but this is not necessary. It is easy to see that

$$\mathcal{L} = \left\{ \prod_{\alpha=1}^{\infty} e^{v_\alpha V_\alpha} : v_\alpha \in \mathbb{R} \right\}. \quad (4)$$

This is what people call “the 2nd type coordinates” for the Lie group, while the presentation $\mathcal{L} = e^{\bar{L}}$ is the “1st type coordinates.

We have:

$$\prod_{\alpha=1}^{\infty} e^{v_\alpha V_\alpha} = 1 + \sum_{\substack{\alpha_1 \leq \dots \leq \alpha_m \\ i_1, \dots, i_m > 0}} \frac{v_{\alpha_1}^{i_1} \dots v_{\alpha_m}^{i_m}}{i_1! \dots i_m!} V_{\alpha_1}^{i_1} \dots V_{\alpha_m}^{i_m}.$$

According to the Poincare–Birkhoff–Witt theorem, the elements

$$V_{\alpha_1}^{i_1} \cdots V_{\alpha_m}^{i_m}, \quad \alpha_1 \leq \cdots \leq \alpha_m, \quad i_1, \dots, i_m > 0, \quad m \geq 0, \quad (5)$$

form an additive basis of $\text{Ass}\{a_1, \dots, a_k\}$, hence

$$\overline{\text{span}\mathcal{L}} = \overline{\text{Ass}(a_1, \dots, a_k)}.$$

Next statement reduces the study of $\text{conv}\{(\text{Ad}z)a_0 : z \in \mathcal{L}\} \subset \overline{L}$ to the study of $\text{conv}\mathcal{L} \subset \overline{\text{Ass}(a_1, \dots, a_k)}$.

Proposition 2. *Linear map $\text{Ad}_0 : \text{Ass}(a_1, \dots, a_k) \rightarrow \text{Lie}(a_0, a_1, \dots, a_k)$ defined by its action on the basis:*

$$\text{Ad}_0 \left(V_{\alpha_1}^{i_1} \cdots V_{\alpha_m}^{i_m} \right) = (\text{ad}V_{\alpha_1})^{i_1} \cdots (\text{ad}V_{\alpha_m})^{i_m} a_0$$

is injective.

Let t_α , $\alpha = 1, 2, \dots$, be coordinates on L induced by the basis V_α . In other words, $t_\alpha \in L^*$, $\langle t_\alpha, V_\beta \rangle = \delta_{\alpha, \beta}$. The basis provides the identification of \bar{L} and L^* . According to this identification, a series $\sum_\alpha v_\alpha V_\alpha$ is identified with the linear function $\sum_\alpha v_\alpha t_\alpha$ on L .

Moreover, the monomials $t_{\alpha_1}^{i_1} \cdots t_{\alpha_m}^{i_m}$ are coordinates on the vector space $\text{Ass}(a_1, \dots, a_k)$. A monomial $t_{\alpha_1}^{i_1} \cdots t_{\alpha_m}^{i_m}$ treated as a linear form on $\text{Ass}(a_1, \dots, a_k)$ annihilates all elements of the basis (5) except of $V_{\alpha_1}^{i_1} \cdots V_{\alpha_m}^{i_m}$ and $\langle t_{\alpha_1}^{i_1} \cdots t_{\alpha_m}^{i_m}, V_{\alpha_1}^{i_1} \cdots V_{\alpha_m}^{i_m} \rangle = 1$.

The basis (5) provides the identification of $\overline{\text{Ass}(a_1, \dots, a_k)}$ with $\text{Ass}(a_1, \dots, a_k)^*$ and eventually with the space of formal power series on the variables t_α , $\alpha = 1, 2, \dots$. Let \mathcal{S} the space of formal power series and

$$\nu : \overline{\text{Ass}(a_1, \dots, a_k)} \rightarrow \mathcal{S}$$

be the continuous isomorphism of vector spaces that realizes the mentioned identification,

$$\nu : V_{\alpha_1}^{i_1} \cdots V_{\alpha_m}^{i_m} \mapsto t_{\alpha_1}^{i_1} \cdots t_{\alpha_m}^{i_m},$$

where $\alpha_1 \leq \cdots \leq \alpha_m$ as in (5). Linear map ν depends on the choice of the basis V_α and it is not a homomorphism of the algebras.

Definition 1. We say that a nonzero function $\varphi : L \rightarrow \mathbb{R}$ is exponential if the restriction of φ to any finite-dimensional subspace of L is continuous and

$$\varphi(z_1 + z_2) = \varphi(z_1)\varphi(z_2), \quad \forall z_1, z_2 \in L.$$

It is easy to see that, written in the coordinates, exponential functions are exactly functions of the form $\varphi(t) = e^{\langle v, t \rangle}$, where

$$v = \{v_\alpha\}_{\alpha=1}^{\infty}, \quad t = \{t_\alpha\}_{\alpha=1}^{\infty}, \quad \langle v, t \rangle = \sum_{\alpha=1}^{\infty} v_\alpha t_\alpha.$$

Recall that an element of L has only a finite number of nonzero coordinates t_α .

The space of exponential functions is denoted by \mathcal{E} . The identification of the exponential function with the exponential series gives the inclusion $\mathcal{E} \subset \mathcal{S}$.

Proposition 3. $\nu(\mathcal{L}) = \mathcal{E}$.

Proof. Indeed,

$$\begin{aligned} \nu \left(\prod_{\alpha=1}^{\infty} e^{v_{\alpha} V_{\alpha}} \right) &= 1 + \sum_{\substack{\alpha_1 \leq \dots \leq \alpha_m \\ i_1, \dots, i_m > 0}} \frac{v_{\alpha_1}^{i_1} \cdots v_{\alpha_m}^{i_m}}{i_1! \cdots i_m!} \nu \left(V_{\alpha_1}^{i_1} \cdots V_{\alpha_m}^{i_m} \right) = \\ &= 1 + \sum_{\substack{\alpha_1 \leq \dots \leq \alpha_m \\ i_1, \dots, i_m > 0}} \frac{v_{\alpha_1}^{i_1} \cdots v_{\alpha_m}^{i_m}}{i_1! \cdots i_m!} t_{\alpha_1}^{i_1} \cdots t_{\alpha_m}^{i_m} = \prod_{\alpha=1}^{\infty} e^{v_{\alpha} t_{\alpha}} = e^{\langle v, t \rangle}. \quad \square \end{aligned}$$

We see that $\nu(\mathcal{L})$ does not depend on the choice of the basis V_{α} , unlikely the isomorphism ν .

The space of formal series \mathcal{S} is the adjoint space to the space of (finite) linear combinations of partial differentials $\frac{\partial^{i_1}}{\partial t_{\alpha_1}^{i_1}} \cdots \frac{\partial^{i_m}}{\partial t_{\alpha_m}^{i_m}} \Big|_{t=0}$, where the pairing of the differential and the series is just the action of the differential on the series.

We set:

$$\mathcal{D} = \text{span} \left\{ \frac{\partial^{i_1}}{\partial t_{\alpha_1}^{i_1}} \cdots \frac{\partial^{i_m}}{\partial t_{\alpha_m}^{i_m}} \Big|_{t=0} : \alpha_1 \leq \cdots \leq \alpha_m, i_1, \dots, i_m > 0, m \geq 0 \right\},$$

$\mathcal{S} = \mathcal{D}^*$. To any $\varphi \in \mathcal{S}$ we associate a quadratic form Q_φ by the following formula:

$$Q_\varphi(\eta) = \eta_t \eta_s \varphi(t + s), \quad \eta \in \mathcal{D},$$

where η_t differentiates with respect to t and η_s differentiates with respect to s .

Theorem 3. Let $\varphi \in \text{span}\mathcal{E}$, $\varphi(0) = 1$. Quadratic form Q_φ is nonnegative if and only if $\varphi \in \text{conv}\mathcal{E}$.

Proof. If $\varphi \in \mathcal{E}$, then $\varphi(t+s) = \varphi(t)\varphi(s)$ and

$$Q_\varphi(\eta) = \eta_t \eta_s \varphi(t+s) = (\eta\varphi)^2 \geq 0.$$

Let $\varphi = \sum_{i=1}^n c_i \varphi_i$. We may assume that $\varphi_i(t) = e^{\langle w_i, t \rangle}$, $i = 1, \dots, n$, where w_1, \dots, w_n are mutually distinct. Of course, there exists $m > 0$ such that the truncation of these infinite vectors to \mathbb{R}^m are also mutually distinct.

We have: $Q_\varphi(\eta) = \sum_{i=1}^n c_i (\eta\varphi_i)^2$. If all c_i are nonnegative, then $Q_\varphi \geq 0$. Assume that $c_{i_0} < 0$. On the other hand, Taylor polynomials of order n of $\varphi_1, \dots, \varphi_n$ at 0 are linearly independent. Hence there exists $\eta_0 \in \mathcal{D}$ such that $\eta_0\varphi_{i_0} = 1$, $\eta_0\varphi_i = 0$, $\forall i \neq i_0$, and $Q_\varphi(\eta_0) = c_{i_0}$. \square

Free Lie algebra is too big. It is more practical and sufficient for many purposes to consider its finite-dimensional nilpotent truncations.

In what follows, we use multi-indices. Given a nonnegative integer m , \mathbb{Z}_+^m is the set of m -dimensional vectors with nonnegative integral coordinates. We extend m -dimensional vectors by zeros and assume that $\mathbb{Z}_+^m \subset \mathbb{Z}_+^{m'}$ if $m \leq m'$; then $\mathbb{Z}_+^\infty = \bigcup_{m \geq 0} \mathbb{Z}_+^m$ is a set of infinite vectors with a finite number nonzero coordinates. If $i = (i_1, \dots, i_m) \in \mathbb{Z}_+^m$, then:

$$t^i = t_1^{i_1} \cdots t_m^{i_m}, \quad |i| = \sum_{j=1}^m i_j, \quad \varphi_0^{(i)} = \left. \frac{\partial^{|i|} \varphi}{\partial t_1^{i_1} \cdots \partial t_m^{i_m}} \right|_{t=0}.$$

Let $I \subset \mathbb{Z}_+^\infty$ be a finite subset, we set:

$$\mathcal{P}(I) = \left\{ \sum_{i \in I} c_i t^i : c_i \in \mathbb{R} \right\},$$

a $\#I$ -dimensional space of polynomials. We denote by $\Pi_I : \mathcal{S} \rightarrow \mathcal{P}(I)$ the continuous linear projector defined by the rule:

$$\Pi_I(t^i) = \begin{cases} t^i, & \text{if } i \in I; \\ 0, & \text{if } i \in \mathbb{Z}_+^\infty \setminus I. \end{cases}$$

Lemma 1. $\text{span} \Pi_I(\mathcal{E}) = \mathcal{P}(I)$, for any $I \subset \mathbb{Z}_+^\infty$ such that $\#I < \infty$.

Let $C \subset \mathbb{R}_+^m$ be a convex compact subset; we set: $I_C = C \cap \mathbb{Z}_+^m$.
Theorem 4. *Let C be a convex compact subset of \mathbb{R}_+^m and $0 \in C$.
If $\phi \in \mathcal{P}(I_C)$ belongs to $\overline{\Pi_{I_C}(\text{conv}\mathcal{E})}$, then*

$$\phi(0) = 1, \quad \sum_{i,j \in I_{\frac{1}{2}C}} \phi_0^{(i+j)} \xi_i \xi_j \geq 0, \quad \forall \xi_i \in \mathbb{R}, i \in I_{\frac{1}{2}C}. \quad (6)$$

Moreover, if $m = 1$ or $|i| \leq 2, \forall i \in I_C$, then condition (6) is sufficient for ϕ to belong to $\overline{\Pi_{I_C}(\text{conv}\mathcal{E})}$.